

Elimination Reaction-Based Benzimidazole Probe for Cysteine Detection and Its Application in Serum Sample Analysis

In-ho Song ^{1,†}, Gyu Seong Yeom ^{1,†}, Anil Kuwar ² and Satish Balasaheb Nimse ^{1,*}

¹ Institute of Applied Chemistry and Department of Chemistry, Hallym University, Chuncheon 24252, Korea; songin-ho@hallym.ac.kr (I.-h.S.); m21019@hallym.ac.kr (G.S.Y.)

² School of Chemical Sciences, KBC-North Maharashtra University, Jalgaon 425001, India; kuwaras@gmail.com

* Correspondence: authors: satish_nimse@hallym.ac.kr

† These authors contributed equally to this work.

Supplementary Information




Figure S1. ¹H-NMR spectrum of compound 3.




Figure S2. ^{13}C NMR spectrum of compound 3.




Figure S3. Mass spectrum of compound 3.




Figure S4. ¹H-NMR spectrum of A-B.




Figure S5. ¹³C NMR spectrum of A-B.




Figure S6. Mass spectrum of A-B.



Figure S7. Image taken under UV lamp depicting the selectivity of A-B for Cys among other amino acids. The addition of Cys (50 μ M) to the solution of A-B (10 μ M) results in fluorescence turn-on effect.