



Supplementary File

Table S1. Chemical constituents of *Helichrysum italicum* (Roth) G. Don essential oil.

No.	Compound	Area percentage (%)
1.	α-pinene	5.6
2.	α -fenchene	0.1
3.	β -pinene	0.3
4.	limonene	2.0
5.	1,8-cineol	0.5
6.	γ -terpinene	0.5
7.	linalool	0.4
8.	2-methylbutyl-2-methylbutanoate	0.3
9.	nerol	0.3
10.	neryl acetate	10.8
11.	α -copaene	2.6
12.	italicene	3.1
13.	<i>cis</i> - α -bergamotene	0.9
14.	<i>trans</i>-β-caryophyllene	8.4
15.	<i>trans</i> - α -bergamotene	0.9
16.	4,6,9-trimethyldec-8-ene-3,5-dione	3.3
17.	neryl propionate	2.0
18.	γ -selinene	2.4
19.	γ-curcumene	18.1
20.	ar-curcumene	5.8
21.	β-selinene	9.9
22.	2,4,6,9-tetramethyldec-8-ene-3,5-dione	2.4
23.	α-selinene	6.9
24.	α -muurolene	0.5
25.	γ -cadinene	0.4
26.	δ -cadinene	1.7
27.	3,5,7,10-tetramethylundec-9-ene-4,6-dione	0.9

Bold means the highest percentage of the essential oil compounds.