antibiotics

MDPI

Correction

Correction: Jensen et al. Prior Antibiotic Use Increases Risk of Urinary Tract Infections Caused by Resistant *Escherichia coli* among Elderly in Primary Care: A Case-Control Study. *Antibiotics* 2022, *11*, 1382

Maria L. V. Jensen [1,2,*], Volkert Siersma [2], Lillian M. Søes [3], Dagny Nicolaisdottir [2], Lars Bjerrum [2] and Barbara J. Holzknecht [1,4]

[1] Department of Clinical Microbiology, Copenhagen University Hospital—Herlev and Gentofte, 2730 Herlev, Denmark
[2] The Research Unit for General Practice and Section of General Practice, Department of Public Health, University of Copenhagen, 1014 Copenhagen, Denmark
[3] Department of Clinical Microbiology, Copenhagen University Hospital—Hvidovre and Amager, 2650 Hvidovre, Denmark
[4] Department of Clinical Medicine, University of Copenhagen, 2200 Copenhagen, Denmark
* Correspondence: mlvj@sund.ku.dk; Tel.: + 45-27-793-223

Error in Figure/Table

In the original publication [1], there was a mistake in Table 2 "Adjusted analyses of total exposure to antibiotics (number of prescriptions and DDD) and time since last prescription, and OR of resistant *E. coli*" as published. The last row of the table is wrongly titled, and the presented results in this row are not correct. The corrected Table 2 appears below. The authors state that the scientific conclusions are unaffected. This correction was approved by the Academic Editor. The original publication has also been updated.

Table 2. Adjusted analyses of total exposure to antibiotics (number of prescriptions and DDD) and time since last prescription, and OR of resistant *E. coli*.

		Mecillinam		Trimethoprim		Nitrofurantoin		Multi-Resistance	
		OR *	CI95% **	OR	CI95%	OR	CI95%	OR	CI95%
Number of prescriptions	0	Ref		Ref		Ref		Ref	
	1	1.32	(1.15;1.53)	1.40	(1.32;1.50)	1.25	(0.97;1.61)	1.35	(0.59;3.08)
	2	1.82	(1.59;2.07)	1.66	(1.56;1.77)	2.10	(1.69;2.60)	2.86	(1.45;5.63)
	≥3	2.47	(2.18;2.80)	3.11	(2.92;3.31)	4.53	(3.76;5.46)	11.53	(6.37;20.85)
Number of DDD ***	0	Ref		Ref		Ref		Ref	
	>0–33.3 percentile	1.56	(1.39;1.76)	1.54	(1.46;1.63)	1.84	(1.51;2.23)	1.34	(0.59;3.05)
	>33.3–66.6 percentile	1.88	(1.63;2.18)	2.03	(1.89;2.17)	2.59	(2.07;3.24)	2.29	(1.07;4.95)
	>66.6 percentile	2.60	(2.26;2.99)	3.22	(2.99;3.46)	4.49	(3.65;5.51)	10.21	(5.78;18.02)
Time since last prescription	No exposure	Ref		Ref		Ref		Ref	
	8–30 days	2.13	(1.91;2.38)	2.20	(2.09;2.32)	2.83	(2.38;3.37)	5.81	(3.37;10.03)
	31–60 days	1.53	(1.32;1.78)	1.71	(1.60;1.83)	2.51	(2.00;3.14)	3.72	(1.93;7.16)
	61–90 days	1.51	(1.27;1.79)	1.49	(1.38;1.62)	1.81	(1.37;2.40)	2.15	(0.85;5.40)

* OR = Odds Ratio. ** CI95% = 95% Confidence interval. *** For the different resistance patterns, the 33.3/66.6 percentiles were as follows: mecillinam 10/21.88, trimethoprim 10/21.88, nitrofurantoin 10.5/22.00, multi-resistance 10/18 DDD.

Citation: Jensen, M.L.V.; Siersma, V.; Søes, L.M.; Nicolaisdottir, D.; Bjerrum, L.; Holzknecht, B.J. Correction: Jensen et al. Prior Antibiotic Use Increases Risk of Urinary Tract Infections Caused by Resistant *Escherichia coli* among Elderly in Primary Care: A Case-Control Study. *Antibiotics* 2022, *11*, 1382. *Antibiotics* **2023**, *12*, 386. https://doi.org/10.3390/antibiotics12020386

Received: 12 January 2023
Accepted: 7 February 2023
Published: 14 February 2023

Reference

1. Jensen, M.L.V.; Siersma, V.; Søes, L.M.; Nicolaisdottir, D.; Bjerrum, L.; Holzknecht, B.J. Prior Antibiotic Use Increases Risk of Urinary Tract Infections Caused by Resistant *Escherichia coli* among Elderly in Primary Care: A Case-Control Study. *Antibiotics* **2022**, *11*, 1382. [CrossRef] [PubMed]