

## Supplementary Materials

**Table S1:** Individual MICs of the antibiotics alone in MAC clinical isolates.

Antibiotic	MIC ( $\mu\text{g/mL}$ )							
	<i>M. intracellulare</i>			<i>M. avium</i>				
	<i>M. intracellulare</i> 1	<i>M. intracellulare</i> 2	<i>M. intracellulare</i> 3	<i>M. intracellulare</i> 4	<i>M. avium</i> 1	<i>M. avium</i> 2	<i>M. avium</i> 3	<i>M. avium</i> 4
CLA	8	8	2	2	4	2	4	<0.25
MOX	16	2	2	4	4	4	4	<0.25
BEDA	2	2	0.5	0.5	0.5	0.5	0.5	1
CLO	16	4	16	16	4	2	4	4
RB	0.25	0.25	0.25	0.5	<0.25	4	0.25	1
RF	0.5	0.25	0.5	0.5	<0.25	2	0.25	<0.25
EB	32	32	2	2	8	8	8	8

MIC: minimum inhibitory concentration ( $\mu\text{g/mL}$ ), MAC: *Mycobacterium avium* complex, CLA: clarithromycin, MOX: moxifloxacin, BED: bedaquiline, CLO: clofazimine, RB: rifabutin, RIF: rifampicin, EMB: ethambutol