

Supplementary Materials: Dermal Targeting Delivery of Terbinafine Hydrochloride Using Novel Multi-Ethosomes: A New Approach to Fungal Infection Treatment

Lijun Zhang ^{1,†}, Xue Li ^{2,†}, Shun Yao Zhu ¹, Ting Zhang ¹, Aikebaier Maimaiti ¹, Meihong Ding ¹ and Senlin Shi ^{1,*}

¹ College of Pharmaceutical Science, Zhejiang Chinese Medical University, Hangzhou 310053, China; lijunzhang@zcmu.edu.cn (L.Z.); zsy@zcmu.edu.cn (S.Z.); zhangting55@163.com (T.Z.); akpar15@163.com (A.M.); dingmeihong0125@163.com (M.D.)

² Institut des Sciences Moléculaires d'Orsay, CNRS, Université Paris-Saclay, 91405 Orsay, France; xue.li@universite-paris-saclay.fr

* Correspondence: pjstone@zcmu.edu.cn; Tel.: +86-0571-61768135

† These authors contributed equally to this work.

1. TH Solubility Test




Figure S1. The TH solubility in different mediums at 37 ± 0.5 °C. Data are shown as the mean \pm SD ($n = 3$).

2. ATR-FTIR Investigation




Figure S2. ATR-FTIR investigation: Amide I (A) and amide II (B) wavenumbers of the rabbit skin treated with different formulations.



© 2019 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).