

# Supplementary Materials: Layered Double Hydroxide Clusters as Precursors of Novel Multifunctional Layers: A Bottom-Up Approach

Cristina S. Neves <sup>1</sup>, Alexandre C. Bastos <sup>1</sup>, Andrei N. Salak <sup>1</sup>, Maksim Strykevich <sup>1</sup>, Daisy Rocha <sup>2</sup>, Mikhail L. Zheludkevich <sup>3</sup>, Angela Cunha <sup>2</sup>, Adelaide Almeida <sup>2</sup>, João Tedim <sup>1,\*</sup> and Mário G.S. Ferreira <sup>1</sup>

<sup>1</sup> Department of Materials Engineering and Ceramics, CICECO, University of Aveiro, 3810-193 Aveiro, Portugal; cristina.neves@ua.pt (C.S.N.); acbastos@ua.pt (A.C.B.); salak@ua.pt (A.N.S.); mstrykevich@ua.pt (M.S.); mgferreira@ua.pt (M.G.S.F.)

<sup>2</sup> Department of Biology, CESAM, University of Aveiro, 3810-193 Aveiro, Portugal; deisyrocha@ua.pt (D.R.); acunha@ua.pt (A.C.); aalmeida@ua.pt (A.A.)

<sup>3</sup> MagIC – Magnesium Innovation Centre, Helmholtz-Zentrum Geesthacht, Max-Planck Straße 1, 21502 Geesthacht, Germany; Mikhail.Zheludkevich@hzg.de

\* Correspondence: joao.tedim@ua.pt




Figure S1. XPS spectra of SR-LDH-MBT after functionalization with HTMS.