

Article

The Effect of Calcium Ions on Resting Membrane Potential

Elizabeth R. Elliott  and Robin L. Cooper * 

Department of Biology, University of Kentucky, Lexington, KY 40506, USA; erel222@uky.edu

* Correspondence: rlcoop1@email.uky.edu

Simple Summary: The factors that influence electrical potential across the cell membrane are important because they affect cellular properties that, in turn, allow tissues, organs, and whole organism to function; as such, it is important to understand them. The distribution and movement of certain ions (i.e., Na^+ , K^+ , Ca^{2+}) determine this electrical potential as they move across the membrane through channels and transport processes, some of which are impeded by the presence of other ions. As a particularly apt example: free Ca^{2+} ions are known to block open Na^+ channels. This study illustrated that raising the Ca^{2+} ion concentration on one side of the membrane led to a larger membrane potential, and this is suggested herein to be caused by blockage of a leaky Na^+ channel. If more such channels are present in the membrane, the effects of altering Ca^{2+} concentration have less impact.

Abstract: Regulating membrane potential is key to cellular function. For many animal cells, resting membrane potential is predominantly driven by a family of K2P (two-pore domain) potassium channels. These channels are commonly referred to as leak channels, as their presence results in the membrane being permeable to K^+ ions. These channels, along with various pumps and exchangers, keep the cell resting membrane potential (R_p) relatively close to potassium's equilibrium potential (E_K); however, in many cells, the resting membrane potential is more depolarized than the E_K due to a small Na^+ ion leak. Raising $[\text{Ca}^{2+}]_O$ (extracellular Ca^{2+} concentration) can result in hyperpolarization of the membrane potential from the resting state. The mechanism for this hyperpolarization likely lies in the blockage of a Na^+ leak channel (NALCN) and/or voltage-gated Na^+ channels. The effects may also be connected to calcium-activated potassium channels. Using *Drosophila melanogaster*, we here illustrate that changing $[\text{Ca}^{2+}]_O$ from 0.5 to 3 mM hyperpolarizes the muscle. Replacing NaCl with LiCl or choline chloride still led to hyperpolarization when increasing $[\text{Ca}^{2+}]_O$. Replacing CaCl_2 with BaCl_2 results in depolarization. K2P channel overexpression in the larval muscle greatly reduces the effects of $[\text{Ca}^{2+}]_O$ on cell membrane potential, likely because potential is heavily driven by the E_K in these muscles. These experiments provide an understanding of the mechanisms behind neuronal hypo-excitability during hypercalcemia, as well as the effects of altered expression of K2P channels on membrane potential.



Citation: Elliott, E.R.; Cooper, R.L. The Effect of Calcium Ions on Resting Membrane Potential. *Biology* **2024**, *13*, 750. <https://doi.org/10.3390/biology13090750>

Academic Editor: Luis Gandía

Received: 5 August 2024

Revised: 29 August 2024

Accepted: 3 September 2024

Published: 23 September 2024

Keywords: calcium; ion substitution; K2P channels; membrane potential; potassium

Copyright: © 2024 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Resting membrane potential (R_p) is known to vary amongst cells, both within an organism and even at the tissue level, which has been attributed to differences in the types of ion channels present and the density of the channels themselves [1]. Some cells utilize ligand-gated channels or those moderated by cellular responses, but the channels primarily involved in the electrical activity for conduction are passive or voltage-gated channels [2]. The specific distribution of these channels varies by cell, as each has a unique molecular composition. However, at rest, the two main channels present are the two pore-domain potassium (K2P) and sodium (NALCN) leak channels [2,3]. Since most cells have a larger P_K (permeability to K^+ ions) than P_{Na} (permeability to Na^+ ions), the cell's R_p is largely driven towards the potassium equilibrium potential (E_K).

Interestingly, variations in extracellular Ca^{2+} ($[\text{Ca}^{2+}]_O$) are known to result in altered R_p without Ca^{2+} flux [4]. $[\text{Ca}^{2+}]_O$ is known to affect conduction in central neurons and play a large role in synaptic transmission [5–7]; it has also been proposed that Ca^{2+} affects R_p through the activation of calcium-activated potassium channels (K_{Ca}) and/or alteration in the kinetics or blockage of voltage-gated Na^+ channels [8,9] and NALCN channels [2,3]. The R_p of a cell is generally more depolarized than the E_K value but raised $[\text{Ca}^{2+}]_O$ can block NALCN channels and cause hyperpolarization [3]. The NALCN channel was biophysically characterized by Lu et al. [10] and the importance of its role in physiology and pathological conditions is discussed in Monteil et al. [3]. Ca^{2+} ions block some types of voltage-gated ion channel, particularly of the Na^+ variety [8–12], but the resulting hyperpolarization can also potentially remove the residual inactivation of voltage-gated Na^+ channels and thus lower the threshold of a cell's excitability. Changes to electrical potential may also occur as Ca^{2+} alters the screening of charges across membranes [13].

This investigation—regarding the influence of Ca^{2+} ions on membrane potential—was conducted with the muscles of larval *Drosophila melanogaster*. The effects of ion substitution were observed through a reduction $[\text{Na}^+]$ of the bathing saline via replacement of the NaCl with choline chloride or LiCl . Ion substitution replacing Ca^{2+} with Ba^{2+} was conducted to address whether similar hyperpolarization is observed during exposure to another divalent cation capable of conduction through Ca^{2+} channels, as well as to investigate whether some small amount of Ca^{2+} might leak into the cell when $[\text{Ca}^{2+}]_O$ is high, since Ba^{2+} would not activate the K_{Ca} channels present [14]. Various buffers were used and brought to an appropriate pH with HCl to maintain physiological levels while avoiding the use of NaOH or NaHCO_3 .

Using *Drosophila* as a model allowed for genetic alteration of channel expression (specifically, that of K2P channel subtypes) and adjustment of the relative relationship between P_{Na} and P_K , which helped determine whether high $[\text{Ca}^{2+}]_O$ conditions still illustrate a reduction of P_{Na} through NALCN channels. Altered K2P expression is known to occur (to varied degree depending on the channel subtype involved) in cancerous/diseased mammalian tissues, though it is yet unknown whether they are a cause or a consequence of these pathologies [15,16]. Hypo- and hypercalcemia are known to have physiological effects on the excitability of human neurons, with the former possibly resulting in Chvostek's sign (twitches of the face due to activation of the motor nerve) or Trousseau's sign (slow contraction of the hand muscles), though the specifics of these effects, other health outcomes, and even the mechanisms behind altered $[\text{Ca}^{2+}]_O$ on membrane excitability are not fully understood [17,18].

Knowing that larval *Drosophila* muscle has an E_{Cl} of around -40 mV, [19,20] an E_K of about -90 mV [21,22], and $[\text{Ca}^{2+}]_O$ of around 1.5 mM in the hemolymph [23] provides a foundational understanding of physiological parameters that would be helpful while trying to address alterations in membrane function caused by changes in $[\text{Ca}^{2+}]_O$. Additionally, the fact that *Drosophila* represent a genetically amenable model allows examination of how altered expression of K2P channels affects R_p and the effects of varied $[\text{Ca}^{2+}]_O$ on it. This study is intended to provide an initial assessment of how altered $[\text{Ca}^{2+}]_O$ affects the R_p of *Drosophila* muscle to foster future studies into the detailed mechanism behind the observations included herein. *Drosophila* genetics being easily manipulable renders them a good model for determining how altered expression of specific proteins affects regulation, allowing the support or refutation of proposed mechanisms.

2. Materials and Methods

2.1. Animals

Drosophila melanogaster Canton S (CS) flies were used in physiological assays. This strain was originally obtained from the Bloomington *Drosophila* Stock Center (BDSC) but has been isogenic in the laboratory since 1996. *Drosophila* CS larvae were used as early third instars (50–70 h post-hatching). Overexpression of the ORK1 protein in larval body wall muscles (m6 and m7) was achieved by crossing homozygous males of BG487 (BDSC

stock # 51634) with female virgins of UAS-ORK1 (BDSC stock # 6586). Progeny carrying one copy each of GAL4 driver and UAS-ORK1, referred to as body muscle m6-m7 > ORK1, were used. Larval body wall muscles 6/7 feature an anteroposterior gradient pattern of BG487-Gal4 expression, allowing BG487 to drive UAS-ORK1 in those muscles [24,25]. The larvae were maintained at room temperature, ~21 °C, in vials partially filled with a cornmeal-agar-dextrose-yeast medium.

2.2. Dissection and Physiology

Similar dissection procedures and electrophysiological measures have been described previously [26]. Briefly, transmembrane potentials were monitored in m6 muscles of early third-instar larvae using sharp intracellular electrodes (30 to 40 megaOhm resistance) filled with 3 M KCl. An Axonclamp 2B (Molecular Devices, Sunnyvale, CA, USA) amplifier and 1 X LU head stage was used. LabChart 7.0 (ADInstruments, Colorado Springs, CO, USA) was used to collect and analyze data.

The saline used for dissection was haemolymph-like 3(HL3) [23,27]: (in mmol/L) 70 NaCl, 5 KCl, 20 MgCl₂, 10 NaHCO₃, 1 CaCl₂, 5 trehalose, 115 sucrose, 25 N,N-bis (2-hydroxyethyl)-2-aminoethane sulfonic acid (BES) and pH at 7.2. The CaCl₂ was varied from 0.5 to 3.0 mM as described in the Results for the various experiments. CaCl₂ was replaced by BaCl₂, and NaCl replaced by LiCl or choline chloride in experiments mentioned. Two other buffers besides BES were examined for their suitability. These were Trizma[®] base (2-amino-2-(hydroxymethyl)-1,3-propanediol) and CAPS (3-(Cyclohexylamino)-1-propanesulfonic acid). All chemical were from Sigma-Aldrich (St. Louis, MO, USA).

2.3. Statistical Analysis

Data are expressed as averages (\pm SEM-standard error of the mean). Response differences before and after solution exchange were quantified with paired *t*-tests. The Shapiro-Wilk test was used to establish normality. When appropriate, the Wilcoxon rank sum, non-parametric test was used. Two-way analysis of variance (ANOVA) was performed with multiple comparisons among different larval strains with Tukey's method since sample sizes were the same. Sigma Stat software (version number 15.0) was used for analysis, and a *p*-value of *p* < 0.05 was considered statistically significant.

3. Results

3.1. Altering [Ca²⁺]_o Effects on Membrane Potential

Membrane potential underwent acute hyperpolarization with increased [Ca²⁺]_o. Since larval body wall muscle was used, the presynaptic motor nerve terminal continued to exhibit spontaneous vesicular fusion events (i.e., minis or quantal events) illustrated by rapid upward deflections on the recording. The motor nerve was transected from the CNS, resulting in a continued observance of spontaneous events that decreased in frequency as time proceeded. Exposure to 3 mM Ca²⁺ affected the muscle fiber membrane within seconds and the effect was generally maintained throughout the full three minutes of observation (Figure 1A). Hyperpolarization occurred when the bath was exchanged from 0.5 mM to 3 mM (Figure 1B), and this effect was found to be significant (paired *t*-test; *p* < 0.05; *n* = 10) with an average percent change of 7.5 (SEM \pm 1.6) mV (Figure 1C,D). Some preparations showed greater changes than others but overall, there was a hyperpolarization with exposure to 3 mM. The rationale behind changing the medium from normal saline (with [Ca²⁺]_o at 1 mM) to the low 0.5 mM experimental concentration was that this alteration would increase the difference between the low and high calcium environments and thus accentuate the observed effects on R_p. If [Ca²⁺]_o is removed from the saline, there is a higher probability of the motor nerve firing action potentials and causing muscle contraction, which can dislodge the intracellular electrode; thus, 0.5 mM was the lowest concentration used in this paradigm.




Figure 1. The effects on resting membrane potential of altered extracellular $[\text{Ca}^{2+}]$. (A) A representative recording of the membrane potential with alterations of the bathing medium. (B) The responses for 10 individual preparations as $[\text{Ca}^{2+}]_O$ was changed. Raised $[\text{Ca}^{2+}]_O$ led to significant hyperpolarization of the membrane (paired t -test; $p < 0.05$) compared to initial values. (C) The average membrane potential for 0.5 mM $[\text{Ca}^{2+}]_O$ and 3 mM $[\text{Ca}^{2+}]_O$ (mean \pm SEM) showed no significant difference due to the large variation in membrane potentials among preparations. (D) The mean percent change for each of the 10 individual preparations (mean \pm SEM). The light blue boxes represent exchange of the medium from one solution to the next.

3.2. Choline Chloride Experiments

A representative recording in which saline was exchanged for that containing increased Ca^{2+} and a replacement of Na^+ with choline chloride is depicted in Figure 2A. With the exchange of NaCl (70 mM) in HL3 saline for choline chloride (70 mM) and of $[\text{Ca}^{2+}]_O$ from 0.5 mM to 3.0 mM, significant hyperpolarization of the membrane was observed (Figure 2A,B; paired t -test; $p < 0.05$; $N = 10$). The average percent change was 7.4 (SEM \pm 2.1)

mV (Figure 2C,D). The response variation observed appears to be a function of normal within-preparation physiological variation.




Figure 2. The effects on resting membrane potential of substituting choline chloride for NaCl and altering extracellular [Ca²⁺]_O, but with Na⁺ present from NaHCO₃⁻ and the NaOH used for pH adjustments. (A) A representative trace of the membrane potential as the medium is changed from saline containing NaCl to that containing choline chloride, to 0.5 mM [Ca²⁺]_O, to 3.0 mM, back to 0.5 mM, and then to HL3 saline once more. (B) The responses for 10 individual preparations during altered [Ca²⁺]_O while exposed to choline chloride instead of NaCl. Raised [Ca²⁺]_O led to significant hyperpolarization of the membrane (paired *t*-test; *p* < 0.05). (C) The average membrane potential for 0.5 mM [Ca²⁺]_O and 3 mM [Ca²⁺]_O (mean ± SEM) showed no significant difference due to the large variation in membrane potentials among preparations. (D) The mean percent change from initial for each of the 10 individual preparations (mean ± SEM). The light blue boxes represent exchange of the medium from one solution to the next. There was a significant difference in the percent change from 0.5 to 3.0 mM (* paired *t*-test; *p* < 0.05).

The hyperpolarization observed after exposure to 3 mM [Ca²⁺]_O (with choline chloride in place of NaCl in the saline) was unexpected for a NaCl-free solution; in such a situation, no significant difference in membrane potential would be expected between 1 mM and 3 mM [Ca²⁺]_O. This response is likely because the HL3 saline used both contained NaHCO₃ and was brought to a pH of 7.2 using NaOH, which would result in the presence of

additional sodium in the solution despite the removal of NaCl. These experiments were thus redone with new solutions that contained choline chloride in place of NaCl while replacing NaHCO₃ with, respectively, Trizma[®] base (Figure 3) and CAPS (Figure 4) in an effort to eliminate sodium from the solution. HCl was used to adjust the pH to 7.2.




Figure 3. The effects on resting membrane potential of substituting choline chloride for NaCl and Trizma[®] base for NaHCO₃, as well as altering [Ca²⁺]_o. (A) A representative trace of the membrane potential as the medium is changed from saline (HL3-BES) containing NaCl to that containing choline chloride/Trizma[®] base, to 1 mM [Ca²⁺]_o, to 3.0 mM, back to 1 mM, and then to HL3 saline once more. (B) The responses for 10 individual preparations during altered [Ca²⁺]_o while exposed to choline chloride and Trizma[®] base. (C) The average membrane potential in HL3 saline, 1 mM [Ca²⁺]_o, 3 mM [Ca²⁺]_o, 1 mM [Ca²⁺]_o again, and HL3 again (mean ± SEM). Significant depolarization of the membrane was observed (* *p* < 0.05, ANOVA for initial HL3-BES to 1 mM after 3 min of incubation and 3 mM [Ca²⁺]_o as well as to returning to 1 mM in Trizma[®]; Paired *t*-test were significant *p* < 0.05 for comparing initial HL3-BES to 1 mM after 3 min of incubation and 3 mM [Ca²⁺]_o as well), though it hyperpolarized again when the medium was returned to HL3-BES saline. (D) The mean percent change from initial for each of the 10 individual preparations (mean ± SEM). The light blue boxes represent exchange of the medium from one solution to the next. There was a significant difference in the percent changes, * paired *t*-test; *p* < 0.05).




Figure 4. The effects on resting membrane potential of substituting choline chloride for NaCl and CAPS for NaHCO₃, as well as altering extracellular [Ca²⁺]_O. (A) A representative trace of the membrane potential as the medium is changed from saline containing NaCl to that containing choline chloride/CAPS, to 1 mM [Ca²⁺]_O, to 3.0 mM, back to 1 mM, and then to HL3 saline once more. (B) The responses for 10 individual preparations during altered [Ca²⁺]_O while exposed to choline chloride and CAPS. Significant depolarization of the membrane was observed (ANOVA for initial HL3-BES to 3 mM [Ca²⁺]_O as well as to returning to 1 mM in CAPS; Paired *t*-test were significant *p* < 0.05 for comparing initial HL3-BES to 1 mM after 3 min of incubation and 3 mM [Ca²⁺]_O as well) and did not re-hyperpolarize even upon return to HL3 saline or examination of adjacent muscle fibers. (C) The average membrane potential in HL3 saline, 1 mM [Ca²⁺]_O, 3 mM [Ca²⁺]_O, 1 mM [Ca²⁺]_O again, and HL3 again (mean ± SEM). Significant differences were observed (* *p* < 0.05, ANOVA for percent difference of HL3-BES 3 min to 1 mM CAPS at 3 min to 3 min of 3 mM [Ca²⁺]_O as well as to returning to 1 mM in CAPS; * Paired *t*-test were significant *p* < 0.05). (D) The mean percent change from initial for each of the 10 individual preparations (mean ± SEM). There was a significant difference in the percent changes, * paired *t*-test; *p* < 0.05. The light blue boxes represent exchange of the medium from one solution to the next.

3.3. LiCl Experiments

Another paradigm involved replacing the NaCl (70 mM) present in HL3 saline with LiCl at 70 mM. A representative recording is shown in Figure 5A. The hyperpolarization was delayed in the Li⁺ saline for some preparations as compared to HL3 (Figure 1A) and choline-chloride-containing saline (Figure 2A). Observations were conducted over a three-minute period and the largest hyperpolarization within that time was used. In some cases, the membrane potential depolarized from its most hyperpolarized state by the end of the three minutes (Figure 5B,C). The saline containing Li⁺ also demonstrated a significant hyperpolarization when the bath was exchanged from one containing 0.5 mM Ca²⁺ to 3.0 mM Ca²⁺ (paired *t*-test; *p* < 0.05; N = 10). The percent change from 0.5 mM to the end of 3 min' incubation in 3.0 mM Ca²⁺ saline solution resulted in a hyperpolarization of 2.9 (SEM ± 2.8) mV (Figure 5D).




Figure 5. The effects of replacing NaCl with LiCl and altering $[Ca^{2+}]_O$ on resting membrane potential. (A) A representative trace of the membrane potential change observed when NaCl is replaced with LiCl and during subsequent exposure to $[Ca^{2+}]_O$ 0.5 mM to 3.0 mM and back. (B) The responses observed across 10 individual preparations as $[Ca^{2+}]_O$ is changed during LiCl exposure. The effect of raised $[Ca^{2+}]_O$ produced significant hyperpolarization of the membrane (paired *t*-test; * *p* < 0.05) compared to initial values. (C) The average membrane potential between 0.5 mM $[Ca^{2+}]_O$ and 3.0 mM $[Ca^{2+}]_O$ (mean ± SEM) showed no significant difference among preparations after 3 min. (D) The mean percent change for each of the 10 individual preparations (mean ± SEM). The light blue boxes represent exchange of the medium from one solution to the next.

3.4. BaCl₂ Experiments

To examine whether the hyperpolarizing effects of increased $[Ca^{2+}]_O$ was specific to the Ca²⁺ ion, another divalent ion was investigated. Ba²⁺ can generally replace Ca²⁺ flux through Ca²⁺ channels, so the CaCl₂ normally present in HL3 saline was replaced

with BaCl₂. The general trends are shown for a representative preparation (Figure 6A). Replacing HL3 saline containing 1.0 mM CaCl₂ with that containing 1 mM BaCl₂ resulted in a significant depolarization of the membrane ($p < 0.05$; paired t -test, $N = 10$). Likewise, raising [Ba²⁺]_O from 1 to 3 mM led to significant depolarization of the membrane (paired t -test; $p < 0.05$). The average membrane potential for 1 mM [Ba²⁺]_O and 3 mM [Ba²⁺]_O (mean ± SEM) was significantly different (Figure 6B,C; The values were not normally distributed; thus, a Wilcoxon Signed Rank Test was used, $p = 0.031$). The average percent change from HL3 with 1 mM CaCl₂ to 1 mM BaCl₂ (7.2 SEM ± 3 mV) and from 1 mM BaCl₂ to 3.0 mM BaCl₂ (6.1 SEM ± 2.1 mV) are shown in Figure 6D. The percent change was taken from the end of the three minutes' exposure in one solution to the end of three minutes' exposure in the next.




Figure 6. The effects on resting membrane potential of substituting BaCl₂ for CaCl₂ and altering [Ba²⁺]_O. (A) A representative trace of the membrane potential as the medium is changed from saline containing CaCl₂ to that containing 1 mM BaCl₂, to 3.0 mM BaCl₂, back to 1 mM BaCl₂, and then to HL3 saline once more. (B) The responses for 10 individual preparations during replacement of CaCl₂ with 1 mM BaCl₂, then 3 mM BaCl₂, back to 1 mM BaCl₂, and then back to HL3. (C) The average membrane potential for 1 mM [Ba²⁺]_O and 3 mM [Ba²⁺]_O (mean ± SEM) was significantly different (Not normally distributed, thus used a Wilcoxon Signed Rank Test, $p = 0.031$) (D) The mean percent change from initial for each of the 10 individual preparations (mean ± SEM). Raised [Ba²⁺]_O led to significant depolarization of the membrane potential for a percent change, * paired t -test; $p < 0.05$). The light blue boxes represent exchange of the medium from one solution to the next.

3.5. The Effects on Membrane Potential in Muscle Overexpressing K2P Channels

The genetically modified *Drosophila* line (m6-m7 > ORK1, bearing an overexpression of the ORK1 K2P channel subtype) underwent no significant effects to membrane potential when exposed to raised $[Ca^{2+}]_O$. The response for a representative preparation is shown (Figure 7A). The muscle fiber membrane potential was not significantly altered by the exchange from 0.5 to 3 mM Ca^{2+} (Figure 7B,C) ($p > 0.05$; Paired *t*-test; $N = 10$), and the average percentage change was also minor, with no significant effect (Figure 7D).




Figure 7. The effects of altering $[Ca^{2+}]_O$ on resting membrane potential for muscle overexpressing K2P channels. (A) A representative trace of the membrane potential changes as $[Ca^{2+}]_O$ is changed from 1.0 mM to 0.5 mM to 3.0 mM and back to 0.5 mM. (B) The responses for 10 individual preparations as $[Ca^{2+}]_O$ were changed. The effect of raising $[Ca^{2+}]_O$ from 0.5 to 3.0 mM had no significant hyperpolarization effect on the membrane (paired *t*-test; $p > 0.05$). (C) The average membrane potential for 0.5 mM $[Ca^{2+}]_O$ and 3 mM $[Ca^{2+}]_O$ (mean \pm SEM) showed no significant difference among preparations. (D) The mean percent change for each of the 10 individual preparations (mean \pm SEM). The light blue boxes represent where the exchange of saline took place.

4. Discussion

In this investigation, it was demonstrated that membrane potential was affected by $[Ca^{2+}]_O$ such that raising the concentration resulted in hyperpolarization, and vice versa. Replacing the NaCl in the saline with LiCl or choline chloride and then exchanging the $CaCl_2$ from 0.5 mM to 3.0 mM caused a hyperpolarization of the membrane. Replacing

CaCl₂ with BaCl₂ led to depolarization of the membrane and increasing the BaCl₂ from 0.5 mM to 3.0 mM resulted in further depolarization. In larval muscle overexpressing K2P channels (ORK1), the effects of increased [Ca²⁺]_o on membrane potential were greatly reduced, likely because high K⁺ permeability with more leak channels causes a strong tendency to remain at E_K; this indicates that the ion channel density and K2P channel subtypes present change how membrane potential is affected by altered [Ca²⁺]_o. The results herein confirm previous findings that larval *Drosophila* muscle hyperpolarizes with raised [Ca²⁺]_o [4].

To investigate potential mechanisms, Ca²⁺ was replaced with Ba²⁺, which resulted in depolarization instead of hyperpolarization. Given that the resting membrane resides not at E_K but a slightly depolarized value, the data suggest an ionic leak, such as a small Na⁺ leak. Considering the possibility that Ca²⁺ ions may block NALCN channels, as has been suggested for other cell types [8,9,11,12], increasing [Ca²⁺]_o would potentially block more NALCN channels, resulting in a greater drive towards E_K. If Ba²⁺ displaced any residual Ca²⁺ ions from the channels, however, and thus reduced the blockage of NALCN channels, it would likely result in depolarization. If Ba²⁺ leaked through a NALCN channel, this would also cause depolarization, but there is no precedent to indicate this.

Experimental paradigms addressing the effects of altered [Ca²⁺]_o on synaptic efficiency should consider this phenomenon, as one might otherwise focus only on the presynaptic influence of increased Ca²⁺ flux via more evoked vesicle fusion events and greater post-synaptic amplitudes. However, alterations in the Ca²⁺ concentration gradient would also affect the electrical driving gradient, such that elevated [Ca²⁺]_o might hyperpolarize the neuronal membrane potential and lead to removal of the inactivation from some voltage-gated Na⁺ channels present, resulting in a lowered excitation threshold for the neuron. Hyperpolarization of the target cell could also influence ionic driving gradients and action potential threshold in the same manner as for the presynaptic neuron. Finally, the membrane potential of larval *Drosophila* muscle is also pH-sensitive [28]; this may impact some K2P channel subtypes, given that the TASK subtype is blocked by acidic conditions [29], and may also affect Ca²⁺ interaction with NALCN channels. Thus far, this topic has largely gone unaddressed, experimentally. It is also of interest to investigate whether the ionized [Ca²⁺] may be decreased by the sodium bicarbonate buffer used in saline or by interactions with other compounds.

The concentration of free Ca²⁺ ions is regulated by various means in intact organisms (hormones, calcium binding proteins) as well as within cells and cellular organelles. The physiological range in human serum is considered to be 2.25 to 2.75 mmol/L for neonates and adults [30,31], while the cerebrospinal fluid (CSF) in healthy humans contains ionized (~1 mM) and total (~1.2 mM) calcium concentration in the CSF to match concentrations in the brain extracellular fluid [32,33]. A major buffer of ionized Ca²⁺ in CSF is carbonate. As is the case for albumin, Ca²⁺ is much lower in CSF than in serum, so regulating saline pH may alter the assumed concentration of ionized Ca²⁺ depending on the buffer used to control pH. Given that chemical synaptic transmission is strongly influenced by Ca²⁺ entry to the presynaptic terminal, the mechanisms behind this may not entirely be due to a concentration gradient, as both the electrical gradient and the modulation of ion channels by the electrostatic interactions can have a role.

Despite years of research with genomic and proteomic tools, it is not yet known how many subtypes of K2P and NALCN channels are expressed and functional within any single cell in which the membrane potential can be measured. In time, it will be possible to know which of the 11 genes known to express K2P channels in *Drosophila melanogaster* are expressed in various cell types across development; in the meantime, however, pharmacological and environmental changes can aid in investigating some of the subtypes present in the membranes. Cold temperatures, acidic conditions, and the compound doxapram (an inhibitor of a K2P-TASK subtype) are known to depolarize larval *Drosophila* muscle [28,34,35]. However, with a pH of 5 or a high concentration of doxapram, the resting membrane does not rapidly reach a potential of zero or E_{Na}, suggesting that

many other factors (such as pumps, exchangers, and different K2P channel subtypes) may be at play in maintaining a membrane potential. On the other hand, overexpressing a dORKA1 K2P channel in larval *Drosophila* muscle produces membrane potentials that are much more negative and closer to E_K . [26]. With more leak channels present and, thus, a higher K^+ permeability, it would be surprising to observe altered $[Ca^{2+}]_O$ (and the resulting reduction or enhancement of Na^+ leak channel blockage) affecting membrane potential. Rather, as demonstrated in this study, the membrane potential of a given cell would vary depending on the density of K^+ and Na^+ leak channels present.

The NALCN channel subtype expressed in the skeletal muscle of larval *Drosophila* has yet to be identified; however, it is likely that a form of Na^+ leak channel is present, as the R_p is not maintained at E_k (~ -90 mV) and overexpression of a K2P channel hyperpolarizes the membrane potential from wild-type *Drosophila*. An NALCN antibody for a mammalian channel subtype is commercially available [2], and it might be worthwhile to accumulate *Drosophila* body wall tissue and test the antibody's effectiveness in the model through the use of Western blots. Some pharmacological agents block NALCN channels in mammalian tissues and could potentially affect *Drosophila* preparations as well [2]. It would also be of interest to use RNA-Seq to examine the effects of altered K2P and NALCN channel expression in other *Drosophila* tissues; this approach would allow RNAi targeting of specific subtypes and tissues, as well as addressing the fact that K2P and NALCN channels are not solely responsible for functional expression because the associated accessory proteins are significant as well. It is even possible that K2P overexpression somehow affected ion channel density or incorporation into the membrane which would not be determined by Western blot or even RNA-Seq. However, these approaches are beyond the scope of this current study. The NALCN protein is associated with various auxiliary subunits (UNC80 and UNC79, as well as others), which form a complex now known as an NALCN channelosome [36]. Examining how these subunits impact channel function could be addressed in future studies with *Drosophila*, as these other proteins are well-established in this model.

Removing all Na^+ from the bathing medium was problematic because it prevented both pH adjustment with NaOH and use of the $NaHCO_3^-$ buffer. Basic buffers such as CAPS and Trizma[®] Base were explored as possible replacements to allow pH adjustment with HCl; however, the larval muscle did not fare well with these buffers, as the membrane would depolarize even with $[Ca^{2+}]_O$ held constant. The molecular structures of CAPS and Trizma[®] Base differ from the BES buffer, which, as has been previously discovered, best maintains physiological function of the larval heart for the hours needed to conduct in situ studies [27,37]. In the same experiments, a wide array of possible buffers was examined, and none performed so well as BES. It is unknown why CAPS and Trizma[®] Base altered the membrane potential in this investigation, as pH was maintained at 7.2 and the other salts remained consistent (save for the aforesaid absence of $NaHCO_3$). Other buffers, such as 4-(2-Hydroxyethyl)piperazine-1-ethanesulfonic acid (HEPES), might be feasible replacements, but attempting to use HEPES resulted in an acidic medium that could not be adjusted without using NaOH. Cesium hydroxide (CsOH) could feasibly be used for pH adjustments, but the effects of Cs^+ in solution would also need to be examined. Other buffers might be useful as replacements, but their effects on cell viability would need to be tested first.

Various schematic models based on the manipulations used in this study are presented to help explain the overall mechanisms behind the alterations in membrane potential observed here. This first model highlights the effects of raised $[Ca^{2+}]_O$ on the cells' membrane potentials (Figure 8).

The second model illustrates the effects of reducing Na^+ through replacement with choline chloride to maintain osmolarity while reducing $[Na^+]_O$ (Figure 9).




Figure 8. The effects of altered $[Ca^{2+}]_o$ on resting membrane potential. Ca^{2+} is believed to influence resting membrane potential (R_p) by reducing Na^+ influx through the corresponding leak channel (NALCN). At rest, when $[Ca^{2+}]_o$ is at physiological levels, the Na^+ leak into the muscle is slightly hindered (**top panel**). As $[Ca^{2+}]_o$ is reduced, the Na^+ leak is enhanced and R_p depolarizes (**middle panel**). As $[Ca^{2+}]_o$ is raised, NALCN channels are blocked to a greater degree than observed at physiological $[Ca^{2+}]_o$ and the R_p is hyperpolarized as it approaches equilibrium potential for K^+ (**bottom panel**). The thickness of the arrows illustrates the changes occurring.




Figure 9. The effects of altered $[Na^+]_o$ on resting membrane potential and spontaneous nerve activity. $[Na^+]_o$ was reduced by replacing the NaCl in physiological saline with equal concentrations of choline chloride (i.e., 70 mM for *Drosophila* saline). Lower $[Na^+]_o$ produced hyperpolarization of the larval muscle's resting membrane potential, likely due to reduced drive on the Na^+ leak channel (NALCN). In addition, spontaneous quantal events occurred, as well as large postsynaptic excitatory junction potentials (EJPs) appearing as nerve-evoked EJPs. It is apparent that more single and multi-quantal events occurred, which indicates a presynaptic response that may be driven by the passive sodium-calcium exchanger (NCX). This response likely results in an influx of Ca^{2+} due to the low driving gradient for Na^+ influx, particularly compared to the stronger gradient for Na^+ efflux. In addition, Na^+ might even efflux from the muscle fiber. The thickness of the arrows illustrates the changes occurring.

The third model illustrates the replacement of NaCl with LiCl and subsequent alterations of $[Ca^{2+}]_O$, which indicates that raised $[Ca^{2+}]_O$ results in more blockage of the Na^+ leak (Figure 10). The explanation behind increased $[Ca^{2+}]_O$ hyperpolarizing the membrane in this paradigm is not fully understood. It is possible that the sodium-potassium pump or other pumps/exchangers work differently in these conditions [38].




Figure 10. Alterations in $[Ca^{2+}]_O$ during substitution of Li^+ for Na^+ affect resting membrane potential. In basal conditions Ca^{2+} is blocking some of the NALCN channels (top panel). Alterations in $[Ca^{2+}]_O$ after replacement of Na^+ with Li^+ resulted in similar responses to those observed when Na^+ was present. Reducing $[Ca^{2+}]_O$ likely decreased the slight blockage of the Na^+ leak channel (NALCN), resulting in depolarization of the muscle fiber (middle panel). On the other hand, raising $[Ca^{2+}]_O$ enhanced the block, promoting the action of K2P channels and resulting in membrane hyperpolarization towards the equilibrium potential for K^+ (bottom panel). The thickness of the arrows illustrates the changes occurring.

The results upon substituting $BaCl_2$ for $CaCl_2$ are interesting, as it was expected that the cellular response would be similar to those observed with varying $[Ca^{2+}]_O$, since Ba^{2+} could potentially block the Na^+ leak channel in a similar manner. Instead, it appears that the cells slightly depolarized in the $BaCl_2$ solution, with greater depolarization observed at higher concentrations (Figure 11). The NALCN channels may be maximally disinhibited even at 0.5 mM Ba, such that no potential change would be expected upon raising Ba^{2+} to 3 mM (Figure 11, bottom panel).

$BaCl_2$ has been used in past studies to address the possibility that calcium influx activates $K_{(Ca)}$ channels from the cytoplasmic side or through direct blockage of voltage-gated K^+ channels [14,39,40]. It appears as though replacing Ca^{2+} with Ba^{2+} enables Ba^{2+} influx through Ca^{2+} channels without activation of $K_{(Ca)}$ channels. In the muscle cells studied during this investigation, it is not likely that the hyperpolarization stems from a Ca^{2+} leak activating $K_{(Ca)}$ channels, as such a leak would likely result in muscle contractions not observed in this study. Spontaneous quantal events weren't observed either, which would also be expected in the presence of increased $[Ca^{2+}]_O$. However, Ba^{2+} might block the K2P channel, as is known to occur for voltage-gated K^+ channels [3].




Figure 11. Replacing Ca^{2+} with Ba^{2+} and its effect on resting membrane potential. When $[\text{Ca}^{2+}]_o$ is at physiological levels, the Na^+ leak (NALCN) channels in the muscle is slightly blocked (**top panel**). After removing Ca^{2+} and replacing it with Ba^{2+} at the same concentrations used for examining the effects of $[\text{Ca}^{2+}]_o$ alteration on resting membrane potential (R_p) (**middle panel**), increasing $[\text{Ba}^{2+}]_o$ resulted in reduced blockage of NALCN channels (**bottom panel**).

Cells would likely exhibit very negative resting membrane potentials if they also demonstrated a reduced level of Na^+ leak or a higher expression of functional K2P channels; the results thus indicate a high level of K^+ leak that renders negligible the effects from both the Na^+ leak and the Ca^{2+} -induced blockage of the Na^+ leak (Figure 12).




Figure 12. When K2P channels are overexpressed in muscle fibers, resting membrane potential (R_p) is not influenced by altered $[\text{Ca}^{2+}]_o$. R_p was observed with Ca^{2+} reducing Na^+ influx through the Na^+ leak channel (NALCN) at 1 mM (**top panel**), with a reduced concentration of 0.5 mM $[\text{Ca}^{2+}]_o$ (**middle panel**), and with a raised concentration of 3 mM $[\text{Ca}^{2+}]_o$ (**bottom panel**). In each condition, the permeability of Na^+ (P_{Na}) was negligible as compared to the high permeability of K^+ (P_{K}), driving R_p towards the equilibrium potential for K^+ (E_K) (**bottom panel**).

It is not likely that $K_{(Ca)}$ -channel activation accounts for the hyperpolarization observed with raised $[Ca^{2+}]_O$. The muscle does not appear to respond well to elevated $[Ca^{2+}]_O$, as the muscle shows granulation over time; it appears as though, over the 10 min of observation, increased $[Ca^{2+}]_O$ damages the muscle cell in CS strains but not in K2P-overexpressing strains. Ca^{2+} ions were noted to block voltage-gated ion channels and have an influence on the kinetics of various channels [8]. While recording compound action potentials (CAPs) in marine crab nerves, it was demonstrated that replacing $CaCl_2$ with $BaCl_2$ resulted in depressed CAP amplitude, possibly indicating a blockage of some voltage-gated Na^+ channels [5]. Earlier studies addressed the potential that Ca^{2+} screens charges on membranes [13]. One might have expected $BaCl_2$ to affect membrane potential similarly to $CaCl_2$.

In the future, it would be of interest to investigate the distribution of ion channel subtypes responsible for the resting membrane potential; it would also be worthwhile to model ion permeabilities and concentration differences through simulations of the Goldman-Hodkin-Katz equation, which would allow examination of how slight differences in leak channel expression (both K2P and NALCN) affect organisms. Since the expression of K2P channels in human tissues varies under pathological conditions, such as cancer [15,16,29], it would also be of interest to investigate the impact of varied ionic concentrations on different cell types with varied membrane potentials.

5. Conclusions

Changing external calcium concentrations from 0.5 to 3 mM led to hyperpolarization of the muscle. Replacing NaCl with LiCl or choline chloride still led to hyperpolarization as the calcium concentration increased. Replacing $CaCl_2$ with $BaCl_2$ results in depolarization. Larval muscle bearing K2P channel overexpression largely saw greatly reduced effects with altered $[Ca^{2+}]_O$, likely because potential is heavily driven by the E_K in these muscles.

Author Contributions: Conceptualization, E.R.E. and R.L.C.; methodology, E.R.E. and R.L.C.; software, E.R.E. and R.L.C.; validation, E.R.E. and R.L.C.; formal analysis, E.R.E. and R.L.C.; investigation, E.R.E. and R.L.C.; resources, R.L.C.; data curation, R.L.C.; writing—original draft preparation, E.R.E. and R.L.C.; writing—review and editing, E.R.E. and R.L.C.; visualization, E.R.E. and R.L.C.; supervision, R.L.C.; project administration, R.L.C.; funding acquisition, E.R.E. and R.L.C. All authors have read and agreed to the published version of the manuscript.

Funding: College of Arts and Sciences Summer Research Fellowship and a Gertrude Flora Ribble Scholarship from the Department of Biology at the University of Kentucky (E.R.E.). Beckman Scholarship (E.R.E.). Chellgren Endowed Funding (R.L.C.). Alumni of the research group (R.L.C.).

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: The data presented in this study are available on request from the corresponding author. The data are not publicly available due to requiring specific software to view the data files. Most of the data are presented within this publication in the line graphs.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Malmivuo, J.; Plonsey, R. *Bioelectromagnetism-Principles and Applications of Bioelectric and Biomagnetic Fields*; Oxford University Press: New York, NY, USA, 1995; ISBN 9780199847839. [[CrossRef](#)]
2. Monteil, A.; Guérineau, N.C.; Gil-Nagel, A.; Parra-Diaz, P.; Lory, P.; Senatore, A. New insights into the physiology and pathophysiology of the atypical sodium leak channel NALCN. *Physiol. Rev.* **2024**, *104*, 399–472. [[CrossRef](#)] [[PubMed](#)]
3. Goldstein, S.A.; Price, L.A.; Rosenthal, D.N.; Pausch, M.H. ORK1, a potassium-selective leak channel with two pore domains cloned from *Drosophila melanogaster* by expression in *Saccharomyces cerevisiae*. *Proc. Natl. Acad. Sci. USA* **1996**, *93*, 13256–13261. [[CrossRef](#)] [[PubMed](#)]
4. Krans, J.L.; Parfitt, K.D.; Gawera, K.D.; Rivlin, P.K.; Hoy, R.R. The resting membrane potential of *Drosophila melanogaster* larval muscle depends strongly on external calcium concentration. *J. Insect Physiol.* **2010**, *56*, 304–313. [[CrossRef](#)]

5. Atkins, D.E.; Bosh, K.L.; Breakfield, G.W.; Daniels, S.E.; Devore, M.J.; Fite, H.E.; Guo, L.Z.; Henry, D.K.J.; Kaffenberger, A.K.; Manning, K.S.; et al. The effect of calcium ions on mechanosensation and neuronal activity in proprioceptive neurons. *NeuroSci* **2021**, *2*, 353–371. [[CrossRef](#)]
6. Augustine, G.J. How does calcium trigger neurotransmitter release? *Curr. Opin. Neurobiol.* **2001**, *11*, 320–326. [[CrossRef](#)]
7. Katz, B. *The Release of Neural Transmitter Substances*; Liverpool University Press: Liverpool, UK, 1969.
8. Armstrong, C.M.; Cota, G. Calcium block of Na⁺ channels and its effect on closing rate. *Proc. Natl. Acad. Sci. USA* **1999**, *96*, 4154–4157. [[CrossRef](#)] [[PubMed](#)]
9. Mert, T.; Gunes, Y.; Guven, M.; Günay, I.; Ozcengiz, D. Effects of calcium and magnesium on peripheral nerve conduction. *Pol. J. Pharmacol.* **2003**, *55*, 25–30. [[PubMed](#)]
10. Lu, B.; Su, Y.; Das, S.; Liu, J.; Xia, J.; Ren, D. The neuronal channel NALCN contributes resting sodium permeability and is required for normal respiratory rhythm. *Cell* **2007**, *129*, 371–383. [[CrossRef](#)] [[PubMed](#)]
11. Segal, M. Calcium stores regulate excitability in cultured rat hippocampal neurons. *J. Neurophysiol.* **2018**, *120*, 2694–2705. [[CrossRef](#)]
12. Woodhull, A.M. Ionic blockage of sodium channels in nerve. *J. Gen. Physiol.* **1973**, *61*, 687–708. [[CrossRef](#)] [[PubMed](#)]
13. Armstrong, C.M. Distinguishing surface effects of calcium ion from pore-occupancy effects in Na⁺ channels. *Proc. Natl. Acad. Sci. USA* **1999**, *96*, 4158–4163. [[CrossRef](#)] [[PubMed](#)]
14. Baraibar, A.M.; de Pascual, R.; Camacho, M.; Domínguez, N.; David Machado, J.; Gandía, L.; Borges, R. Distinct patterns of exocytosis elicited by Ca²⁺, Sr²⁺ and Ba²⁺ in bovine chromaffin cells. *Pflugers. Arch.* **2018**, *470*, 1459–1471. [[CrossRef](#)]
15. Lee, L.M.; Müntefering, T.; Budde, T.; Meuth, S.G.; Ruck, T. Pathophysiological role of K₂P channels in human diseases. *Cell Physiol. Biochem.* **2021**, *55*, 65–86. [[PubMed](#)]
16. Wiedmann, F.; Frey, N.; Schmidt, C. Two-Pore-Domain potassium (K₂P⁻) channels: Cardiac expression patterns and disease-specific remodelling processes. *Cells* **2021**, *10*, 2914. [[CrossRef](#)] [[PubMed](#)]
17. Kamalanathan, S.; Balachandran, K.; Parthan, G.; Hamide, A. Chvostek’s sign: A video demonstration. *BMJ Case Rep.* **2012**, *2012*, 1–2. [[CrossRef](#)] [[PubMed](#)]
18. Patel, M.; Hu, E.W. Trousseau Sign. 2023 May 1. In *StatPearls*; StatPearls Publishing: Treasure Island, FL, USA, 2024. Available online: <https://pubmed.ncbi.nlm.nih.gov/32491764/> (accessed on 26 June 2024).
19. Rose, U.; Derst, C.; Wanischek, M.; Marinc, C.; Walther, C. Properties and possible function of a hyperpolarisation-activated chloride current in *Drosophila*. *J. Exp. Biol.* **2007**, *210 Pt 14*, 2489–2500. [[CrossRef](#)]
20. Stanley, C.E.; Mauss, A.S.; Borst, A.; Cooper, R.L. The effects of chloride flux on *Drosophila* heart rate. *Methods Protoc.* **2019**, *2*, 73. [[CrossRef](#)]
21. Ikeda, K.; Ozawa, S.; Hagiwara, S. Synaptic transmission reversibly conditioned by single-gene mutation in *Drosophila melanogaster*. *Nature* **1976**, *259*, 489–491. [[CrossRef](#)]
22. Salkoff, L.B.; Wyman, R.J. Ion currents in *Drosophila* flight muscles. *J. Physiol.* **1983**, *337*, 687–709. [[CrossRef](#)]
23. Stewart, B.A.; Atwood, H.L.; Renger, J.J.; Wang, J.; Wu, C.F. Improved stability of *Drosophila* larval neuromuscular preparations in haemolymph-like physiological solutions. *J. Comp. Physiol. A* **1994**, *175*, 179–191. [[CrossRef](#)]
24. Budnik, V.; Koh, Y.H.; Guan, B.; Hartmann, B.; Hough, C.; Woods, D.; Gorczyca, M. Regulation of synapse structure and function by the *Drosophila* tumor suppressor gene *dlg*. *Neuron* **1996**, *17*, 627–640. [[CrossRef](#)] [[PubMed](#)]
25. Sulkowski, M.; Kim, Y.J.; Serpe, M. Postsynaptic glutamate receptors regulate local BMP signaling at the *Drosophila* neuromuscular junction. *Development* **2014**, *141*, 436–447. [[CrossRef](#)] [[PubMed](#)]
26. Elliott, E.R.; Cooper, R.L. Fluoxetine antagonizes the acute response of LPS: Blocks K₂P channels. In revision. *Comp. Biochem. Physiol. C* **2024**, (in press). [[CrossRef](#)] [[PubMed](#)]
27. de Castro, C.; Titlow, J.; Majeed, Z.R.; Cooper, R.L. Analysis of various physiological salines for heart rate, CNS function, and synaptic transmission at neuromuscular junctions in *Drosophila melanogaster* larvae. *J. Comp. Physiol. A Neuroethol. Sens. Neural. Behav. Physiol.* **2014**, *200*, 83–92. [[CrossRef](#)]
28. Badre, N.H.; Martin, M.E.; Cooper, R.L. The physiological and behavioral effects of carbon dioxide on *Drosophila melanogaster* larvae. *Comp. Biochem. Physiol. A Mol. Integr. Physiol.* **2005**, *140*, 363–376. [[CrossRef](#)]
29. Bustos, D.; Bedoya, M.; Ramírez, D.; Concha, G.; Zúñiga, L.; Decher, N.; Hernández-Rodríguez, E.W.; Sepúlveda, F.V.; Martínez, L.; González, W. Elucidating the Structural Basis of the Intracellular pH Sensing Mechanism of TASK-2 K₂P Channels. *Int. J. Mol. Sci.* **2020**, *21*, 532. [[CrossRef](#)] [[PubMed](#)]
30. Levaillant, L.; Linglart, A.; Gajdos, V.; Benachi, A.; Souberbielle, J.C. Reference values for serum calcium in neonates should be established in a population of vitamin D-replete subjects. *J. Clin. Endocrinol. Metab.* **2024**, dgae167. [[CrossRef](#)]
31. Goldstein, D.A. Serum Calcium. In *Clinical Methods: The History, Physical, and Laboratory Examinations*, 3rd ed.; Walker, H.K., Hall, W.D., Hurst, J.W., Eds.; Butterworths: Boston, MA, USA, 1990; Chapter 143. Available online: <https://www.ncbi.nlm.nih.gov/books/NBK250/> (accessed on 26 June 2024).
32. Forsberg, M.; Seth, H.; Björefeldt, A.; Lyckenvik, T.; Andersson, M.; Wasling, P.; Zetterberg, H.; Hanse, E. Ionized calcium in human cerebrospinal fluid and its influence on intrinsic and synaptic excitability of hippocampal pyramidal neurons in the rat. *J. Neurochem.* **2019**, *149*, 452–470. [[CrossRef](#)]
33. Hunter, G.; Smith, H.V. Calcium and magnesium in human cerebrospinal fluid. *Nature* **1960**, *186*, 161–162. [[CrossRef](#)]

34. Zhu, Y.C.; Cooper, R.L. Cold exposure effects on cardiac function and synaptic transmission at the neuromuscular junction in invertebrates. *Int. J. Zool. Res.* **2018**, *14*, 49–60. [[CrossRef](#)]
35. Elliott, E.R.; Brock, K.E.; Vacassenno, R.M.; Harrison, D.A.; Cooper, R.L. The effects of doxapram and its potential interactions with K2P channels in experimental model preparations. *J. Comp. Physiol. A Neuroethol. Sens. Neural. Behav. Physiol.* **2024**. [[CrossRef](#)] [[PubMed](#)]
36. Xie, J.; Ke, M.; Xu, L.; Lin, S.; Huang, J.; Zhang, J.; Yang, F.; Wu, J.; Yan, Z. Structure of the human sodium leak channel NALCN in complex with FAM155A. *Nat. Commun.* **2020**, *11*, 5831. [[CrossRef](#)] [[PubMed](#)] [[PubMed Central](#)]
37. De Castro, C.; Titlow, J.S.; Majeed, Z.R.; Malloy, C.; King, K.E.; Cooper, R.L. Chemical and mechanical factors required for maintaining cardiac rhythm in *Drosophila melanogaster* larva. *J. Entomol.* **2019**, *16*, 62–73. [[CrossRef](#)]
38. Cheslock, A.; Andersen, M.K.; MacMillan, H.A. Thermal acclimation alters Na⁺/K⁺-ATPase activity in a tissue-specific manner in *Drosophila melanogaster*. *Comp. Biochem. Physiol. A Mol. Integr. Physiol.* **2021**, *256*, 110934. [[CrossRef](#)]
39. Carvacho, I.; Gonzalez, W.; Torres, Y.P.; Brauchi, S.; Alvarez, O.; Gonzalez-Nilo, F.D.; Latorre, R. Intrinsic electrostatic potential in the BK channel pore: Role in determining single channel conductance and block. *J. Gen. Physiol.* **2008**, *131*, 147–161. [[CrossRef](#)]
40. Guo, R.; Zeng, W.; Cui, H.; Chen, L.; Ye, S. Ionic interactions of Ba²⁺ blockades in the MthK K⁺ channel. *J. Gen. Physiol.* **2014**, *144*, 193–200. [[CrossRef](#)]

Disclaimer/Publisher’s Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.