

Table S2. Pollen viability, germinability, and stigma receptivity (duration in hours) of *Turnera ulmifolia*.

Season	Pollen viability (%)		Pollen germinability (%)	Stigma receptivity (h)
	TTC	IKI		
Summer	77.92 ± 6.31	84.88 ± 6.26	73.34 ± 6.29	5.88 ± 0.34
Monsoon	78.21 ± 6.08	85.45 ± 6.27	74.17 ± 6.15	6.12 ± 0.34
Autumn	78.04 ± 6.33	85.12 ± 6.31	73.81 ± 6.43	7.38 ± 0.50
Late autumn	77.76 ± 6.06	84.57 ± 6.62	72.62 ± 5.92	8.44 ± 0.51
Winter	75.87 ± 6.28	82.62 ± 6.46	70.28 ± 5.94	9.44 ± 0.51
Spring	77.56 ± 6.07	84.32 ± 6.67	72.40 ± 6.15	6.44 ± 0.51
Throughout year	77.56 ± 6.08	84.49 ± 6.33	72.77 ± 6.14	7.24 ± 1.37
Statistical analysis	$\chi^2 = 1.94$, df = 5, p = 0.86	$\chi^2 = 2.20$, df = 5, p = 0.82	$\chi^2 = 1.87$, df = 5, p = 0.88	$\chi^2 = 84.32$, df = 5, p < 0.001