

Quantitative Assessment of Occipital Metabolic and Energetic Changes in Parkinson's Patients using *In Vivo* ³¹P MRS-based Metabolic Imaging at 7T

¹Xiao-Hong Zhu*, ¹Byeong-Yeul Lee, ²Paul Tuite, ³Lisa Coles, ³Abhishek G. Sathe, ⁴Chi Chen, ³Jim Cloyd, ⁵Walter C. Low, ⁶Clifford J. Steer and ¹Wei Chen*

¹Center for Magnetic Resonance Research, Department of Radiology; ²Department of Neurology; ³Department of Experimental and Clinical Pharmacology; ⁴Department of Food Science and Nutrition; ⁵Department of Neurosurgery; ⁶Departments of Medicine and Genetics, Cell Biology and Development; University of Minnesota, Minneapolis, MN, USA

*Correspondence: Xiao-Hong Zhu, PhD
 or Wei Chen, PhD

Supplemental Material (four figures):




Figure S1: Distribution of the intracellular NAD⁺ (A), NADH (B) and total NAD (C) contents, as well as the NAD⁺/NADH redox ratio (D) determined in individual PD patients (n=8) and healthy controls (n=8) of **Cohort I**. *** p<0.001 indicates that significant differences between the PD and CT groups were detected with 2-tailed Student t-test.




Figure S2: Distribution of the forward reaction rate constant of $k_{f,ATPase}$ (A) and $k_{f,CK}$ (B) and cerebral metabolic rate of ATP production CMR_{ATP} (C) and CMR_{CK} (D) in individual PD patients (n=11) and healthy controls (n=11) of **Cohort II**. * $p < 0.02$ indicates that significant differences between the PD and CT groups were detected with 2-tailed Student t-test.




Figure S3: Summary of phosphorous metabolites concentration (A), intracellular pH (B), forward rate constant (C) and cerebral metabolic rate of ATPase (D) and CK (E) reactions measured in a subset of the *Cohort II-PD* brains (n=3) before and after 6-week of oral UDCA treatment. All data are presented as Mean \pm SD.




Figure S4: (A) Bioenergetic changes between the PD (n=11) and control (n=11) groups of **Cohort II**, and (B) the pre- and post-UDCA conditions in *subset of Cohort II-PD* (n=3), represented by the *PD/CT* or *pre-/post-UDCA ratios* of metabolites concentration, forward rate constant and metabolic rate of ATPase and CK reactions, respectively. The red and blue arrows indicate the elevated and decreased values of the particular metabolic parameter of interest, while the dashed lines mark unchanged parameter values between the two groups or conditions.