

Article

Gauge Couplings Evolution from the Standard Model, through Pati–Salam Theory, into E_8 Unification of Families and Forces

Alfredo Aranda ^{1,2}, Francisco J. de Anda ³, António P. Morais ^{4,5} and Roman Pasechnik ^{6,*} ¹ Facultad de Ciencias-CUICBAS, Universidad de Colima, Colima 28045, Mexico² Dual CP Institute of High Energy Physics, Colima 28045, Mexico³ Tepatitlán's Institute for Theoretical Studies, Tepatitlán 47600, Mexico⁴ Departamento de Física, Universidade de Aveiro and CIDMA, Campus de Santiago, 3810-183 Aveiro, Portugal⁵ Theoretical Physics Department, European Organization for Nuclear Research, 1211 Geneva, Switzerland⁶ Department of Physics, Lund University, SE-223 62 Lund, Sweden

* Correspondence: roman.pasechnik@hep.lu.se

Abstract: We explore the potential of ultimate unification of the Standard Model matter and gauge sectors into a single E_8 superfield in ten dimensions via an intermediate Pati–Salam gauge theory. Through a consistent realisation of a $\mathbb{T}^6 / (\mathbb{Z}_6 \times \mathbb{Z}_2 \times \mathbb{Z}_2)$ orbifolding procedure and renormalisation group evolution of gauge couplings, we establish several novel benchmark scenarios for New Physics that are worth further phenomenological exploration.

Keywords: grand unified theories; supersymmetry; phenomenology of new physics

1. Introduction

Grand Unified Theories (GUTs) aim to unify the three independent gauge interactions of the Standard Model (SM) into a single one. Among the simplest ways to achieve this is via an $SU(5)$ gauge symmetry [1]. One can also extend the gauge symmetry to unify all the SM fermions of a single family into one $SO(10)$ representation [2]. Enlarging the gauge group even further into E_6 , one could unify the SM Higgs sector and a full family of fermions into a single representation [3] by means of the simple $\mathcal{N} = 1$ supersymmetry (SUSY).

Such a step-by-step enlarging of the gauge symmetry group can be studied through its Dynkin diagram and is known to follow the exceptional chain [4,5],

$$SU(3)_C \times SU(2)_L \times U(1)_Y \subset SU(5) \subset SO(10) \subset E_6 \subset E_7 \subset E_8. \quad (1)$$

Note the group E_8 is the largest group of the chain and is especially relevant as its adjoint representation (248) is the same as the fundamental one [6]. This suggests that the SM gauge fields can be in principle unified with the SM fermions provided that a maximal $\mathcal{N} = 4$ SUSY is realised. Furthermore, it also provides an $SU(3)$ flavor (or family) symmetry as a cost of E_8 to E_6 reduction.

There is a plethora of models in the literature that aim to build GUTs including flavor symmetries [7–18,18–20] and extra dimensions (EDs) [21–31]). In order to achieve viability, most require a number of independent groups and quite a large number of fields. As alluded above, the group E_8 seems to be a good bet for a complete unification of SM vectors, fermions and scalars and has indeed been widely studied both in the context of string theory [32,33] and within the framework of quantum field theory (QFT) [18,34–46].

The first challenge posed by E_8 is that it is a real group (as is extended SUSY), while the SM requires chiral representations. A way forward is to assume the existence of extra dimensions which are orbifolded in such a way that, after their compactification, the massless chiral representations containing the SM fermion sectors remain. Furthermore, extended SUSY can also be generated from the EDs [47,48]. A big challenge is to avoid the presence



Citation: Aranda, A.; Anda, F.J.d.; Morais, A.P.; Pasechnik, R. Gauge Couplings Evolution from the Standard Model, through Pati–Salam Theory, into E_8 Unification of Families and Forces. *Universe* **2023**, *9*, 90. <https://doi.org/10.3390/universe9020090>

Academic Editors: Jinmin Yang and Lorenzo Viliani

Received: 13 December 2022

Revised: 18 January 2023

Accepted: 3 February 2023

Published: 8 February 2023



Copyright: © 2023 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

of too many massless states in the low-energy limit of the theory typically originating by an orbifolding procedure of E_8 . A consistent resolution may be found by a combination of orbifolding and the Wilson line symmetry-breaking mechanism, also associated to the specific orbifold structure, providing a reduction of the symmetry group and generation of large masses for many of the unobserved states [49–51].

One of the main requirements for a consistent GUT is gauge couplings unification, i.e., when the SM gauge couplings are evolved from their measured values at the electroweak (EW) scale up to the high-energy scales, and they match into a single coupling of a unified gauge group. Recently [52,53], some of us presented a framework where the E_8 gauge symmetry with $\mathcal{N} = 1$ SUSY is considered in 10d corresponding to an extended $\mathcal{N} = 4$ SUSY in 4d, where the EDs are orbifolded so that after the compactification stage, only simple SUSY and Pati–Salam symmetry remain [54]. For an alternative example considering an E_6 SUSY GUT with 6d orbifold breaking via an intermediate Pati–Salam symmetry, see [55].

In this paper, we study for the first time a particular realisation of the GUT with E_8 gauge symmetry in 10d, where the full SM (Higgs, gauge, fermion) field content is unified into a single E_8 gauge superfield. Through the $\mathbb{T}^6/(\mathbb{Z}_6 \times \mathbb{Z}_2 \times \mathbb{Z}_2)$ orbifold, the E_8 symmetry is broken down into $SU(4)_{PS} \times SU(2)_L \times SU(2)_R \times U(1)_X \times U(1)_F \times SU(2)_{F'}$, with no remaining SUSY. In order to consistently derive the 4d Pati–Salam theory from E_8 in 10d, the ED compactification procedure invokes the presence of several extra fields for the model to be anomaly free at every step¹. Thus, we assume the existence of additional chiral superfields, consistent with the E_8 symmetry, so that the anomalies are manifestly cancelled at the Pati–Salam level. The symmetry is further broken down to the SM one through Wilson lines and VEVs of the extra fields of the model, thus leaving only the SM field content as massless.

The SM gauge and matter sectors originate from a single gauge superfield and remain massless up to the EW symmetry breaking scale. The additional chiral fields have arbitrary masses and thus may be chosen to not be present at low energies. In this work, we study different benchmark scenarios of New Physics where some of the fields survive below the compactification scale and provide important contributions to the renormalization group (RG) evolution of the gauge couplings necessary to achieve the exact gauge couplings unification below the Planck scale.

The layout of the paper is as follows: In Section 2, we show the basics of the orbifolding mechanism employed in this work. In Section 3, we demonstrate how the specific $\mathbb{T}^6/(\mathbb{Z}_6 \times \mathbb{Z}_2 \times \mathbb{Z}_2)$ orbifold breaks the E_8 gauge symmetry and SUSY in 10d. In Section 4, we introduce the extra superfields that are needed to cancel anomalies at the Pati–Salam level. In Section 5, we present the benchmark scenarios relevant for further explorations of New Physics phenomenology where the exact gauge couplings unification is achieved. Finally, in Section 6 the basic conclusions are summarised.

2. Orbifolding

Let us start by considering a theory in 10d spacetime with $\mathcal{N} = 1$ SUSY. Six extra spatial dimensions are assumed to be orbifolded as \mathbb{T}^6/F , where F is a discrete subgroup of the extra dimensional Poincarè symmetry $O(6) \ltimes T^6$, such that $O(6) \simeq SO(6) \times \mathbb{Z}_2 \simeq SU(4) \times \mathbb{Z}_2$ is the rotation group, while T^6 is the translation group. The translation group is molded by the lattice vectors $\Gamma = \mathbb{Z}^6$ compactifying it as $\mathbb{R}^6 \rightarrow \mathbb{T}^6 \simeq \mathbb{R}^6/\Gamma$ into a 6d torus. The group F must leave the lattice invariant, i.e., $F\Gamma = \Gamma$. If the discrete subgroup of the rotations is such that $F \subset SU(3)$, it would preserve simple $\mathcal{N} = 1$ SUSY, leaving an invariant $U(1)$ [56,57]. The remaining SUSY must be broken in order to generate a realistic theory. It could be broken by orbifolding through a larger discrete group $F \in SU(4)$ or through Wilson lines. In the next section it will be shown that the latter is preferable. Therefore, in the considered model, simple SUSY is preserved by the rotation subgroup F but broken by the translation one via Wilson lines at the compactification scale.

A simple orbifolding that preserves $\mathcal{N} = 1$ SUSY reads

$$F \simeq \mathbb{Z}_N \subset SU(3), \tag{2}$$

with a positive integer N , where a generic \mathbb{Z}_N orbifolding can be defined by identifying

$$\begin{aligned} (x, z_1, z_2, z_3) &\sim (x, \omega_1 z_1, \omega_2 z_2, \omega_3 z_3), & \omega_i &\equiv e^{2i\pi n_i/N}, \\ \mathcal{V}(x, z_1, z_2, z_3) &= R(\omega_1, \omega_2, \omega_3) \mathcal{V}(x, \omega_1 z_1, \omega_2 z_2, \omega_3 z_3). \end{aligned} \tag{3}$$

Here, R corresponds to a representation of the \mathbb{Z}_N rotation acting on the 10d vector superfield \mathcal{V} . Such a transformation would belong to $SU(3)$ and preserve SUSY as long as

$$n_1 + n_2 + n_3 = 0 \pmod{2N}, \tag{4}$$

as fermions rotate twice as slow [58].

Considering only an Abelian orbifolding that preserves $\mathcal{N} = 1$ SUSY, the most general one reads as $F \simeq \mathbb{Z}_N \times \mathbb{Z}_M \subset SU(3)$, where N, M are positive integers [59–61]. If the boundary condition that breaks the gauge group is imposed, the orbifolding must be accompanied by an $E_8 \supset U(1)_f \supset \mathbb{Z}_N$ transformation. As a result, the decomposed 10d superfield is transformed as follows

$$\begin{aligned} V(x, z_1, z_2, z_3) &= e^{2i\pi q_a^f/N} V(x, \omega_1 z_1, \omega_2 z_2, \omega_3 z_3), \\ \phi^i(x, z_1, z_2, z_3) &= \omega_i e^{2i\pi q_a^f/N} \phi^i(x, \omega_1 z_1, \omega_2 z_2, \omega_3 z_3), \end{aligned} \tag{5}$$

where q_a^f is the $U(1)_f$ charge of the corresponding representation. Here, V lies in the adjoint representation of the unbroken gauge group, while the light chiral superfields ϕ_i belong to the corresponding fundamental representation with charge $q_a^f = -n_i \pmod{N}$. The desired light fields are then specified by an appropriate choice of n_i . This fully determines the orbifolding procedure.

The underlined gauge symmetry can be broken and its rank reduced by adding a gauge transformation to the EDs translations through the so-called Wilson line mechanism. The latter generates a mass splitting similar to the one a Vacuum Expectation Value (VEVs) would generate. It is therefore usual to parametrize this symmetry breaking by an effective VEV. It should be remembered that it is not actually a VEV, as it does not come from the minimization of a potential, but from the ED profiles of the fields, determined by boundary conditions. Consistent with the orbifold boundary conditions, the effective VEVs should obey the rotation–translation commutation relations coming from the Poincaré algebra and hence emerge in chiral supermultiplets that have a zero mode. An effective potential for the fields is obtained by integrating out all the other fields (for more details, see, e.g., Refs. [62–67]).

The full \mathcal{R} symmetry without orbifolding is $SU(4)_{\mathcal{R}}$ of the effective $\mathcal{N} = 4$ SUSY. We can add a further orbifolding $\mathbb{Z}_2 \subset SU(4)_{\mathcal{R}}$ but $\mathbb{Z}_2 \not\subset SU(3)_{\mathcal{R}}$. Let us define S_i as the complex scalar field inside the chiral superfield ϕ_i (up to a phase $e^{i\zeta_i}$ which depends on the definition of the base) and ψ_i the chiral fermion (up to the same phase $e^{i\zeta_i}$). Let us also define as A_μ the vector field inside the gauge superfield and λ as the gaugino inside it.

The fermions transform as a 4 of $SU(4)_{\mathcal{R}}$, and they can be written as a 4 vector

$$(\psi_1, \psi_2, \psi_3, \lambda)^T, \tag{6}$$

while the real scalars transform as a 6 of $SU(4)_{\mathcal{R}}$, and it can be written as a 4×4 antisymmetric matrix

$$\begin{pmatrix} 0 & \Re S_3 & \Im S_2 & \Re S_1 \\ -\Re S_3 & 0 & \Im S_1 & \Re S_2 \\ -\Im S_2 & -\Im S_1 & 0 & \Im S_3 \\ -\Re S_1 & -\Re S_2 & -\Im S_3 & 0 \end{pmatrix}. \tag{7}$$

This would break the remaining SUSY.

3. The $T^6/(\mathbb{Z}_6 \times \mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_2)$ Orbifold with E_8

Now, we consider an E_8 gauge theory in 10d spacetime. The E_8 gauge symmetry has rank 8, and the orbifolding must preserve the rank-4 SM gauge symmetry, $SU(3)_C \times SU(2)_L \times U(1)_Y$. In this work, we employ the following decomposition [6]

$$\begin{aligned} E_8 &\supset SO(10) \times U(1)_{X'} \times SU(3)_F \\ &\supset SO(10) \times U(1)_{X'} \times U(1)_F \times SU(2)_F \\ &\supset G_{PS} \equiv SU(4)_{PS} \times SU(2)_L \times SU(2)_R \times U(1)_{X'} \times U(1)_F \times SU(2)_F, \end{aligned} \tag{8}$$

and consider the $T^6/(\mathbb{Z}_6 \times \mathbb{Z}_2)$ orbifold whose compactification triggers the breaking $E_8 \rightarrow G_{PS}$, i.e., featuring an intermediate Pati–Salam SUSY theory and a flavor $U(1)_F \times SU(2)_F$ symmetry in 4d. Subsequent reduction of this symmetry occurs in the following steps,

$$\begin{aligned} SU(4)_{PS} &\rightarrow SU(3)_C \times U(1)_{B-L}, \quad SU(2)_R \rightarrow U(1)_{T_3^R}, \\ SU(4)_{PS} \times SU(2)_R &\rightarrow SU(3)_C \times U(1)_Y \times U(1)_X, \end{aligned} \tag{9}$$

with

$$q_Y = 6q_{T_3^R} + 3q_{B-L}, \quad q_X = 4Q_{T_3^R} - 3q_{B-L}, \tag{10}$$

where one recovers the color $SU(3)_C$ and hypercharge $U(1)_Y$ groups of the SM.

The orbifold boundary conditions which provide such a breaking pattern read

$$\mathbb{Z}_6 : \phi \rightarrow e^{2i\pi q_{X'}/6} \phi, \quad \mathbb{Z}_2 : \phi \rightarrow e^{2i\pi(q_Y + q_{T_3^R})/2} \phi, \tag{11}$$

with the orbifold transformations

$$\begin{aligned} (x, z_1, z_2, z_3) &\sim (x, \alpha^2 z_1, \alpha^5 z_2, \alpha^5 z_3), \\ (x, z_1, z_2, z_3) &\sim (x, -z_1, -z_2, (-1)^2 z_3), \end{aligned} \tag{12}$$

where $\alpha = e^{2i\pi/6}$, and $-1 = e^{2i\pi/2}$. The breaking described in the first line of Equation (8) is achieved by the boundary condition given by $q_{X'}/6$. It is further broken to the second line by the boundary condition $q_{T_3^R}/2$. The third line is achieved by further applying $q_Y/2$. The above conditions enable us to decompose the **248** representation of E_8 into the representations of unbroken G_{PS} symmetry summarised in Table 1, together with the corresponding charges ² (for a better presentation we color code the representations containing **SM fermions**, **right handed neutrinos**, **SM Higgses**, **gauge fields in the adjoint**, **mirror fermions**, **mirror Higgses**, **flavons**, leptoquarks and vector-like triplets).

Table 1. The $\mathbb{Z}_6 \times \mathbb{Z}_2$ orbifold charges of each $G_{PS} \mathcal{N} = 1$ superfield. Only the superfields with both charges equal to unity (the singlets 1, 1) have zero modes. The colors indicate where the different fields of interest are contained: **SM fermions**, **right handed neutrinos**, **SM Higgses**, **gauge fields in the adjoint**, **mirror fermions**, **mirror Higgses**, **flavons**, leptoquarks and vector-like triplets.

	V	ϕ_1	ϕ_2	ϕ_3
$\mathcal{V}_{(15,1,1,0,0,1)}$	1, 1	$\alpha^2, -1$	$\alpha^5, -1$	$\alpha^5, 1$
$\mathcal{V}_{(1,3,1,0,0,1)}$	1, 1	$\alpha^2, -1$	$\alpha^5, -1$	$\alpha^5, 1$
$\mathcal{V}_{(1,1,3,0,0,1)}$	1, 1	$\alpha^2, -1$	$\alpha^5, -1$	$\alpha^5, 1$
$\mathcal{V}_{(1,1,1,0,0,1)}$	1, 1	$\alpha^2, -1$	$\alpha^5, -1$	$\alpha^5, 1$
$\mathcal{V}_{(1,1,1,0,0,3)}$	1, 1	$\alpha^2, -1$	$\alpha^5, -1$	$\alpha^5, 1$
$\mathcal{V}_{(1,1,1,0,0,1)}$	1, 1	$\alpha^2, -1$	$\alpha^5, -1$	$\alpha^5, 1$
$\mathcal{V}_{(1,1,1,0,-3,2)}$	1, -1	$\alpha^2, 1$	$\alpha^5, 1$	$\alpha^5, -1$
$\mathcal{V}_{(1,1,1,3,0,2)}$	1, -1	$\alpha^2, 1$	$\alpha^5, 1$	$\alpha^5, -1$

Table 1. Cont.

	V	ϕ_1	ϕ_2	ϕ_3
$\mathcal{V}_{(6,2,2,0,0,1)}$	1, −1	$\alpha^2, 1$	$\alpha^5, 1$	$\alpha^5, -1$
$\mathcal{V}_{(4,2,1,-3,0,1)}$	$\alpha^3, -1$	$\alpha^5, 1$	$\alpha^2, 1$	$\alpha^2, -1$
$\mathcal{V}_{(\bar{4},1,2,-3,0,1)}$	$\alpha^3, 1$	$\alpha^5, -1$	$\alpha^2, -1$	$\alpha^2, 1$
$\mathcal{V}_{(\bar{4},2,1,3,0,1)}$	$\alpha^3, -1$	$\alpha^5, 1$	$\alpha^2, 1$	$\alpha^2, -1$
$\mathcal{V}_{(4,1,2,3,0,1)}$	$\alpha^3, 1$	$\alpha^5, -1$	$\alpha^2, -1$	$\alpha^2, 1$
$\mathcal{V}_{(6,1,1,-2,1,2)}$	$\alpha^4, -1$	1, 1	$\alpha^3, 1$	$\alpha^3, -1$
$\mathcal{V}_{(6,1,1,-2,-2,1)}$	$\alpha^4, 1$	1, −1	$\alpha^3, -1$	$\alpha^3, 1$
$\mathcal{V}_{(6,1,1,2,-1,2)}$	$\alpha^2, -1$	$\alpha^4, 1$	$\alpha, 1$	$\alpha, -1$
$\mathcal{V}_{(6,1,1,2,2,1)}$	$\alpha^2, 1$	$\alpha^4, -1$	$\alpha, -1$	$\alpha, 1$
$\mathcal{V}_{(4,2,1,1,1,2)}$	$\alpha, 1$	$\alpha^3, -1$	1, −1	1, 1
$\mathcal{V}_{(4,2,1,1,-2,1)}$	$\alpha, -1$	$\alpha^3, 1$	1, 1	1, −1
$\mathcal{V}_{(\bar{4},1,2,1,1,2)}$	$\alpha, -1$	$\alpha^3, 1$	1, 1	1, −1
$\mathcal{V}_{(\bar{4},1,2,1,-2,1)}$	$\alpha, 1$	$\alpha^3, -1$	1, −1	1, 1
$\mathcal{V}_{(1,2,2,-2,1,2)}$	$\alpha^4, 1$	1, −1	$\alpha^3, -1$	$\alpha^3, 1$
$\mathcal{V}_{(1,2,2,-2,-2,1)}$	$\alpha^4, -1$	1, 1	$\alpha^3, 1$	$\alpha^3, -1$
$\mathcal{V}_{(1,1,1,4,1,2)}$	$\alpha^4, -1$	1, 1	$\alpha^3, 1$	$\alpha^3, -1$
$\mathcal{V}_{(1,1,1,4,-2,1)}$	$\alpha^4, 1$	1, −1	$\alpha^3, -1$	$\alpha^3, 1$
$\mathcal{V}_{(\bar{4},2,1,-1,-1,2)}$	$\alpha^5, 1$	$\alpha, -1$	$\alpha^4, -1$	$\alpha^4, 1$
$\mathcal{V}_{(\bar{4},2,1,-1,2,1)}$	$\alpha^5, -1$	$\alpha, 1$	$\alpha^4, 1$	$\alpha^4, -1$
$\mathcal{V}_{(4,1,2,-1,-1,2)}$	$\alpha^5, -1$	$\alpha, 1$	$\alpha^4, 1$	$\alpha^4, -1$
$\mathcal{V}_{(4,1,2,-1,2,1)}$	$\alpha^5, 1$	$\alpha, -1$	$\alpha^4, -1$	$\alpha^4, 1$
$\mathcal{V}_{(1,2,2,2,-1,2)}$	$\alpha^2, 1$	$\alpha^4, -1$	$\alpha, -1$	$\alpha, 1$
$\mathcal{V}_{(1,2,2,2,2,1)}$	$\alpha^2, -1$	$\alpha^4, 1$	$\alpha, 1$	$\alpha, -1$
$\mathcal{V}_{(1,1,1,-4,-1,2)}$	$\alpha^2, -1$	$\alpha^4, 1$	$\alpha, 1$	$\alpha, -1$
$\mathcal{V}_{(1,1,1,-4,2,1)}$	$\alpha^2, 1$	$\alpha^4, -1$	$\alpha, -1$	$\alpha, 1$

The zero modes have the following representations of the residual symmetry group (recall they correspond to those with 1, 1 charges in Table 1):

$$\begin{aligned}
 V_\mu &: (\mathbf{15}, \mathbf{1}, \mathbf{1}, \mathbf{0}, \mathbf{0}, \mathbf{1}) + (\mathbf{1}, \mathbf{3}, \mathbf{1}, \mathbf{0}, \mathbf{0}, \mathbf{1}) + (\mathbf{1}, \mathbf{1}, \mathbf{3}, \mathbf{0}, \mathbf{0}, \mathbf{1}) \\
 &\quad + (\mathbf{1}, \mathbf{1}, \mathbf{1}, \mathbf{0}, \mathbf{0}, \mathbf{1}) + (\mathbf{1}, \mathbf{1}, \mathbf{1}, \mathbf{0}, \mathbf{0}, \mathbf{1}) + (\mathbf{1}, \mathbf{1}, \mathbf{1}, \mathbf{0}, \mathbf{0}, \mathbf{3}), \\
 \phi_1 &: (\mathbf{6}, \mathbf{1}, \mathbf{1}, -2, \mathbf{1}, \mathbf{2}) + (\mathbf{1}, \mathbf{2}, \mathbf{2}, -2, -2, \mathbf{1}) + (\mathbf{1}, \mathbf{1}, \mathbf{1}, \mathbf{4}, \mathbf{1}, \mathbf{2}), \\
 \phi_2 &: (\mathbf{4}, \mathbf{2}, \mathbf{1}, \mathbf{1}, -2, \mathbf{1}) + (\bar{\mathbf{4}}, \mathbf{1}, \mathbf{2}, \mathbf{1}, \mathbf{1}, \mathbf{2}), \\
 \phi_3 &: (\mathbf{4}, \mathbf{2}, \mathbf{1}, \mathbf{1}, \mathbf{1}, \mathbf{2}) + (\bar{\mathbf{4}}, \mathbf{1}, \mathbf{2}, \mathbf{1}, -2, \mathbf{1}).
 \end{aligned}
 \tag{13}$$

which can be named as

$$\begin{aligned}
 V_\mu &: G_\mu + W_\mu^L + W_\mu^R + Z'_\mu + Z_\mu^F + W_\mu^F, \\
 \phi_1 &: T + h + \Phi, \\
 \phi_2 &: f + F^c, \\
 \phi_3 &: F + f^c,
 \end{aligned}
 \tag{14}$$

where uppercase letters T, F, Φ are used to denote $SU(2)_F$ doublets, and lowercase letters f, h correspond to $SU(2)_F$ singlets.

In order to further break the intermediate symmetry group down to the SM, one could use each Wilson line to give an effective VEV to the SM singlet fields. If they are not available, there are extra fields described below that can provide such breaking.

We now analyze each field to find the SM singlets. The field $T_{1,2}$ contains a color triplet and antitriplet; therefore, it has no SM singlets. The field

$$h \sim \begin{pmatrix} H_u \\ H_d \end{pmatrix}, \tag{15}$$

only contains the Higgs $SU(2)_L$ doublets and cannot obtain a Wilson line VEV. Otherwise, the electroweak symmetry would be broken at the compactification scale. The fields

$$F, f \sim (Q_{1,2,3}^r, Q_{1,2,3}^g, Q_{1,2,3}^b, L_{1,2,3}), \tag{16}$$

contain the quark and lepton $SU(2)_L$ doublets (where the r, g, b are the color indices) and cannot obtain a Wilson line VEV, as it would break the SM symmetry. Finally,

$$F^c, f^c \sim \begin{pmatrix} u_{1,2,3}^{cr} & u_{1,2,3}^{cg} & u_{1,2,3}^{cb} & \nu_{1,2,3}^c \\ d_{1,2,3}^{cr} & d_{1,2,3}^{cg} & d_{1,2,3}^{cb} & e_{1,2,3}^c \end{pmatrix}, \tag{17}$$

where the $\nu_{1,2,3}^c$ are SM singlets, and the sneutrino can obtain a VEV. They are charged under $SU(4)_{PS} \times SU(2)_R \times U(1)_{X'} \times U(1)_F \times SU(2)_F$ so their VEVs break it. A VEV in a 4 dimensional representation of $SU(4)_{PS}$ would break $SU(4)_{PS} \rightarrow SU(3)_C$, while a VEV in a 2 dimensional representation of $SU(2)_R$ would break it completely. Together, both sneutrino VEVs break $SU(4)_{PS} \times SU(2)_R \times U(1)_{X'} \times U(1)_F \times SU(2)_F \rightarrow SU(3)_C \times SU(2)_L \times U(1)_Y \times U(1)_N$, where it preserves an extra $U(1)_N$ whose charge is defined by

$$q_N = q_X + 5q_{X'}. \tag{18}$$

The field $\Phi \sim \varphi_{1,2}$ contains two SM singlets and can get a Wilson line VEV. Note that the VEV $\langle \Phi \rangle = (\langle \varphi_2 \rangle, \langle \varphi_3 \rangle)$ has a non-zero q_N charge and hence breaks the remaining $U(1)_N$ symmetry yielding the SM gauge symmetry at low energies.

Note that the symmetry breaking is triggered by the geometry of the orbifold. This means that the original symmetries (both the E_8 and SUSY) are not exact as they are broken explicitly on the singular points of the orbifold. Therefore, after compactification, all effective calculations are performed without SUSY and E_8 .

In this model, we can impose an extra rotational \mathbb{Z}_2 boundary condition

$$P_{SS} = \text{diag}(-1, 1, 1, -1), \tag{19}$$

which would only leave as massless modes the gauge vectors (but not the gauginos), the scalars of ϕ_1 (which lie in the T, h, Φ representations) and the chiral fermions $\phi_{2,3}$ (which lie in the representations f, f^c, F, F^c). Therefore, the whole SM field content lies in the zero modes. As the right-handed sneutrinos now do not have a zero mode, there can not be a Wilson line aligned with them; therefore, the Pati–Salam symmetry is unbroken. One may safely assume that the fields in Section 4 can break the remaining symmetry.

The sneutrino effective VEV does break the usual R-parity. It affects the phenomenology by generating the R-parity violating (RPV) terms $\sim \langle \tilde{\nu} \rangle \tilde{H}_u L$. The latter can be effectively “rotated” away by a unitary transformation which then generates terms such as $\sim y_{\text{RPV}}^{ijk} Q_i L_j d_k^c$. The strongest experimental bound on this type of interactions implies that $y_{\text{RPV}}^{111} < 0.001$ [68,69]. As such, an interaction term is proportional to the first-family neutrino Yukawa couplin; it is expected to be well below this bound.

4. Anomaly-Cancelling Sector

In the previous section, we have shown that the orbifold can break E_8 into the SM directly, leaving no remaining SUSY. We have also shown that the full SM field content arises from the single gauge supermultiplet. Therefore, the full low energy phenomenology is determined by the orbifold compactification. However, one has to ensure that any

potential anomalies are cancelled at every symmetry breaking step. While this may require the addition of new fields, we will assume that they do not survive at low energies, so that the predictivity shown in the previous section is not spoiled. Let us discuss the anomaly cancellation in more detail.

The rotation boundary conditions split the masses of different representations by integer multiples of the compactification scale $n\Lambda$. The Wilson lines split the masses by an effective VEV with scale $y\Lambda$, where y is an arbitrary real parameter. The latter can, in principle, be as small as desired, generating a certain hierarchy between the scales. Therefore, one could think of the subsequential breaking $E_8 \rightarrow G_{PS} \rightarrow G_{SM}$ at well-separated energy scales as a New Physics scenario potentially relevant for phenomenology.

The field content in the Pati–Salam phase would be composed of zero modes from Equation (13). However, we can easily notice that this phase is not fully consistent as the field content generates gauge anomalies, while the SM phase is anomaly free. In order to resolve this issue, one can add more fields so that the considered two-step breaking is consistently realised in an anomaly-free way.

There are two sets of fields that cancel the anomalies generated by the zero modes. First, one would add 27 copies of the pair of representations $(\mathbf{1}, \mathbf{1}, \mathbf{1}, 0, -1, \mathbf{2}) + (\mathbf{1}, \mathbf{1}, \mathbf{1}, 0, \mathbf{2}, \mathbf{1})$ as the 4d chiral superfields located at the $z_i = 0$ origin brane. These would cancel the flavor-specific anomalies. Note that they are all SM singlets and do not affect the low energy phenomenology. The second set of fields consists of 5 copies of 10d chiral superfields in the bulk living in the **248** representation of E_8 . They have the charges under the orbifold rotations $(1, \pm 1), (\alpha^4, \pm 1), (\alpha^2, -1)$. These will add a bunch of zero-mode chiral superfields (51 in total) which are either the real representations (i.e., vector-like pairs) or the chiral representations $(\mathbf{6}, \mathbf{1}, \mathbf{1}, -2, -2, \mathbf{1}) + (\mathbf{1}, \mathbf{2}, \mathbf{2}, -2, -2, \mathbf{1}) + (\mathbf{1}, \mathbf{1}, \mathbf{1}, \mathbf{4}, -2, \mathbf{1})$, as summarised in Table 2.

The full SM field content comes from the gauge superfield. While one can add a symmetry to restrict the couplings, they will not change the couplings to the gauge superfield. This does not reduce the predictivity of the setup. The added fields will have arbitrary couplings, and since they live in real representations, they also have explicit arbitrary masses. The extra fields in the bulk have real representations and therefore allow explicit mass terms. They are naturally heavy, and no hierarchies between them will be assumed; thus they do not affect the low energy physics.

Table 2. The $\mathbb{Z}_6 \times \mathbb{Z}_2$ orbifold charges of each G_{PS} $\mathcal{N} = 1$ superfield. Only the superfields with highlighted charges have zero modes from the 10d chiral superfields. The same colored highlighted fields would come from the same 10d chiral superfield.

	ϕ_0	ϕ_1	ϕ_2	ϕ_3
$\Phi_{(\mathbf{15}, \mathbf{1}, \mathbf{1}, 0, \mathbf{0}, \mathbf{1})}$	1, 1	$\alpha^2, -1$	$\alpha^5, -1$	$\alpha^5, 1$
$\Phi_{(\mathbf{1}, \mathbf{3}, \mathbf{1}, 0, \mathbf{0}, \mathbf{1})}$	1, 1	$\alpha^2, -1$	$\alpha^5, -1$	$\alpha^5, 1$
$\Phi_{(\mathbf{1}, \mathbf{1}, \mathbf{3}, 0, \mathbf{0}, \mathbf{1})}$	1, 1	$\alpha^2, -1$	$\alpha^5, -1$	$\alpha^5, 1$
$\Phi_{(\mathbf{1}, \mathbf{1}, \mathbf{1}, 0, \mathbf{0}, \mathbf{1})}$	1, 1	$\alpha^2, -1$	$\alpha^5, -1$	$\alpha^5, 1$
$\Phi_{(\mathbf{1}, \mathbf{1}, \mathbf{1}, 0, \mathbf{0}, \mathbf{3})}$	1, 1	$\alpha^2, -1$	$\alpha^5, -1$	$\alpha^5, 1$
$\Phi_{(\mathbf{1}, \mathbf{1}, \mathbf{1}, 0, \mathbf{0}, \mathbf{1})}$	1, 1	$\alpha^2, -1$	$\alpha^5, -1$	$\alpha^5, 1$
$\Phi_{(\mathbf{1}, \mathbf{1}, \mathbf{1}, 0, -\mathbf{3}, \mathbf{2})}$	1, -1	$\alpha^2, 1$	$\alpha^5, 1$	$\alpha^5, -1$
$\Phi_{(\mathbf{1}, \mathbf{1}, \mathbf{1}, 0, \mathbf{3}, \mathbf{2})}$	1, -1	$\alpha^2, 1$	$\alpha^5, 1$	$\alpha^5, -1$
$\Phi_{(\mathbf{6}, \mathbf{2}, \mathbf{2}, 0, \mathbf{0}, \mathbf{1})}$	1, -1	$\alpha^2, 1$	$\alpha^5, 1$	$\alpha^5, -1$
$\Phi_{(\mathbf{4}, \mathbf{2}, \mathbf{1}, -\mathbf{3}, \mathbf{0}, \mathbf{1})}$	$\alpha^3, -1$	$\alpha^5, 1$	$\alpha^2, 1$	$\alpha^2, -1$
$\Phi_{(\mathbf{4}, \mathbf{1}, \mathbf{2}, -\mathbf{3}, \mathbf{0}, \mathbf{1})}$	$\alpha^3, 1$	$\alpha^5, -1$	$\alpha^2, -1$	$\alpha^2, 1$
$\Phi_{(\mathbf{4}, \mathbf{2}, \mathbf{1}, \mathbf{3}, \mathbf{0}, \mathbf{1})}$	$\alpha^3, -1$	$\alpha^5, 1$	$\alpha^2, 1$	$\alpha^2, -1$
$\Phi_{(\mathbf{4}, \mathbf{1}, \mathbf{2}, \mathbf{3}, \mathbf{0}, \mathbf{1})}$	$\alpha^3, 1$	$\alpha^5, -1$	$\alpha^2, -1$	$\alpha^2, 1$
$\Phi_{(\mathbf{6}, \mathbf{1}, \mathbf{1}, -\mathbf{2}, \mathbf{1}, \mathbf{2})}$	$\alpha^4, -1$	1, 1	$\alpha^3, 1$	$\alpha^3, -1$
$\Phi_{(\mathbf{6}, \mathbf{1}, \mathbf{1}, -\mathbf{2}, -\mathbf{2}, \mathbf{1})}$	$\alpha^4, 1$	1, -1	$\alpha^3, -1$	$\alpha^3, 1$

Table 2. Cont.

	ϕ_0	ϕ_1	ϕ_2	ϕ_3
$\Phi_{(6,1,1,2,-1,2)}$	$\alpha^2, -1$	$\alpha^4, 1$	$\alpha, 1$	$\alpha, -1$
$\Phi_{(6,1,1,2,2,1)}$	$\alpha^2, 1$	$\alpha^4, -1$	$\alpha, -1$	$\alpha, 1$
$\Phi_{(4,2,1,1,1,2)}$	$\alpha, 1$	$\alpha^3, -1$	$1, -1$	$1, 1$
$\Phi_{(4,2,1,1,-2,1)}$	$\alpha, -1$	$\alpha^3, 1$	$1, 1$	$1, -1$
$\Phi_{(\bar{4},1,2,1,1,2)}$	$\alpha, -1$	$\alpha^3, 1$	$1, 1$	$1, -1$
$\Phi_{(\bar{4},1,2,1,-2,1)}$	$\alpha, 1$	$\alpha^3, -1$	$1, -1$	$1, 1$
$\Phi_{(1,2,2,-2,1,2)}$	$\alpha^4, 1$	$1, -1$	$\alpha^3, -1$	$\alpha^3, 1$
$\Phi_{(1,2,2,-2,-2,1)}$	$\alpha^4, -1$	$1, 1$	$\alpha^3, 1$	$\alpha^3, -1$
$\Phi_{(1,1,1,4,1,2)}$	$\alpha^4, -1$	$1, 1$	$\alpha^3, 1$	$\alpha^3, -1$
$\Phi_{(1,1,1,4,-2,1)}$	$\alpha^4, 1$	$1, -1$	$\alpha^3, -1$	$\alpha^3, 1$
$\Phi_{(\bar{4},2,1,-1,-1,2)}$	$\alpha^5, 1$	$\alpha, -1$	$\alpha^4, -1$	$\alpha^4, 1$
$\Phi_{(\bar{4},2,1,-1,2,1)}$	$\alpha^5, -1$	$\alpha, 1$	$\alpha^4, 1$	$\alpha^4, -1$
$\Phi_{(4,1,2,-1,-1,2)}$	$\alpha^5, -1$	$\alpha, 1$	$\alpha^4, 1$	$\alpha^4, -1$
$\Phi_{(4,1,2,-1,2,1)}$	$\alpha^5, 1$	$\alpha, -1$	$\alpha^4, -1$	$\alpha^4, 1$
$\Phi_{(1,2,2,2,-1,2)}$	$\alpha^2, 1$	$\alpha^4, -1$	$\alpha, -1$	$\alpha, 1$
$\Phi_{(1,2,2,2,2,1)}$	$\alpha^2, -1$	$\alpha^4, 1$	$\alpha, 1$	$\alpha, -1$
$\Phi_{(1,1,1,-4,-1,2)}$	$\alpha^2, -1$	$\alpha^4, 1$	$\alpha, 1$	$\alpha, -1$
$\Phi_{(1,1,1,-4,2,1)}$	$\alpha^2, 1$	$\alpha^4, -1$	$\alpha, -1$	$\alpha, 1$

Below the compactification scale, we have the chiral multiplets

$$\begin{aligned}
 & 2 \times \Phi_{(15,1,1,0,0,1)}, & 1 \times \Phi_{(6,2,2,0,0,1)}, & 3 \times \Phi_{(4,2,1,1,1,2)}, & 2 \times \Phi_{(\bar{4},2,1,-1,-1,2)}, \\
 & 2 \times \Phi_{(1,3,1,0,0,1)}, & 0 \times \Phi_{(4,2,1,-3,0,1)}, & 3 \times \Phi_{(4,2,1,1,-2,1)}, & 2 \times \Phi_{(\bar{4},2,1,-1,2,1)}, \\
 & 2 \times \Phi_{(1,1,3,0,0,1)}, & 0 \times \Phi_{(\bar{4},1,2,-3,0,1)}, & 3 \times \Phi_{(\bar{4},1,2,1,1,2)}, & 2 \times \Phi_{(4,1,2,-1,-1,2)}, \\
 & 2 \times \Phi_{(1,1,1,0,0,1)}, & 0 \times \Phi_{(\bar{4},2,1,3,0,1)}, & 3 \times \Phi_{(\bar{4},1,2,1,-2,1)}, & 2 \times \Phi_{(4,1,2,-1,2,1)}, \\
 & 2 \times \Phi_{(1,1,1,0,0,3)}, & 0 \times \Phi_{(4,1,2,3,0,1)}, & 2 \times \Phi_{(1,2,2,-2,1,2)}, & 1 \times \Phi_{(1,2,2,2,-1,2)}, \\
 & 2 \times \Phi_{(1,1,1,0,0,1)}, & 3 \times \Phi_{(6,1,1,-2,1,2)}, & 3 \times \Phi_{(1,2,2,-2,-2,1)}, & 2 \times \Phi_{(1,2,2,2,2,1)}, \\
 & 1 \times \Phi_{(1,1,1,0,-3,2)}, & 2 \times \Phi_{(6,1,1,-2,-2,1)}, & 3 \times \Phi_{(1,1,1,4,1,2)}, & 2 \times \Phi_{(1,1,1,-4,-1,2)}, \\
 & 1 \times \Phi_{(1,1,1,0,3,2)}, & 2 \times \Phi_{(6,1,1,2,-1,2)}, & 2 \times \Phi_{(1,1,1,4,-2,1)}, & 1 \times \Phi_{(1,1,1,-4,2,1)}, \\
 & & 1 \times \Phi_{(6,1,1,2,2,1)}, & 27 \times \Phi_{(1,1,1,0,-1,2)}, & 27 \times \Phi_{(1,1,1,0,2,1)},
 \end{aligned} \tag{20}$$

where all the vector-like pairs have an arbitrary mass determined by the abovementioned parameters. The unpaired fields contain the SM sectors plus SM singlets as we further discuss below in Section 5. While the masses of the KK modes are fixed by the compactification scale, the ones in Equation (20) are determined by the arbitrary parameters of the superpotential. Furthermore, a Wilson line VEV does not preserve SUSY; therefore, the chiral multiplets will be split into their scalar and fermion components.

For completeness, let us also show the branching rules of the Pati–Salam blue representations in terms of the SM gauge group identifying them with standard chiral matter and right-handed neutrinos,

$$\begin{aligned}
 (4, 2, 1, 1, 1, 2) & \rightarrow (3, 2, -\frac{1}{6}) + (1, 2, \frac{1}{2}) \equiv Q_{1,2} + L_{1,2} \\
 (4, 2, 1, 1, -2, 1) & \rightarrow (3, 2, -\frac{1}{6}) + (1, 2, \frac{1}{2}) \equiv Q_3 + L_3 \\
 (\bar{4}, 1, 2, 1, 1, 2) & \rightarrow (\bar{3}, 1, -\frac{2}{3}) + (\bar{3}, 1, \frac{1}{3}) + (1, 1, 1) + (1, 1, 0) \equiv u_{1,2}^c + d_{1,2}^c + e_{1,2}^c + \nu_{1,2}^c \\
 (\bar{4}, 1, 2, 1, -2, 1) & \rightarrow (\bar{3}, 1, -\frac{2}{3}) + (\bar{3}, 1, \frac{1}{3}) + (1, 1, 1) + (1, 1, 0) \equiv u_3^c + d_3^c + e_3^c + \nu_3^c
 \end{aligned} \tag{21}$$

with the labels 1, 2, 3 denoting the three families.

5. RG Evolution of Gauge Couplings

In this section, we study the RG evolution of the gauge couplings in our model with the purpose of finding possible low-energy scale scenarios compatible with an exact unification of all forces, including flavour, into E_8 . Our strategy for the current anal-

ysis consists of searching for those extensions of the SM with a minimal field content. Note that for a consistent SM-like fermion mass spectrum, one needs two Higgs doublets. The need for two Higgs doublets comes from the holomorphicity of the superpotential when there is SUSY. In this model, SUSY is broken through a non-vanishing $\langle D \rangle$ which provides specific soft SUSY breaking terms. As $\langle D \rangle$ also breaks Pati–Salam and flavor symmetry, the structure of the SUSY breaking terms is also determined by their GUT representations. These do not allow non-holomorphic Yukawa couplings; therefore, two Higgs doublets are needed. While one is responsible for giving masses to up-type quarks, the other is needed for their down-type partners. Therefore, the minimal framework to consider contains two Higgs doublets, commonly dubbed two-Higgs doublet model (2HDM) (for a detailed review, see, e.g., refs. [70–72]). The SUSY breaking mechanism, coming from the Wilson line effective VEVs, may decouple one of the Higgs doublets and create large mass difference between the two doublets. A precise analysis of their masses and their running lies beyond the scope of this paper. Due to the large amount of parameters coming from the extra fields, it will be assumed that there are scenarios where they can be tuned to bring down the scale of some extra Higgs doublets.

In what follows, we study whether a 2HDM scenario is already consistent with gauge couplings unification under E_8 , and if not, how many extra Higgs doublets or generations of vector-like fermions are needed to fulfil the unification condition. As we discuss below, we will only consider low-scale scenarios with up to a maximum of three generations of vector-like quarks (VLQ) and vector-like leptons (VLL) that can be either $SU(2)_L$ doublets or singlets. We also allow up to one additional Higgs doublet in the low-energy scale theory on top of the two doublets already mentioned above.

In this work, the unification scale is assumed to be below the compactification scale through a simplifying assumption of a small Wilson line scale. Therefore, unification of gauge couplings will be studied before the KK modes enter into effect. The running of the gauge couplings is calculated at one-loop order where the value of the inverse structure constants at a given scale μ is given by

$$\alpha_A^{-1}(\mu) = \alpha_0^{-1} + \frac{b_A}{2\pi} \log\left(\frac{\mu}{\mu_0}\right), \tag{22}$$

where a label A identifies a given gauge group and $\alpha_A = g_A^2/(4\pi)$, while α_0^{-1} denotes the value of the inverse structure constant at the initial energy-scale μ_0 . The value of the b_A coefficients will determine how fast a given gauge coupling evolves between any two scales. For non-Abelian gauge groups, these are given by

$$b_A = \frac{11}{3}C_2(G) - \frac{4}{3}\kappa T(F) - \frac{1}{3}T(S), \tag{23}$$

where $\kappa = \frac{1}{2}$ for Weyl fermions, $C_2(G)$ is a group Casimir in the adjoint representation, $T(F)$ and $T(S)$ are the Dynkin indices for fermions and complex scalars, respectively. For the case of U(1) symmetries, the beta-function coefficients read as

$$b'_A = -\frac{4}{3}\kappa \sum_f \left(\frac{Q_f}{2}\right)^2 - \frac{1}{3} \sum_s \left(\frac{Q_s}{2}\right)^2, \tag{24}$$

with Q_f and Q_s the Abelian charges of the fermions and scalars in the theory.

In our RG analysis, we consider three distinct regions:

1. The first one corresponds to the running of the gauge couplings of the G_{PS} symmetry emergent from the orbifold compactification of the 10-dimensional E_8 theory as described above. In such a region, we label the beta-function coefficients as b_A^I and denote the universal E_8 inverse structure constant at the GUT scale as α_8^{-1} . At this

stage, all representations identified in Equation (20) contribute to the b_A^I coefficients with the indicated multiplicity. Knowing that for SU(4) we have

$$T(\mathbf{15}) = C_2(\mathbf{15}) = 4, \quad T(\mathbf{6}) = 1, \quad T(\mathbf{4}) = \frac{1}{2}, \quad (25)$$

and that for SU(2)

$$T(\mathbf{3}) = C_2(\mathbf{3}) = 2, \quad T(\mathbf{2}) = \frac{1}{2}, \quad (26)$$

we obtain the coefficients

$$b_{PS}^I = -43, \quad b_L^I = b_R^I = -45, \quad b_F^I = -76. \quad (27)$$

Taking into account the $U(1)_{X'}$ and $U(1)_F$ charges and respective multiplicities also in Equation (20), the slopes of the RGEs of the Abelian inverse structure constants read as

$$b_{X'}^I = b_F^I = -234. \quad (28)$$

while $U(1)_{X'}$ and $U(1)_F$ are considered to be trivially embedded into E_8 .

- The second region corresponds to the stage after the three Wilson lines $\langle \phi_{1,2,3} \rangle$ give VEVs to the SM singlet directions. The gauge group after this stage is that of the SM, and as discussed above, we only study the following possibilities:

- A scalar sector with either two or three Higgs doublets that we denote as H in what follows;
- New exotic quarks containing either none or up to three generations of $SU(2)_L$ doublet VLQs denoted as Q_V ;
- New exotic up-type quarks containing either none or up to three generations of $SU(2)_L$ singlet VLQs and denoted as U_V ;
- New exotic down-type quarks containing either none or up to three generations of $SU(2)_L$ singlet VLQs and denoted as D_V ;
- New exotic leptons containing either none or up to three generations of $SU(2)_L$ doublet VLLs denoted as L_V ;
- New exotic leptons containing either none or up to three generations of $SU(2)_L$ singlet VLLs denoted as E_V .

Note that the choice of including up to three generations of vector-like fermions in the low-energy spectrum is not arbitrary. To see this, let us consider the possible bilinear terms involving the red and blue fields in Equation (20) that can be cast as

$$\sum_{i=1}^3 \sum_{k=1}^2 \left(\mu_{ik} F_i \bar{F}_k + \mu'_{ik} f_i \bar{f}_k + \mu''_{ik} F_i^c \bar{F}_k^c + \mu'''_{ik} f_i^c \bar{f}_k^c \right). \quad (29)$$

If we specialize on the first term, we can write a 5×5 mass matrix in the basis $\{F_1, F_2, F_3, \bar{F}_1, \bar{F}_2\}$ as

$$m_F = \begin{pmatrix} 0 & 0 & 0 & \mu_{11} & \mu_{12} \\ 0 & 0 & 0 & \mu_{21} & \mu_{22} \\ 0 & 0 & 0 & \mu_{31} & \mu_{32} \\ \mu_{11} & \mu_{21} & \mu_{31} & 0 & 0 \\ \mu_{12} & \mu_{22} & \mu_{32} & 0 & 0 \end{pmatrix}, \quad (30)$$

whose rank is 4. Furthermore, and assuming μ_{ik} real for illustration purposes, we see that $m_F^2 = m_F \cdot m_F^\dagger$ has two degenerate eigenvalues which means that in the mass basis, we end up with one massless chiral fermion F as well as two vector-like states

F_V^1 and F_V^2 . Inspired by the unified origin of our model under E_8 , in the limit where all μ_{ik} can be thought as approximately degenerate, one can write

$$\mu_{ik} = \mu + \epsilon m_{ik}, \tag{31}$$

where for $\epsilon \ll 1$ the squared masses read as

$$m_F^2 = 0 \quad m_{F_V^1}^2 = 0 + \mathcal{O}(\epsilon^2) \quad m_{F_V^2}^2 = 6\mu^2 + \mathcal{O}(\epsilon). \tag{32}$$

This implies that, for a μ of the order of the compactification scale and for sufficiently small ϵ , we can have $m_{F_V^1} \ll m_{F_V^2}$. In turn, it may result in up to two generations of vector-like fermions of the type F_V^1 not far from the TeV scale. Note that both F_i and \bar{F}_k are $SU(2)_F$ doublets and so is F_V^1 . The exact same reasoning can be applied to the μ'_{ik} , μ''_{ik} and μ'''_{ik} yielding up to two generations of the Pati–Salam fermions F_V^{1c} and up to one generation of f_V^1 and f_V^{1c} , motivating our choices in the bullet points above. Note that the $SU(2)_L$ doublet VLQs, Q_V , and VLLs L_V belong to F_V (two generations) and f_V (one generation), while their singlet counterparts, U_V , D_V and E_V are embedded in F_V^c and f_V^c . Similarly, all chiral matter belongs to the massless eigenstates and transforms according to the blue quantum numbers in Equation (20). With this in mind, the coefficients of the G_{SM} RGEs read as

$$\begin{aligned} b_C^{\text{II}} &= 11 - \frac{2}{3} \left(6 + n_{Q_V} + \frac{n_{D_V}}{2} + \frac{n_{U_V}}{2} \right), \\ b_L^{\text{II}} &= \frac{22}{3} - \frac{1}{3} \left(\frac{n_H}{2} + 1 \right) - \frac{2}{3} \left(6 + \frac{n_{L_V}}{2} + 3 \frac{n_{Q_V}}{2} \right), \\ b_Y^{\text{II}} &= -\frac{1}{3} \left(\frac{11}{4} + \frac{n_H}{16} \right) - \frac{2}{3} \left(\frac{11}{4} + \frac{n_{D_V}}{36} + \frac{n_{E_V}}{4} + \frac{n_{L_V}}{16} + \frac{n_{Q_V}}{144} + \frac{n_{U_V}}{9} \right), \end{aligned} \tag{33}$$

with the various n_X , encoding the number of extra Higgs and vector-like fermions at the low scale according to the label $X = H, Q_V, U_V, D_V, L_V, E_V$.

3. Finally, we consider a third region below the mass threshold of the vector-like fermions and where the only New Physics states are either one or two additional Higgs doublets, i.e., a 2HDM or a 3HDM EW-scale theory. Note that the presence of three Higgs doublets can be advantageous for the generation of a realistic CKM mixing in the quark sector as discussed in refs. [16,17]. With this in mind, the beta-function coefficients in this region are

$$\begin{aligned} b_C^{\text{III}} &= 7, \\ b_L^{\text{III}} &= \frac{10}{3} - \frac{1}{3} \left(\frac{n_H}{2} + 1 \right), \\ b_Y^{\text{III}} &= -\frac{11}{6} - \frac{1}{3} \left(\frac{n_H}{16} + \frac{11}{4} \right). \end{aligned} \tag{34}$$

The $U(1)_Y$ generators and the inverse structure constants matched at tree-level with those of the Pati–Salam theory read as

$$T_Y = \sqrt{\frac{2}{3}} T_{\text{PS}}^{15} - T_R^3 \quad \alpha_Y^{-1} = \frac{2}{3} \alpha_{\text{PS}}^{-1} + \alpha_R^{-1}, \tag{35}$$

resulting in the RGEs

$$\begin{aligned}
 \alpha_Y^{-1}(\mu) &= \frac{2}{3}\alpha_{PS}^{-1}(\mu_8) + \alpha_R^{-1}(\mu_8) + \left(\frac{b_{PS}^I}{3\pi} + \frac{b_R^I}{2\pi}\right) \log\left(\frac{\mu_{PS}}{\mu_8}\right) + \frac{b_Y^{II}}{2\pi} \log\left(\frac{\mu_{VLF}}{\mu_{PS}}\right) \\
 &\quad + \frac{b_Y^{III}}{2\pi} \log\left(\frac{\mu}{\mu_{VLF}}\right), \\
 \alpha_L^{-1}(\mu) &= \alpha_L^{-1}(\mu_8) + \frac{b_L^I}{2\pi} \log\left(\frac{\mu_{PS}}{\mu_8}\right) + \frac{b_L^{II}}{2\pi} \log\left(\frac{\mu_{VLF}}{\mu_{PS}}\right) + \frac{b_L^{III}}{2\pi} \log\left(\frac{\mu}{\mu_{VLF}}\right), \\
 \alpha_C^{-1}(\mu) &= \alpha_{PS}^{-1}(\mu_8) + \frac{b_{PS}^I}{2\pi} \log\left(\frac{\mu_{PS}}{\mu_8}\right) + \frac{b_C^{II}}{2\pi} \log\left(\frac{\mu}{\mu_{VLF}}\right) + \frac{b_C^{III}}{2\pi} \log\left(\frac{\mu}{\mu_{VLF}}\right),
 \end{aligned}
 \tag{36}$$

where $\mu_8 \equiv \Lambda$ is the orbifold compactification scale, μ_{PS} represents the Pati–Salam breaking scale via the Wilson line mechanism whereas μ_{VLF} is the mass threshold scale at which all vector-like fermions are integrated out.

We have performed a scan over the number of vector-like fermion generations such that $0 \leq n_X \leq 3$ for $X = Q_V, U_V, D_V, L_V, E_V$ and additional Higgs doublets $0 \leq n_H \leq 1$, i.e., considering a scalar $(n_H + 2)$ HDM sector. In addition, we have selected two possible cases: one where $\mu_{VLF} = 1$ TeV and so the new VLLs and VLQs may be at the reach of the LHC (see ref. [73] for a recent study on VLLs phenomenology in the GUT context) and another where $\mu_{VLF} = 10$ TeV, which may only be accessible at future colliders. Defining as valid low-scale models (we call each set of (n_X, n_H, μ_{VLF}) a “model”) those that are compatible with an exact unification of all gauge interactions at the E_8 breaking scale μ_8 , we have found that there are only 17 such models ((n_X, n_H, μ_{VLF}) sets). Of those, 3 work for the case where $\mu_{VLF} = 1$ TeV, while only 4 do so for the case where $\mu_{VLF} = 10$ TeV. Our results are presented in Tables 3 and 4.

Table 3. Benchmark models consistent with the unification of gauge couplings at μ_8 for $\mu_{VLF} = 1$ TeV. All three viable scenarios have $n_{Q_V} = 0$ and $n_H = 1$.

Model	$\log_{10} \frac{\mu_8}{\text{GeV}}$	$\log_{10} \frac{\mu_{PS}}{\text{GeV}}$	$\alpha_8^{-1}(\mu_8)$	n_H	n_{U_V}	n_{D_V}	n_{L_V}	n_{E_V}
1	18.33	18.24	44.96	1	1	0	0	2
2	18.68	17.77	31.12	1	1	0	0	3
3	18.56	17.93	35.64	1	0	1	0	3

Table 4. Benchmark models consistent with the unification of gauge couplings at μ_8 for $\mu_{VLF} = 10$ TeV. All viable scenarios have $n_{Q_V} = 0$. Models 5 and 6 share the same field content as Models 2 and 3 in Table 3.

Model	$\log_{10} \frac{\mu_8}{\text{GeV}}$	$\log_{10} \frac{\mu_{PS}}{\text{GeV}}$	$\alpha_8^{-1}(\mu_8)$	n_H	n_{U_V}	n_{D_V}	n_{L_V}	n_{E_V}
4	19.07	18.66	41.03	0	0	1	0	1
5	18.30	17.84	38.54	1	1	0	0	3
6	18.19	17.98	42.77	1	0	1	0	3
7	18.98	18.80	41.58	0	2	0	1	3

We limit the scan to solutions where gauge coupling unification happens below the Planck scale $M_P = 1.22 \times 10^{19}$ GeV, to avoid potentially significant corrections from quantum gravity.

We have found that at least one generation of $SU(2)_L$ singlet VLQs and VLLs is needed (Model 4), while none of the viable scenarios allow for $SU(2)_L$ doublet VLQs and only one (Model 7) do so for doublet VLLs in the low-energy scale spectrum. Model 7 is also the only one with more than one generation of U_V VLQs. The compactification scale always satisfies $\mu_8 > 10^{18}$ GeV, while the Wilson line scale $\mu_{PS} > 10^{17.7}$ GeV, not too far from μ_8 . The value of the universal E_8 gauge coupling is also limited to a small range $33 \lesssim \alpha_8^{-1}(\mu_8) \lesssim 45$. Finally, we show in Figure 1 six representative examples of Models 1, 2 and 3 for the case of $\mu_{VLF} = 1$ TeV and Models 4, 6 and 7 for the case of $\mu_{VLF} = 10$ TeV,

whereas in Figure 2, we show how the gauge couplings evolve between the E_8 and the G_{PS} scales. Notice that in such a region, the only difference resides in the starting μ_8 and stopping μ_{PS} scales as shown in Tables 3 and 4, while the coefficients of the beta functions are the same for all considered models. In particular, the coefficient of the $SU(2)_L$ and $SU(2)_R$ gauge couplings are identical $b_{L,R}^I = -45$, while that of $SU(4)_{PS}$ is $b_{PS}^I = -43$, thus closely superimposed with the latter two.




Figure 1. RG evolution of the gauge couplings in the non-minimal Pati–Salam GUT for six viable low-scale models compatible with exact unification of all interactions. In (a–c), we have $\mu_{VLF} = 1$ TeV, whereas in (d–f) $\mu_{VLF} = 10$ TeV, according to Tables 3 and 4. The orange dot represents the unification point with a universal gauge coupling g_8 .




Figure 2. Running of the gauge couplings between the μ_8 and μ_{PS} scales for the case of Model 4. The only difference between all considered scenarios in Tables 3 and 4 is the initial and final scales.

6. Conclusions

In this paper, as a further development of the previous work by some of the authors [52], we have classified for the first time possible anomaly-free realisations of non-minimal Pati–Salam GUTs emerging in four dimensions and studied the corresponding gauge couplings’ unification under the E_8 symmetry in ten dimensions. The 10-dimensional theory contains one vector **248** as well as five chiral **248**-plets. The four-dimensional limit of the E_8 theory upon $\mathbb{Z}_6 \times \mathbb{Z}_2 \times \mathbb{Z}_2$ orbifold compactification contains massless zero modes that can describe all SM gauge fields, chiral matter and right-handed neutrinos. We have shown that in the limit of approximately degenerate superpotential mass parameters, our Pati–Salam GUT can naturally contain vector-like fermions at low-energy scales and at the reach of LHC or future collider experiments. In particular, the unification of the gauge couplings under E_8 requires that such exotic fermions can either be $SU(2)_L$ singlet VLQs or both singlet and doublet VLLs. In total, we have found 7 viable models with New Physics manifest at 1 TeV or at 10 TeV, making our model falsifiable at the LHC or future colliders.

Author Contributions: Conceptualization, A.A., F.J.d.A., A.P.M. and R.P.; Methodology, A.A., F.J.d.A., A.P.M. and R.P.; Validation, A.A., F.J.d.A., A.P.M. and R.P.; Formal analysis, F.J.d.A. and A.P.M.; Investigation, A.A., F.J.d.A., A.P.M. and R.P.; Writing—original draft preparation, A.A., F.J.d.A., A.P.M. and R.P.; Writing—review and editing, A.A., F.J.d.A., A.P.M. and R.P.; Supervision, A.A., F.J.d.A., A.P.M. and R.P.; Project administration, A.A., F.J.d.A., A.P.M. and R.P. All authors have read and agreed to the published version of the manuscript.

Funding: A.A. is supported by CONACYT-SNI (México). A.P.M. is supported by the Center for Research and Development in Mathematics and Applications (CIDMA) through the Portuguese Foundation for Science and Technology (FCT—Fundação para a Ciência e a Tecnologia), references UIDB/04106/2020 and UIDP/04106/2020. A.P.M. is also supported by the projects PTDC/FIS-PAR/31000/2017, CERN/FIS-PAR/0027/2019, CERN/FISPAR/0002/2017 and by national funds (OE), through FCT, I.P., in the scope of the framework contract foreseen in the numbers 4, 5 and 6 of the article 23, of the Decree-Law 57/2016, of August 29, changed by Law 57/2017, of July 19. R. P. is supported in part by the Swedish Research Council grants, contract numbers 621-2013-4287 and 2016-05996, as well as by the European Research Council (ERC) under the European Union’s Horizon 2020 research and innovation programme (grant agreement No 668679).

Data Availability Statement: No new data were created.

Acknowledgments: The authors want to thank Stephen F. King for thorough and insightful discussions about the problems addressed in this manuscript.

Conflicts of Interest: The authors declare no conflict of interest.

Notes

- ¹ We are indebted to Stephen F. King for important discussions on this topic at early stages of this work.
- ² The orbifold rotational conditions in Equation (11), are slightly different from the previous work in [52], which preserved the flavor symmetry $SU(3)_F$. In this work the boundary conditions only preserve $SU(2)_F \times U(1)_F$ but, as will be seen below, these allow the Wilson lines to completely break the remaining symmetry into the SM one. This was not possible in the previous setup, making this one preferable.

References

1. Georgi, H.; Glashow, S.L. Unity of All Elementary Particle Forces. *Phys. Rev. Lett.* **1974**, *32*, 438–441. [\[CrossRef\]](#)
2. Fritzsch, H.; Minkowski, P. Unified Interactions of Leptons and Hadrons. *Annals Phys.* **1975**, *93*, 193–266. [\[CrossRef\]](#)
3. King, S.F.; Moretti, S.; Nevzorov, R. Exceptional supersymmetric standard model. *Phys. Lett.* **2006**, *B634*, 278–284. [\[CrossRef\]](#)
4. Buchmuller, W.; Nappi, O. Exceptional Coset Spaces and the Spectrum of Quarks and Leptons. *Phys. Lett. B* **1985**, *163*, 161. [\[CrossRef\]](#)
5. Koca, M. Explicit Realization of E8. *Lect. Notes Phys.* **2005**, *180*, 356–359. [\[CrossRef\]](#)
6. Slansky, R. Group Theory for Unified Model Building. *Phys. Rept.* **1981**, *79*, 1–128. [\[CrossRef\]](#)
7. King, S.; Ross, G.G. Fermion masses and mixing angles from SU(3) family symmetry. *Phys. Lett. B* **2001**, *520*, 243–253. [\[CrossRef\]](#)
8. King, S. Unified Models of Neutrinos, Flavour and CP Violation. *Prog. Part. Nucl. Phys.* **2017**, *94*, 217–256. [\[CrossRef\]](#)
9. Hagedorn, C.; King, S.F.; Luhn, C. A SUSY GUT of Flavour with $S_4 \times SU(5)$ to NLO. *J. High Energy Phys.* **2010**, *06*, 48. [\[CrossRef\]](#)
10. Antusch, S.; de Medeiros Varzielas, I.; Maurer, V.; Sluka, C.; Spinrath, M. Towards predictive flavour models in SUSY SU(5) GUTs with doublet-triplet splitting. *J. High Energy Phys.* **2014**, *9*, 141. [\[CrossRef\]](#)
11. Björkeröth, F.; de Anda, F.J.; de Medeiros Varzielas, I.; King, S.F. Towards a complete $A_4 \times SU(5)$ SUSY GUT. *J. High Energy Phys.* **2015**, *6*, 141. [\[CrossRef\]](#)
12. Björkeröth, F.; de Anda, F.J.; de Medeiros Varzielas, I.; King, S.F. Towards a complete $\Delta(27) \times SO(10)$ SUSY GUT. *Phys. Rev. D* **2016**, *94*, 16006. [\[CrossRef\]](#)
13. Björkeröth, F.; de Anda, F.J.; King, S.F.; Perdomo, E. A natural $S_4 \times SO(10)$ model of flavour. *J. High Energy Phys.* **2017**, *10*, 148. [\[CrossRef\]](#)
14. de Anda, F.J.; King, S.F.; Perdomo, E. $SO(10) \times S_4$ grand unified theory of flavour and leptogenesis. *J. High Energy Phys.* **2017**, *12*, 075. [\[CrossRef\]](#)
15. Cárcamo Hernández, A.; Huang, D.; Kovalenko, S.; Morais, A.P.; Pasechnik, R.; Schmidt, I. How low-scale trinification sheds light in the flavor hierarchies, neutrino puzzle, dark matter, and leptogenesis. *Phys. Rev. D* **2020**, *102*, 95003. [\[CrossRef\]](#)
16. Morais, A.P.; Pasechnik, R.; Porod, W. Prospects for New Physics from Gauge Left-Right-Colour-Family Grand Unification. *Eur. Phys. J.* **2020**, *C80*, 1162. [\[CrossRef\]](#)
17. Morais, A.P.; Pasechnik, R.; Porod, W. Grand Unified origin of gauge interactions and families replication in the Standard Model. *Universe* **2021**, *7*, 461. [\[CrossRef\]](#)
18. Camargo-Molina, J.E.; Morais, A.P.; Ordell, A.; Pasechnik, R.; Sampaio, M.O.; Wessén, J. Reviving trinification models through an E6 -extended supersymmetric GUT. *Phys. Rev.* **2017**, *D95*, 75031. [\[CrossRef\]](#)
19. Camargo-Molina, J.E.; Morais, A.P.; Ordell, A.; Pasechnik, R.; Wessén, J. Scale hierarchies, symmetry breaking and particle spectra in SU(3)-family extended SUSY trinification. *Phys. Rev.* **2019**, *D99*, 35041. [\[CrossRef\]](#)
20. Camargo-Molina, J.E.; Morais, A.P.; Pasechnik, R.; Wessén, J. On a radiative origin of the Standard Model from Trinification. *J. High Energy Phys.* **2016**, *9*, 129. [\[CrossRef\]](#)
21. Altarelli, G.; Feruglio, F.; Hagedorn, C. A SUSY SU(5) Grand Unified Model of Tri-Bimaximal Mixing from A_4 . *J. High Energy Phys.* **2008**, *3*, 52. [\[CrossRef\]](#)
22. Burrows, T.; King, S. A(4) Family Symmetry from SU(5) SUSY GUTs in 6d. *Nucl. Phys. B* **2010**, *835*, 174–196. [\[CrossRef\]](#)
23. Burrows, T.; King, S. $A_4 \times SU(5)$ SUSY GUT of Flavour in 8d. *Nucl. Phys. B* **2011**, *842*, 107–121. [\[CrossRef\]](#)
24. de Anda, F.J.; King, S.F. An $S_4 \times SU(5)$ SUSY GUT of flavour in 6d. *J. High Energy Phys.* **2018**, *7*, 57. [\[CrossRef\]](#)
25. Altarelli, G.; Feruglio, F.; Lin, Y. Tri-bimaximal neutrino mixing from orbifolding. *Nucl. Phys. B* **2007**, *775*, 31–44. [\[CrossRef\]](#)
26. Adulpravitchai, A.; Schmidt, M.A. Flavored Orbifold GUT—An $SO(10) \times S_4$ model. *J. High Energy Phys.* **2011**, *1*, 106. [\[CrossRef\]](#)
27. Adulpravitchai, A.; Blum, A.; Lindner, M. Non-Abelian Discrete Flavor Symmetries from $T^{**2}/Z(N)$ Orbifolds. *J. High Energy Phys.* **2009**, *7*, 53. [\[CrossRef\]](#)
28. Asaka, T.; Buchmuller, W.; Covi, L. Gauge unification in six-dimensions. *Phys. Lett. B* **2001**, *523*, 199–204. [\[CrossRef\]](#)
29. de Anda, F.J.; Valle, J.W.; Vaquera-Araujo, C.A. Flavour and CP predictions from orbifold compactification. *Phys. Lett. B* **2020**, *801*, 135195. [\[CrossRef\]](#)
30. de Anda, F.J.; King, S.F.; Perdomo, E. $SU(5)$ grand unified theory with A_4 modular symmetry. *Phys. Rev. D* **2020**, *101*, 15028. [\[CrossRef\]](#)

31. de Anda, F.J.; King, S.F. $SU(3) \times SO(10)$ in 6d. *J. High Energy Phys.* **2018**, *10*, 128. [[CrossRef](#)]
32. Ibanez, L.E.; Mas, J.; Nilles, H.P.; Quevedo, F. Heterotic Strings in Symmetric and Asymmetric Orbifold Backgrounds. *Nucl. Phys. B* **1988**, *301*, 157–196. [[CrossRef](#)]
33. Parr, E.; Vaudrevange, P.K.; Wimmer, M. Predicting the orbifold origin of the MSSM. *Fortsch. Phys.* **2020**, *68*, 2000032. [[CrossRef](#)]
34. Adler, S.L. Should $E(8)$ SUSY Yang-Mills be reconsidered as a family unification model? *Phys. Lett.* **2002**, *B533*, 121–125. [[CrossRef](#)]
35. Adler, S.L. Further Thoughts on Supersymmetric $E(8)$ as a Family and Grand Unification Theory 2004. Available online: <https://arxiv.org/abs/hep-ph/0401212> (accessed on 12 December 2022).
36. Garibaldi, S. E_8 , the most exceptional group. *Bull. Am. Math. Soc.* **2016**, *53*, 643–671. [[CrossRef](#)]
37. Thomas, S. Softly Broken $N = 4$ and E_8 . *J. Phys. A* **1986**, *19*, 1141. [[CrossRef](#)]
38. Konshtein, S.; Fradkin, E. Asymptotically Supersymmetric Model of unified Interaction Based On E_8 . *Pisma Zh. Eksp. Teor. Fiz.* **1980**, *32*, 575–578. (In Russian)
39. Baaklini, N. Supersymmetric Exceptional Gauge Unification. *Phys. Rev. D* **1980**, *22*, 3118–3127. [[CrossRef](#)]
40. Baaklini, N. Supergrand Unification in E_8 . *Phys. Lett. B* **1980**, *91*, 376–378. [[CrossRef](#)]
41. Barr, S.M. E_8 family unification, mirror fermions, and new low-energy physics. *Phys. Rev. D* **1988**, *37*, 204. [[CrossRef](#)]
42. Bars, I.; Gunaydin, M. Grand Unification With the Exceptional Group E_8 . *Phys. Rev. Lett.* **1980**, *45*, 859. [[CrossRef](#)]
43. Koca, M. On Tumbling E_8 . *Phys. Lett. B* **1981**, *107*, 73–76. [[CrossRef](#)]
44. Mahapatra, S.; Deo, B. Supergravity Induced $E(8)$ Gauge Hierarchies. *Phys. Rev. D* **1988**, *38*, 3554. [[CrossRef](#)]
45. Ong, C.L. Supersymmetric Models for Quarks and Leptons With Nonlinearly Realized E_8 Symmetry. *Phys. Rev. D* **1985**, *31*, 3271. [[CrossRef](#)] [[PubMed](#)]
46. Olive, D.I.; West, P.C. The $N = 4$ Supersymmetric $E(8)$ Gauge Theory and Coset Space: Dimensional Reduction. *Nucl. Phys. B* **1983**, *217*, 248–284. [[CrossRef](#)]
47. Arkani-Hamed, N.; Gregoire, T.; Wacker, J.G. Higher dimensional supersymmetry in 4-D superspace. *J. High Energy Phys.* **2002**, *3*, 55. [[CrossRef](#)]
48. Brink, L.; Schwarz, J.H.; Scherk, J. Supersymmetric Yang-Mills Theories. *Nucl. Phys. B* **1977**, *121*, 77–92. [[CrossRef](#)]
49. Kawamura, Y. Triplet doublet splitting, proton stability and extra dimension. *Prog. Theor. Phys.* **2001**, *105*, 999–1006. [[CrossRef](#)]
50. Hall, L.J.; Nomura, Y. Gauge unification in higher dimensions. *Phys. Rev. D* **2001**, *64*, 55003. [[CrossRef](#)]
51. Hall, L.J.; Murayama, H.; Nomura, Y. Wilson lines and symmetry breaking on orbifolds. *Nucl. Phys. B* **2002**, *645*, 85–104. [[CrossRef](#)]
52. Aranda, A.; de Anda, F.J.; King, S.F. Exceptional Unification of Families and Forces. *Nucl. Phys. B* **2020**, *960*, 115209. [[CrossRef](#)]
53. Aranda, A.; de Anda, F.J. Complete E_8 Unification in 10 Dimensions *Int. J. Mod. Phys.* **2021**, *A36*, 2150112. [[CrossRef](#)]
54. Pati, J.C.; Salam, A. Unified Lepton-Hadron Symmetry and a Gauge Theory of the Basic Interactions. *Phys. Rev. D* **1973**, *8*, 1240–1251. [[CrossRef](#)]
55. Braam, F.; Knochel, A.; Reuter, J. An Exceptional SSM from E_6 Orbifold GUTs with intermediate LR symmetry. *J. High Energy Phys.* **2010**, *6*, 13. [[CrossRef](#)]
56. Dixon, L.J.; Harvey, J.A.; Vafa, C.; Witten, E. Strings on Orbifolds. *Nucl. Phys.* **1985**, *B261*, 678–686. [[CrossRef](#)]
57. Dixon, L.J.; Harvey, J.A.; Vafa, C.; Witten, E. Strings on Orbifolds. 2. *Nucl. Phys.* **1986**, *B274*, 285–314. [[CrossRef](#)]
58. Groot Nibbelink, S.; Loukas, O.; Mütter, A.; Parr, E.; Vaudrevange, P.K. Tension Between a Vanishing Cosmological Constant and Non-Supersymmetric Heterotic Orbifolds. *Fortsch. Phys.* **2020**, *68*, 2000044. [[CrossRef](#)]
59. De Anda, F.J.; King, S.F.; Perdomo, E.; Vaudrevange, P.K. Flavour alignments from orbifolding: $SU(5) \times SU(3)$ model with $T^6/\Delta(54)$. *J. High Energy Phys.* **2019**, *12*, 55. [[CrossRef](#)]
60. Fischer, M.; Ratz, M.; Torrado, J.; Vaudrevange, P.K. Classification of symmetric toroidal orbifolds. *J. High Energy Phys.* **2013**, *1*, 84. [[CrossRef](#)]
61. Fischer, M.; Ramos-Sanchez, S.; Vaudrevange, P.K.S. Heterotic non-Abelian orbifolds. *J. High Energy Phys.* **2013**, *7*, 80. JHEP07(2013)080. [[CrossRef](#)]
62. Hosotani, Y. Dynamical Mass Generation by Compact Extra Dimensions. *Phys. Lett. B* **1983**, *126*, 309–313. [[CrossRef](#)]
63. Hosotani, Y. Dynamical Gauge Symmetry Breaking as the Casimir Effect. *Phys. Lett. B* **1983**, *129*, 193–197. [[CrossRef](#)]
64. Hosotani, Y.; Noda, S.; Takenaga, K. Dynamical gauge-Higgs unification in the electroweak theory. *Phys. Lett. B* **2005**, *607*, 276–285. [[CrossRef](#)]
65. Hosotani, Y.; Noda, S.; Takenaga, K. Dynamical gauge symmetry breaking and mass generation on the orbifold T^2/Z_2 . *Phys. Rev. D* **2004**, *69*, 125014. [[CrossRef](#)]
66. Haba, N.; Hosotani, Y.; Kawamura, Y.; Yamashita, T. Dynamical symmetry breaking in gauge Higgs unification on orbifold. *Phys. Rev. D* **2004**, *70*, 15010. [[CrossRef](#)]
67. Haba, N.; Harada, M.; Hosotani, Y.; Kawamura, Y. Dynamical rearrangement of gauge symmetry on the orbifold $S^1/Z(2)$. *Nucl. Phys. B* **2003**, *657*, 169–213. [[CrossRef](#)]
68. Dreiner, H.K. An Introduction to explicit R-parity violation. *Adv. Ser. Direct. High Energy Phys.* **2010**, *21*, 565–583. [[CrossRef](#)]
69. Barbier, R.; et al. R-parity violating supersymmetry. *Phys. Rept.* **2005**, *420*, 1–202. [[CrossRef](#)]
70. Branco, G.C.; Lavoura, L.; Silva, J.P. CP Violation. *Int. Ser. Monogr. Phys.* **1999**, *103*, 1–536. [[CrossRef](#)]

71. Branco, G.C.; Ferreira, P.M.; Lavoura, L.; Rebelo, M.N.; Sher, M.; Silva, J.P. Theory and phenomenology of two-Higgs-doublet models. *Phys. Rept.* **2012**, *516*, 1–102. [[CrossRef](#)]
72. Ivanov, I.P. Building and testing models with extended Higgs sectors. *Prog. Part. Nucl. Phys.* **2017**, *95*, 160–208. [[CrossRef](#)]
73. Freitas, F.F.; Gonçalves, J.; Morais, A.P.; Pasechnik, R. Phenomenology of Vector-Like Leptons with Deep Learning at the Large Hadron Collider. *J. High Energy Phys.* **2021**, *01*, 076. [[CrossRef](#)]

Disclaimer/Publisher’s Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.