

## Supplementary Materials



# Co-delivery of Imiquimod and Curcumin by Nanoemugel for Improved Topical Delivery and Reduced Psoriasis-like Skin Lesions

Mohammed S. Algahtani <sup>1</sup>, Mohammad Zaki Ahmad <sup>1</sup>, Ihab Hamed Nourein <sup>2</sup> and Javed Ahmad <sup>1,\*</sup>

<sup>1</sup> Department of Pharmaceutics, College of Pharmacy, Najran University, Najran 11001, Saudi Arabia; msalqahtane@nu.edu.sa (M.S.A.); zaki.manipal@gmail.com (M.Z.A.); ihab213@gmail.com (I.H.N.)

<sup>2</sup> Department of Clinical Laboratory (Histopathology and Cytology), College of Applied Medical Sciences, Najran University, Najran 11001, Saudi Arabia

\* Correspondence: jaahmed@nu.edu.sa or jahmad18@gmail.com; Tel.: +966-17542-8744



**Figure S1.** (a) Cumulative amount of IMQ permeated from IMQ-NEG, IMQ-CUR-NEG, and IME-CUR gel ( $n=3$ ). (b) Cumulative amount of CUR permeated from IMQ-CUR-NEG and IME-CUR gel ( $n = 3$ ).