

Supplementary table 1. Electrophysiological effects of vehicle (n=6) and loperamide (n=6) on the Isolated Arterially Perfused Rabbit Left Ventricular Wedge.

	Vehicle					Loperamide				
Concentrations	0.00%	0.00%	0.01%	0.03%	0.10%	0.1 μ M	0.3 μ M	1 μ M	3 μ M	10 μ M
JT-interval	1 \pm 1	2 \pm 2	2 \pm 2	3 \pm 3	3 \pm 3	3 \pm 2	6 \pm 4	3 \pm 5	-12 \pm 9*	-
rTp-Te	0 \pm 1	-1 \pm 2	-2 \pm 2	-3 \pm 4	-5 \pm 2	1 \pm 2	-6 \pm 4	-19 \pm 9*	-37 \pm 15*	-
TdP Score	0 \pm 0	0 \pm 0	0 \pm 0	0 \pm 0	-1 \pm 1	0 \pm 0	0 \pm 1	-1 \pm 0*	-1 \pm 0*	-1 \pm 0
QRS rate dependency	0 \pm 0	0 \pm 0	0 \pm 0	0 \pm 0	0 \pm 0	0 \pm 0	1 \pm 1	6 \pm 4*	30 \pm 23*	-
JTp	1 \pm 1	2 \pm 2	2 \pm 2	4 \pm 4	5 \pm 4	3 \pm 2	8 \pm 5*	8 \pm 6	-5 \pm 13	-

Data are expressed as % changes of the baseline value, and in MEAN \pm SD ; -: no measurable; * p<0.05