

***In vitro* evaluation of antidiabetic potential of *Cleistocalyx nervosum* var. *paniala* fruit extract**

Suttida Chukiatsiri^{1*}, Nattakarn Wongsrangsap¹, Siriluk Ratanabunyong¹, Kiattawee Choowongkomon¹

¹Department of Biochemistry, Faculty of Science, Kasetsart University, Bangkok, Thailand

*Correspondence to: Suttida Chukiatsiri, Department of Biochemistry, Faculty of Science, Kasetsart University, 50 Ngam Wong Wan Road, Ladyao, Chatuchak, 10900, Bangkok, Thailand. E-mail: suttida.e@ku.th




Figure S1. The cytotoxicity effect of water, ethanol and hexane extracts of *C. nervosum* at concentration of 0.5 mg/mL, 1.0 mg/mL, 1.5 mg/mL and 2.0 mg/ml on HepG2 after 72-h exposure assessed by MTT assay. Data are mean \pm SD of three independent experiments.

Cytotoxicity test L6




Figure S2. The cytotoxicity effect of water, ethanol and hexane extracts of *C. nervosum* at concentration of 0.5 mg/mL, 1.0 mg/mL, 1.5 mg/mL and 2.0 mg/ml on L6 myoblasts after 72-h exposure assessed by MTT assay. Data are mean \pm SD of three independent experiments.