

Supplementary table S2. Primers used for qRT-PCR

Gene name	Forward primer (5'-3')	Reverse primer (5'-3')
<i>qPCR-CsaRL1a</i>	F: TGGCTCTGGCAGTTGAA	R: CTAGTTCCTCCTGTAGTTTGGG
<i>qPCR-CsaRL1b</i>	F: GAAAAGGCTCTGGCTGTGTAT	R: AATTACTACTACTACCTGTGTAGT TTGG
<i>qPCR-CsaRL3a</i>	F: CTCATGTAGCTCCGGCTCT	R: GATCTGTAATTGGGAAATGGGAC AC
<i>qPCR-CsaRL3b</i>	F: GAGAGGTGGGAGAACATTGCG	R: CATTGCAGTTTCAGACACTTCAG
<i>qPCR-CsaRL3c</i>	F: ATGGCATCAAGTTCTCTCAGTT CAC	R: TCAGTGCTGCTTAAGGAACC
<i>qPCR-CsaRL4a \ CsaRL4b</i>	F: ATGGCATCGAACTCTTTCAG	R: TTAGAGATAATTACCACCACTCTG CC
<i>qPCR-CsaRL4c</i>	F: ATGGCATCAAGCTCTATGAAAT C	R: TTACCAACCTCTACCATTGCTAGC
<i>Csaβ-actin</i>	F: GCCATCTTTGATTGGAATGG	R: GGTGCCACAACCTTGATCTT