

Article

The Metrization Problem in $[0, 1]$ -Topology

Peng Chen 

College of Big Data, Fuzhou University of International Studies and Trade, Fuzhou 350202, China; chenpengbeijing@sina.com or chenpeng@ime.ac.cn

Abstract: This paper discusses the classification of fuzzy metrics based on their continuity conditions, dividing them into Erceg, Deng, Shi, and Chen metrics. It explores the relationships between these types of fuzzy metrics, concluding that a Deng metric in $[0, 1]$ -topology must also be Erceg, Chen, and Shi metrics. This paper also proves that the product of countably many Deng pseudo-metric spaces remains a Deng pseudo-metric space, and demonstrates some σ -locally finite properties of Deng metric space. Additionally, this paper constructs two interrelated mappings based on normal space and concludes that, if a $[0, 1]$ -topological space is T_1 and regular, and its topology has a σ -locally finite base, then it is Deng-metrizable, and thus Erceg-, Shi-, and Chen-metrizable as well.

Keywords: fuzzy point; $[0, 1]$ -topology; Deng pseudo-metric; σ -locally finite base; regular; T_1 -space; distance; metrizable

MSC: 54A40; 03E72; 54E35

1. Introduction

In general topology, given a topological space (X, δ) , it is natural to ask whether there is a metric for X such that δ is the metric topology. Such a metric metrizes the topological space and the space is said to be metrizable. Around the 1950s, through the efforts of R.H. Bing [1], Y.M. Smirnov and C.H. Dowker [2], J. Nagata [3], and M.H. Stone [4], this problem was satisfactorily solved and, eventually, their comprehensive proposition is called Nagata–Smirnov metrization admittedly in general topology, unquestionably, which is the most important theorem of topology. By that time, the main theory of topology had been perfected. However, scholars engaged in academic research never stopped exploring the unknown areas and sought new ways to gain a breakthrough in topological theory. In 1968, C.L. Chang [5] introduced the fuzzy set theory of Zadeh [6] into topology for the first time, which declared the birth of $[0, 1]$ -topology. Soon after that, J.A. Goguen [7] further generalized L -fuzzy set to the proposed $[0, 1]$ -topology and his theory is now recognized as L -topology. From then on, this kind of lattice-valued topology formed another important branch of topology, and thereafter many creative results and original thoughts were presented (see [8–31], etc.).

Nevertheless, how to generalize classical metrics to the lattice-valued topology reasonably has always been a great challenge. So far, there have been some fuzzy metrics introduced in the branch of learning (see [9,13,15,16,28,32–34], etc.). Considering the codomain is either an ordinary number or a fuzzy number, these metrics are roughly divided into two types.

One type is composed of these metrics, each of which is defined by such a function whose distance between objects is fuzzy, while the objects themselves are crisp. Additionally, each of them always induces a fuzzifying topology. In recent years, these metrics have been promoted by quite a few experts, such as I. Kramosil, J. Michalek, A. George, P. Veeramani, V. Gregori, S. Romaguera, J. Gutiérrez García, S. Morillas, and F.G. Shi, etc. (see [16,17,27,35–46], etc.).



Citation: Chen, P. The Metrization Problem in $[0, 1]$ -Topology. *Mathematics* **2023**, *11*, 4430. <https://doi.org/10.3390/math11214430>

Academic Editor: Gia Sirbiladze

Received: 4 September 2023

Revised: 18 October 2023

Accepted: 19 October 2023

Published: 26 October 2023



Copyright: © 2023 by the author. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

The other type consists of these metrics, each of which is defined by such a mapping $p : M \times M \rightarrow [0, +\infty)$, where M is the set of all standard fuzzy points of the underlying classical set X . In this case, every such fuzzy metric always induces a fuzzy topology (see [9,13–15,28,30], etc.).

About the latter, there are roughly three kinds of fuzzy metrics in history, with which the academic community has gradually been familiar. In addition, there is the recently discovered fourth metric. We will list the four fuzzy metrics below, one by one.

The first is the Erceg metric, which was presented in 1979 by M.A. Erceg [15]. Since then, many scholars have been engaged in this research and have obtained many beautiful results. Among them, a typical proposition can be found in [24], where J.H. Liang showed Urysohn’s metrization theorem in 1984: an L -topological space is Erceg metrizable if it is T_1 , regular, and C_{II} . In 1985, M.K. Luo [25] constructed an example of the Erceg metric on I^X whose metric topology has no σ -locally finite base, which implies that the $[0, 1]$ -topological space of this example is not C_{II} of course, but then Liang’s conclusion is still the best one. In this paper, Liang’s conclusion is only a corollary of our result in $[0, 1]$ -topology. Later on, based on Peng’s simplification method [47], the Erceg metric was further simplified by P. Chen and F.G. Shi (see [12,48]):

(I) An Erceg pseudo-metric on L^X is a mapping $p : M \times M \rightarrow [0, +\infty)$ satisfying

(A1) If $a \geq b$, then $p(a, b) = 0$;

(A2) $p(a, c) \leq p(a, b) + p(b, c)$;

(B1) $p(a, b) = \bigvee_{c \ll b} p(a, c)$;

(A3) $\forall a, b \in M, \exists x \not\leq a' \text{ s.t. } p(b, x) < r \Leftrightarrow \exists y \not\leq b' \text{ s.t. } p(a, y) < r$.

An Erceg pseudo-metric p is called an Erceg metric if it further satisfies

(A4) If $p(a, b) = 0$, then $a \geq b$,

where “ \ll ” is the way-below relation in Domain Theory and L^X is a completely distributive lattice [49,50].

The second is Shi metric (or p.q. metric), which is proposed in 1988 by L.C. Yang [30]. It was proved by Yang that each topological molecular lattice with C_{II} property is p.q.-metrizable (refer to [30,50] for details). After that, this kind of metric was studied in depth by F.G. Shi (see [12,28,48,51,52], etc.). Its definition is as follows:

(II) A Shi pseudo-metric (resp., Shi metric) on L^X is a mapping $p : M \times M \rightarrow [0, +\infty)$ satisfying (A1)–(A3) (resp., (A1)–(A4)) and the following

(B2) $p(a, b) = \bigwedge_{c \ll a} p(c, b)$.

Similarly, according to our later proofs in this paper, Yang’s proposition is still a corollary of our result in $[0, 1]$ -topology.

The third is Deng metric supplied in 1982 by Z.K. Deng [13], where Deng [14] showed that if a $[0, 1]$ -topological space is T_1 , regular and C_{II} then it is Deng metrizable. In this paper, we will extend this result substantially. Incidentally, Y.Y. Lan and F. Long also provided a result about Deng pseudo-metrization problem [53]. However, the proof was not completely right after careful checking pointed out by us. It is worth mentioning that, since Deng’s research is only limited to this special lattice I^X , Deng pseudo-metric was later extended to L^X by P. Chen [54]:

(III) An extended Deng pseudo-metric (resp., extended Deng metric) on L^X is a mapping $p : M \times M \rightarrow [0, +\infty)$ satisfying (A1)–(A3) (resp., (A1)–(A4)) and the following:

(B3) $p(a, b) = \bigwedge_{b \ll c} p(a, c)$.

In summary, the above three kinds of fuzzy metrics are defined by using (A1)–(A4) and different (B1)–(B3), respectively. Inspired by this, we conclude that there is another new metric defined as follows:

(IV) A Chen pseudo-metric (resp., Chen metric) on L^X is a mapping $p : M \times M \rightarrow [0, +\infty)$ satisfying (A1)–(A3) (resp., (A1)–(A4)) and the following:

$$(B4) \quad p(a, b) = \bigvee_{a \ll c} p(c, b).$$

Some elementary properties related to it have been introduced in [9].

In this paper, we will focus mainly on the latter and study its metrization problem in $[0, 1]$ -topology. For this reason, we investigate the relationships between (I)–(IV) on I^X and fortunately acquire such a profound result: let $C = \{ p \mid p \text{ be a Chen metric} \}$, $E = \{ p \mid p \text{ is an Erceg metric} \}$, $D = \{ p \mid p \text{ is a Deng metric} \}$ and $Y = \{ p \mid p \text{ is a Shi metric} \}$ on I^X . Then $D = C \cap Y \cap E$.

Consequently, if a given $[0, 1]$ -topology is Deng-metrizable, then it must be Erceg-, Shi-, and Chen-metrizable. Thus, this paper mainly will discuss Deng metric and its metrization problem in $[0, 1]$ -topology.

To sum up, although so many scholars have been engaged in the study of fuzzy metrics, it is a little pity that the metrization problem in $[0, 1]$ -topology remains unsolved now. This paper aims to study the metrization problem in $[0, 1]$ -topology and will obtain the generalization of Nagata–Smirnov metrization theorem in $[0, 1]$ -topology.

2. Preliminaries

In this section, we cite the fundamental definitions that will be used in the sequel. The letter X always refers to a nonempty set throughout this paper, and I denotes the unit interval $[0, 1]$.

A fuzzy set of X is a mapping $A : X \rightarrow I$, which forms the family I^X . The constant fuzzy set of X with the value 1 (resp., 0) is denoted by $\underline{1}$ (resp., $\underline{0}$). A fuzzy point (resp., standard fuzzy point) x_λ in X is a fuzzy set defined by $x_\lambda(x) = \lambda$ and $x_\lambda(y) = 0$ if $y \neq x$, where λ is a fixed number in $(0, 1)$ (resp., $(0, 1]$). The set of all fuzzy points (resp., all standard fuzzy points) of X is denoted by M_0 (resp., M). M_0 is a subfamily of M . Naturally, these properties of M_0 and M are also suitable for L -topology.

A subfamily δ of I^X is called a $[0, 1]$ -topology if it satisfies the following three conditions: (O1) $\underline{1}, \underline{0} \in \delta$; (O2) if $A, B \in \delta$, then $A \wedge B \in \delta$; (O3) if $\{A_\lambda \mid \lambda \in \Lambda\} \subseteq \delta$, then $\bigvee_{\lambda \in \Lambda} A_\lambda \in \delta$. The pair (X, δ) is called a $[0, 1]$ -topological space (a space for short). If $\delta \subseteq I^X$, then for each $A \in \delta$, A and A' are called a δ -fuzzy open set and a δ -fuzzy closed set (open set and closed set for short), respectively.

Two fuzzy sets A and B are called quasi-coincident if there exists x belonging to X such that $A(x) + B(x) > 1$ (see [55]). Let x_α be a fuzzy point and let A be a fuzzy set of X . The notation $x_\alpha \in A$ means $\alpha < A(x)$ [13]. The closure of a fuzzy set A of (X, δ) is the intersection of the members of the family of all closed sets containing A , denoted by \overline{A} [13]. A fuzzy point x_α is called a *cluster point* of a fuzzy set U of (X, δ) if each open neighborhood of $x_{1-\alpha}$ is quasi-coincident with U . Consequently, $x_\alpha \leq \overline{A}$ if and only if x_α is a cluster point of A . Therefore, $\overline{A} = \bigvee \{y_\beta \mid y_\beta \text{ is a cluster point of } A\}$ [13].

The space (X, δ) is called regular (resp., normal) if for any $x_\lambda \in M_0$ (resp., $\tau' \in \delta$), $\mu \in \delta$ with $x_\lambda \in \mu$ (resp., $\tau \leq \mu$), there exists v belonging to δ such that $x_\lambda \in v \leq \overline{v} \leq \mu$ (resp., $\tau \leq v \leq \overline{v} \leq \mu$) [21]. A $[0, 1]$ -topological space is T_1 if and only if x_λ is closed for each fuzzy point $x_\lambda \in M_0$. A family ψ of fuzzy sets is a base of δ if ψ is a subfamily of δ and for each fuzzy point x_λ and each open neighborhood μ of x_λ , there is a member v of δ such that $x_\lambda \in v \leq \mu$. A family κ of fuzzy sets is a subbase of δ if the family of finite intersections of members of κ is a base of δ [13,56]. The space (X, δ) is C_{II} , or called second-countable if the $[0, 1]$ -topology δ has a countable basis.

A family of fuzzy sets Ψ is called locally finite (resp., discrete) in a space (X, δ) if and only if each fuzzy point x_λ of the space has its an open neighborhood which is quasi-coincident with only finitely many members (resp., at most one member) of Ψ (see [50]).

A family of fuzzy sets is called σ -locally finite (resp., σ -discrete) in a space (X, δ) if and only if it is the union of a countable number of locally finite (resp., discrete) subfamilies. A subfamily σ of I^X (resp., σ of δ) is called a (resp., an open) cover of a fuzzy set A in a space (X, δ) if for each $x_\alpha \in A$, there exists B belonging to σ such that $x_\alpha \in B$. Furthermore, if $A = \underline{1}$, then σ is called a cover of (X, δ) . A cover \mathcal{B} of a fuzzy set A is called a refinement of a cover \mathcal{D} if each member of \mathcal{B} is a subset of a member of \mathcal{D} [50].

Let $\{X_t\}_{t \in T}$ be an indexed family of sets. The Cartesian product of this indexed family, denoted by $\prod_{t \in T} X_t$, is the set of all functions $x : T \rightarrow \bigcup_{t \in T} X_t$ such that $x(t) \in X_t$ for each $t \in T$.

Let $X = \prod_{t \in T} X_t$. Then, the t -th projection $J_t : I^X \rightarrow I^{X_t}$ is defined by $J_t(A)(y_t) = \sup\{A(x) \mid x_t = y_t\}$ for each $y_t \in X_t$ and let $J_t^{-1}(B) = \bigvee\{C \in I^X \mid J_t(C) \leq B\}$. The product space of $\{(X_t, \delta_t) \mid t \in T\}$ is defined by $\{J_t^{-1}(A_t) \mid A_t \in \delta_t, t \in T\}$ as a subbase [50].

Other unexplained terminologies and notations and further details can be found in [7,9,13,50,56].

Definition 1 ([9,13]). A Deng pseudo-metric on I^X is a mapping $p : M_0 \times M_0 \rightarrow [0, +\infty)$ satisfying

(D1) If $\lambda_1 \geq \lambda_0$, then $p(x_{\lambda_1}, x_{\lambda_0}) = 0$;

(D2) $p(x_{\lambda_1}, z_{\lambda_3}) \leq p(x_{\lambda_1}, y_{\lambda_2}) + p(y_{\lambda_2}, z_{\lambda_3})$;

(D3) $p(x_{\lambda_1}, y_{\lambda_2}) = \bigwedge_{\lambda > \lambda_2} p(x_{\lambda_1}, y_\lambda)$;

(D4) $p(x_{\lambda_1}, y_{\lambda_2}) = p(y_{1-\lambda_2}, x_{1-\lambda_1})$.

A Deng pseudo-metric p is called a Deng metric if it further satisfies the following:

(D5) If $p(x_{\lambda_1}, y_{\lambda_2}) = 0$, then $x = y, \lambda_1 \geq \lambda_2$.

Remark 1. In [54], we have proved the following results: (1) a Deng metric p on I^X can be extended to an extended Deng metric p^* ; (2) $p = p^* \mid M_0 \times M_0$; (3) p^* and p induce the same metric topology.

Based on the above (1)–(3), it is much easier to study the Deng metric by using Definition 1 instead of (III) on I^X as its definition. A similar treatment to (I), (II), and (VI) on I^X is to restrict their domains to M_0 and use (D4) instead of (A3) while other conditions remain unchanged.

Theorem 1 ([13]). Let p be a Deng pseudo-metric (resp., a Deng metric) on I^X . For each $r \in [0, 1)$ define $U_r(a) = \bigvee\{b \in M_0 \mid p(a, b) < r\}$. Then the family $\{U_r(a) \mid a \in M_0, r \in [0, +\infty)\}$ forms a base of δ_p , called the $[0, 1]$ -topology induced by p . The space (X, δ_p) is called a Deng pseudo-metric space (resp., a Deng metric space).

Theorem 2 ([13]). If p is a Deng pseudo-metric on I^X , then (X, δ_p) is regular, normal.

In [13], Deng has proved such a result: If (X, δ) is regular and C_{II} , then it is normal [13]. It is a special case of the following result:

Theorem 3 ([9,50]). If (X, δ) is regular, and δ has a σ -locally finite base, then it is normal.

Theorem 4 ([13,50]). If $\{A_\lambda \mid \lambda \in \Gamma\}$ is locally finite in a space (X, δ) , then $\overline{\bigvee_{\lambda \in \Gamma} A_\lambda} = \bigvee_{\lambda \in \Gamma} \overline{A_\lambda}$.

Theorem 5 ([52]). Suppose that (X, δ) is normal, and let a closed set A and an open set B satisfy $A \leq B$. Then there is a family $\{U_r \mid r \in \mathbb{Q}_{[0,1]}\}$ such that each element is an open neighborhood of A and satisfies the following properties:

- (a) $U_0 = A, U_1 = B;$
- (b) If $r < s$, then $U_r \leq \overline{U_r} \leq U_s.$

Theorem 6 ([52]). Let p be a Shi pseudo-metric on L^X and define $P_r(b) = \bigvee \{c \in M \mid p(c, b) \geq r\}$. Then for $c, b \in M, c \leq P_r(b) \Leftrightarrow p(c, b) \geq r.$

Theorem 7. Let p be a Deng pseudo-metric on I^X . For any $a \in M_0$ and each $r \in [0, 1)$ define $B_r(a) = \bigvee \{b \in M_0 \mid p(a, b) \leq r\}$. Then

- (1) $\overline{B_r(a)} = B_r(a);$
- (2) $b \leq B_r(a) \Leftrightarrow p(a, b) \leq r.$

Proof. Because of the following Theorem 9, Theorem 10, and the existing proposition [12]: If p is an Erceg pseudo-metric on I^X , then it satisfies (1) and (2). This theorem holds as desired. \square

3. The Relationships between Four Kinds of Fuzzy Metrics on I^X

In this section, we will investigate the relationships between the four kinds of metrics: Erceg, Shi, Deng, and Chen metrics. First of all, we expose the main result as follows:

Theorem 8. On I^X , let $C = \{p \mid p \text{ be a Chen metric}\}; E = \{p \mid p \text{ is an Erceg metric}\}; D = \{p \mid p \text{ is a Deng metric}\}; Y = \{p \mid p \text{ is a Shi metric}\}.$ Then $D = C \cap Y \cap E.$

Proof. It can be obtained from the following Theorem 9–12. \square

Theorem 9. If p is a Shi pseudo-metric on I^X , then it is an Erceg pseudo-metric.

Proof. To prove that p is an Erceg pseudo-metric on I^X , we only need to prove that $p(x_\alpha, y_\beta) = \bigvee_{\gamma < \beta} p(x_\alpha, y_\gamma).$ The proof is as follows:

By (A1) and (A2), when $\gamma < \beta$, we have $p(x_\alpha, y_\gamma) \leq p(x_\alpha, y_\beta).$ Hence $p(x_\alpha, y_\beta) \geq \bigvee_{\gamma < \beta} p(x_\alpha, y_\gamma).$ If $p(x_\alpha, y_\beta) > \bigvee_{\gamma < \beta} p(x_\alpha, y_\gamma),$ then we may take two different numbers $s, r > 0$ such that

$$p(x_\alpha, y_\beta) > s > r \geq \bigvee_{\gamma < \beta} p(x_\alpha, y_\gamma).$$

In addition, for each $\gamma < \beta$, by triangle inequality $p(x_\alpha, y_\beta) \leq p(x_\alpha, y_\gamma) + p(y_\gamma, y_\beta),$ we have

$$s < p(x_\alpha, y_\beta) \leq p(y_\gamma, y_\beta) + r.$$

Therefore, $p(y_\gamma, y_\beta) > s - r$, so that $p(y_\beta, y_\beta) = \bigwedge_{\gamma < \beta} p(y_\gamma, y_\beta) \geq s - r > 0.$ But this contradicts (A1), as desired. \square

However, the converse is not true. Such a counterexample is given below.

Example 1. Let $L = [0, 1]$ and $X = \{x\}.$ For convenience, we denote L^X and x_λ for L and λ respectively. Define a mapping $p : (0, 1] \times (0, 1] \rightarrow [0, +\infty)$ by:

$$p(a, b) = \begin{cases} 0, & \text{if } a \geq b; \\ 1, & \text{if } a < b. \end{cases}$$

Firstly, let us verify that p is an Erceg pseudo-metric on $[0, 1].$ (A1) and (A2) are trivial.

(B1) if $a, b \in (0, 1]$ and $a \geq b$, then $p(a, b) = 0$. Therefore, we have $\bigvee_{x < b} p(a, x) = 0$, so that in this case $p(a, b) = \bigvee_{x < b} p(a, x)$. Similarly, when $a < b$, we can prove $p(a, b) = \bigvee_{x < b} p(a, x) = 1$. Consequently, p satisfies (B1).

(A3) we only need to prove that $\bigwedge_{y > 1-b} p(a, y) = \bigwedge_{x > 1-a} p(b, x)$, which can be obtained from the following implications: $\bigwedge_{y > 1-b} p(a, y) = 1 \Leftrightarrow y > 1 - b$ implies $y > a \Leftrightarrow 1 - b \geq a \Leftrightarrow x > 1 - a$ implies $x > b \Leftrightarrow \bigwedge_{x > 1-a} p(b, x) = 1$.

Secondly, we assert that p is not a Shi pseudo-metric. In fact, for any $a \in (0, 1]$, we have $p(a, a) = 0$. But $\bigwedge_{c < a} p(c, a) = 1$. Thus $p(a, b) \neq \bigwedge_{c < a} p(c, b)$, as desired.

Theorem 10. *If p is a Deng pseudo-metric on I^X , then it is a Shi pseudo-metric.*

Proof. For any two fuzzy points x_a and y_b , we only need to prove $p(x_a, y_b) = \bigwedge_{c < a} p(x_c, y_b)$. If $c < a$, then $p(x_a, y_b) \leq p(x_c, y_b)$. So $p(x_a, y_b) \leq \bigwedge_{c < a} p(x_c, y_b)$. If $p(x_a, y_b) = r < \bigwedge_{c < a} p(x_c, y_b) = t$, then by (D4) we have $p(y_{1-b}, x_{1-a}) = r < t$, so that by (D3) there exists a number $s > 1 - a$ such that $p(y_{1-b}, x_s) < t$, i.e., $p(x_{1-s}, y_b) < t$. But that contradicts $\bigwedge_{c < a} p(x_c, y_b) = t$. Consequently, $p(x_a, y_b) = \bigwedge_{c < a} p(x_c, y_b)$, as desired. \square

Conversely, we have the following proposition:

Theorem 11. *If a Shi pseudo-metric p further is a Chen pseudo-metric on I^X , then p is a Deng pseudo-metric.*

To prove this, we first need to prove the following two Lemmas 1 and 2.

Lemma 1. *Let p be a Shi pseudo-metric on I^X and for each $r \in [0, 1)$ define $U_r(a) = \bigvee \{b \in I^X \mid p(a, b) < r\}$. Then $U_r(y_\lambda) = \bigvee_{\alpha > 1-\lambda} P_r(y_\alpha)'$.*

Proof. Let $x_\beta \in \bigvee_{\alpha > 1-\lambda} P_r(y_\alpha)'$ and take γ such that $x_\beta < x_\gamma \leq \bigvee_{\alpha > 1-\lambda} P_r(y_\alpha)'$. Because $1 - \gamma \geq \bigwedge_{\alpha > 1-\lambda} P_r(y_\alpha)(x)$, there exists a number $\alpha > 1 - \lambda$ such that $1 - \gamma \geq P_r(y_\alpha)(x)$, and then for each $\delta > 1 - \gamma$ we have $\delta > P_r(y_\alpha)(x)$. Therefore by Theorem 6, we can obtain $p(x_\delta, y_\alpha) < r$. Again by (A3) in (I) ((A3) on the special case I^X of L^X is for any $x_{\lambda_1}, y_{\lambda_2}, \exists t > 1 - \lambda_1$ s.t. $p(y_{\lambda_2}, x_t) < r \Leftrightarrow \exists s > 1 - \lambda_2$ s.t. $p(x_{\lambda_1}, y_s) < r$), there exists $x_\omega(x_\delta)$ (x_ω has something to do with x_δ) with $\omega > 1 - \delta$ such that $p(y_\lambda, x_\omega) < r$. Let $x_q = \bigvee \{x_\omega(x_\delta) \mid \delta > 1 - \gamma\}$. Then $x_\delta \not\leq x_{1-q}$, i.e., $x_\delta > x_{1-q}$. This implies that, as long as $x_\delta > x_{1-\gamma}$, it must hold that $x_\delta > x_{1-q}$. Thus $x_\gamma \leq x_q$. Since $x_\beta < x_\gamma \leq x_q$, there exists $x_\omega(x_\delta)$ such that $x_\beta \leq x_\omega$, and so $p(y_\lambda, x_\beta) \leq p(y_\lambda, x_\omega) < r$. Hence $x_\beta \leq U_r(y_\lambda)$. Because x_β is arbitrary, we have $\bigvee_{\alpha > 1-\lambda} P_r(y_\alpha)' \leq U_r(y_\lambda)$.

Conversely, let $x_\alpha \in U_r(y_\lambda)$. Then $p(y_\lambda, x_\alpha) < r$. For each $x_\beta > x_{1-\alpha}$, i.e., $\alpha > 1 - \beta$, by (A3) there exists $\gamma > 1 - \lambda$ such that $p(x_\beta, y_\gamma) < r$, and then by Theorem 6, $x_\beta \not\leq P_r(y_\gamma)$. Hence $x_\beta \not\leq \bigwedge_{\gamma > 1-\lambda} P_r(y_\gamma)$. That is to say, as long as $x_\beta > x_{1-\alpha}$, i.e., $x_\beta \not\leq x_{1-\alpha}$, it is true that $x_\beta \not\leq \bigwedge_{\gamma > 1-\lambda} P_r(y_\gamma)$. Consequently, $\bigwedge_{\gamma > 1-\lambda} P_r(y_\gamma)(x) \leq x_{1-\alpha}$, i.e., $x_\alpha \leq \bigvee_{\gamma > 1-\lambda} P_r(y_\gamma)'$. Because x_α is arbitrary, we have $U_r(y_\lambda) \leq \bigvee_{\gamma > 1-\lambda} P_r(y_\gamma)'$, as desired. \square

Lemma 2. *If p is a Shi pseudo-metric on I^X , then $\bigvee_{\alpha > 1-\lambda_1} p(x_\alpha, y_{\lambda_2}) = \bigvee_{\beta > 1-\lambda_2} p(y_{\lambda_\beta}, x_{\lambda_1})$.*

Proof. Denote $\bigvee_{\alpha > 1-\lambda_1} p(x_\alpha, y_{\lambda_2}) = \bigvee_{\beta > 1-\lambda_2} p(y_{\lambda_\beta}, x_{\lambda_1})$ as (H1). Then it is easy to check that (H1) is equivalent to the following property:

$$(H1)^* \exists \alpha > 1 - \lambda_1 \text{ s.t. } p(x_\alpha, y_{\lambda_2}) > r \Leftrightarrow \exists \beta > 1 - \lambda_2 \text{ s.t. } p(y_{\lambda_\beta}, x_{\lambda_1}) > r.$$

Now, let us prove (H1)*.

Assume that there is α with $\alpha > 1 - \lambda_1$ such that $p(x_\alpha, y_{\lambda_2}) > r$. Take a number s such that $p(x_\alpha, y_{\lambda_2}) > s > r$. By the process of proving of Theorems 7 and 9, we assert that $\lambda_2 > B_s(x_\alpha)(y)$. Therefore, by Lemma 1, we can obtain the following formula:

$$\lambda_2 > B_s(x_\alpha)(y) \geq U_s(x_\alpha)(y) = \bigvee_{\gamma > 1-\alpha} P_s(x_\gamma)'(y).$$

Thus, for every $\gamma > 1 - \alpha$ it is true that $\lambda_2 > P_s(x_\gamma)'(y)$. That is to say, as long as $\alpha > 1 - \lambda_1$, i.e., $x_{\lambda_1} \not\leq x_{1-\alpha}$ such that $p(x_\alpha, y_{\lambda_2}) > r$, it is true that $\lambda_2 > P_s(x_{\lambda_1})'(y)$, i.e., $1 - \lambda_2 < P_s(x_{\lambda_1})(y)$. So there exists y_ω such that $y_{1-\lambda_2} < y_\omega \leq P_s(x_{\lambda_1})$, and then $p(y_\omega, x_{\lambda_1}) \geq s > r$ by Theorem 6; similarly, so is the reverse, as desired. \square

Proof. The proof of Theorem 11 is as follows:

Let p be a Shi pseudo-metric on I^X and it satisfies $p(x_{\lambda_2}, y_{\lambda_1}) = \bigvee_{s > \lambda_2} p(x_s, y_{\lambda_1})$. Then we only need to prove that p satisfies (D3) and (D4).

(D4). Given any $x_{\lambda_1}, y_{\lambda_2} \in M_0$. According to Lemma 2, we have

$$\bigvee_{\alpha > 1-\lambda_1} p(x_\alpha, y_{\lambda_2}) = \bigvee_{\beta > 1-\lambda_2} p(y_{\lambda_\beta}, x_{\lambda_1}),$$

and then $p(x_{1-\lambda_1}, y_{\lambda_2}) = p(y_{1-\lambda_2}, x_{\lambda_1})$.

(D3). By (D1) and (D2), if $\lambda_3 > \lambda_1$, then $p(y_{\lambda_2}, x_{\lambda_1}) \leq p(y_{\lambda_2}, x_{\lambda_3})$. Thus, $p(y_{\lambda_2}, x_{\lambda_1}) \leq \bigwedge_{\lambda_3 > \lambda_1} p(y_{\lambda_2}, x_{\lambda_3})$.

Conversely, take any r with $r \in (0, +\infty)$ such that $p(y_{\lambda_2}, x_{\lambda_1}) < r$. Then by (D4) and (B1) we have

$$p(y_{\lambda_2}, x_{\lambda_1}) = p(x_{1-\lambda_1}, y_{1-\lambda_2}) = \bigwedge_{h < 1-\lambda_1} p(x_h, y_{1-\lambda_2}) < r.$$

Therefore, there at least exists h with $h < 1 - \lambda_1$ such that $p(x_h, y_{1-\lambda_2}) < r$, i.e., $p(y_{\lambda_2}, x_{1-h}) < r$. Let $1 - h = \lambda_3$. Then $h < 1 - \lambda_1 \Leftrightarrow \lambda_1 < 1 - h = \lambda_3$ and $p(y_{\lambda_2}, x_{\lambda_3}) < r$. Consequently, $p(y_{\lambda_2}, x_{\lambda_1}) \geq \bigwedge_{\lambda_3 > \lambda_1} p(y_{\lambda_2}, x_{\lambda_3})$, as desired. \square

Theorem 12. If p is a Deng pseudo-metric on I^X , then it is a Chen pseudo-metric.

Proof. We only need to prove that $p(x_\alpha, y_\beta) = \bigvee_{\alpha < \gamma} p(x_\gamma, y_\beta)$. By (D1) and (D2) we have $p(x_\alpha, y_\beta) \geq \bigvee_{\alpha < \gamma} p(x_\gamma, y_\beta)$. If $p(x_\alpha, y_\beta) > \bigvee_{\alpha < \gamma} p(x_\gamma, y_\beta)$, then there exist two numbers s and r such that $p(x_\alpha, y_\beta) > s > r \geq \bigvee_{\alpha < \gamma} p(x_\gamma, y_\beta)$. Therefore, for any $\gamma > \alpha$ we have $s < p(x_\alpha, y_\beta) \leq p(x_\alpha, x_\gamma) + p(x_\gamma, y_\beta) \leq p(x_\alpha, x_\gamma) + r$, and then $s - r < p(x_\alpha, x_\gamma)$. Hence $0 < s - r \leq \bigwedge_{\alpha < \gamma} p(x_\alpha, x_\gamma) = p(x_\alpha, x_\alpha) = 0$. But this is a contradiction, and then it must hold $p(x_\alpha, y_\beta) = \bigvee_{\alpha < \gamma} p(x_\gamma, y_\beta)$, as desired. \square

Theorem 13. If p is a Chen pseudo-metric on I^X and satisfies the property $\bigvee_{s > 1-\lambda_1} p(x_s, y_{\lambda_2}) = \bigvee_{t > 1-\lambda_2} p(y_t, x_{\lambda_1})$, then p is an Erceg pseudo-metric.

Proof. From $\bigvee_{s>1-\lambda_1} p(x_s, y_{\lambda_2}) = \bigvee_{t>1-\lambda_2} p(y_t, x_{\lambda_1})$ and $p(x_\alpha, y_\beta) = \bigvee_{\gamma>\alpha} p(x_\gamma, y_\beta)$, we can obtain $p(x_{1-\lambda_1}, y_{\lambda_2}) = p(y_{1-\lambda_2}, x_{\lambda_1})$, and then

$$\begin{aligned} & \bigvee_{s<\lambda_1} p(y_{\lambda_2}, x_s) = \bigvee_{s<\lambda_1} p(x_{1-s}, y_{1-\lambda_2}) \\ &= \bigvee_{1-s>1-\lambda_1} p(x_{1-s}, y_{1-\lambda_2}) = p(x_{1-\lambda_1}, y_{1-\lambda_2}) = p(y_{\lambda_2}, x_{\lambda_1}). \end{aligned}$$

Consequently, p is an Erceg pseudo-metric, as desired. \square

Conversely, we have the following result:

Theorem 14. *If p is an Erceg pseudo-metric on I^X and satisfies the property $p(x_{1-\lambda_1}, y_{\lambda_2}) = p(y_{1-\lambda_2}, x_{\lambda_1})$, then p is a Chen pseudo-metric.*

Proof. Since $p(x_{1-\lambda_1}, y_{\lambda_2}) = p(y_{1-\lambda_2}, x_{\lambda_1})$, we have the following equation:

$$\begin{aligned} & \bigvee_{t>\lambda_2} p(y_t, x_{\lambda_1}) = \bigvee_{t>\lambda_2} p(x_{1-\lambda_1}, y_{1-t}) \\ &= \bigvee_{1-t<1-\lambda_2} p(x_{1-\lambda_1}, y_{1-t}) = p(x_{1-\lambda_1}, y_{1-\lambda_2}) = p(y_{\lambda_2}, x_{\lambda_1}). \end{aligned}$$

Therefore, p is a Chen pseudo-metric, as desired. \square

In summary, because of Theorem 8 in this section, we have asserted that, if a given $[0, 1]$ -topology δ is Deng-metrizable, then δ must be Erceg-, Shi-, and Chen-metrizable. For this reason, next, we will mainly focus on the Deng metric and its metrization in $[0, 1]$ -topology.

4. The Product of Countable Metric Spaces

In this section, let $Q_{[0,1]}$ be the set of all rational numbers in $[0, 1]$, let $\omega = \{1, 2, \dots, n, \dots\}$, and let $\mathcal{X} = \prod_{n \in \omega} X_n$, where X_n ($n \in \omega$) is a nonempty set. For clarity, the set of all fuzzy points in \mathcal{X} is denoted by $M_0(\mathcal{X})$, and for $y \in \mathcal{X}$, $y = (y^1, y^2, \dots, y^n, \dots)$.

Theorem 15. *Let p be a Deng pseudo-metric on I^X and let $e(x_\alpha, y_\beta) = \min[1, p(x_\alpha, y_\beta)]$. Then (X, δ_e) is a Deng pseudo-metric space whose topology δ_e is identical to that of (X, δ_p) .*

Consequently, each pseudo-metric space (X, δ_p) is homomorphic to a pseudo-metric space (X, δ_e) , where the range of e is the unit interval $[0, 1]$.

Proof. The proof is trivial and omitted. \square

Theorem 16. *Let $\{(X_n, \delta_{p_n}) \mid n \in \omega\}$ be a sequence of Deng pseudo-metric spaces, and the range of p_n ($n \in \omega$) is the unit interval $[0, 1]$. Define a mapping $p : M_0(\mathcal{X}) \times M_0(\mathcal{X}) \rightarrow [0, 1]$ by:*

$$p(x_\alpha, y_\beta) = \sum_{n \in \omega} 2^{-n} p_n(J_n(x_\alpha), J_n(y_\beta)),$$

where $J_n : I^X \rightarrow I^{X_n}$ is the n -th projection (see 2. Preliminaries on J_n). Then

- (1) For each $n \in \omega$, $p_n(J_n(x_\alpha), J_n(y_\beta)) = \bigwedge_{\tau>\beta} p_n(J_n(x_\alpha), J_n(y_\tau))$;
- (2) The mapping p is a Deng pseudo-metric on I^X ;
- (3) The space (\mathcal{X}, δ_p) is the product space of $\{(X_n, \delta_{p_n}) \mid n \in \omega\}$.

Proof. (1) Since $J_n(y_\beta) = y_\beta^n$ ($n \in \omega$), we have

$$p_n(J_n(x_\alpha), J_n(y_\beta)) = p_n(J_n(x_\alpha), y_\beta^n) = \bigwedge_{\tau > \beta} p_n(J_n(x_\alpha), y_\tau^n) = \bigwedge_{\tau > \beta} p_n(J_n(x_\alpha), J_n(y_\tau)).$$

(2) Since p_n ($n \in \omega$) satisfies (D1) and (D2), it is easy to check that p also satisfies (D1) and (D2).

(D3) First, by the definition of p and (1), we have

$$\begin{aligned} p(x_\alpha, y_\beta) &= \sum_{n \in \omega} 2^{-n} p_n(J_n(x_\alpha), J_n(y_\beta)) = \sum_{n \in \omega} 2^{-n} \bigwedge_{\tau > \beta} p_n(J_n(x_\alpha), J_n(y_\tau)) \\ &\leq \bigwedge_{\tau > \beta} \sum_{n \in \omega} 2^{-n} p_n(J_n(x_\alpha), J_n(y_\tau)) \\ &= \bigwedge_{\tau > \beta} p(x_\alpha, y_\tau). \end{aligned}$$

Conversely, let $p(x_\alpha, y_\beta) = r$. Then, for any $\varepsilon > 0$ we have

$$p(x_\alpha, y_\beta) = \sum_{n \in \omega} 2^{-n} p_n(J_n(x_\alpha), J_n(y_\beta)) = \sum_{n \in \omega} 2^{-n} \bigwedge_{\tau > \beta} p_n(J_n(x_\alpha), J_n(y_\tau)) < r + \varepsilon.$$

Let $r_n = 2^{-n} \bigwedge_{\tau > \beta} p_n(J_n(x_\alpha), J_n(y_\tau))$. Then $\bigwedge_{\tau > \beta} p_n(J_n(x_\alpha), J_n(y_\tau)) < 2^n \times r_n + \varepsilon$. Therefore, for each n there is a number τ_n with $\tau_n > \beta$ such that $p_n(J_n(x_\alpha), J_n(y_{\tau_n})) < 2^n \times r_n + \varepsilon$. Hence we have

$$\sum_{n \in \omega} 2^{-n} p_n(J_n(x_\alpha), J_n(y_{\tau_n})) < \sum_{n \in \omega} r_n + \varepsilon \sum_{n \in \omega} 2^{-n} = r + \varepsilon.$$

Given every fixed natural number n , we can take a number μ_n with $\mu_n > \beta$ such that $\mu_n \leq \min\{\tau_1, \tau_2, \dots, \tau_n\}$. Thus for any natural number m , we have

$$\begin{aligned} \sum_{i=1}^m 2^{-i} p_i(J_i(x_\alpha), J_i(y_{\mu_m})) + \sum_{i=m+1}^\infty 2^{-i} p_i(J_i(x_\alpha), J_i(y_{\tau_i})) \\ \leq \sum_{i=1}^\infty 2^{-i} p_i(J_i(x_\alpha), J_i(y_{\tau_i})) < r + \varepsilon, \end{aligned}$$

which is equivalent to the following inequality:

$$\sum_{i=1}^\infty 2^{-i} p_i(J_i(x_\alpha), J_i(y_{\mu_m})) + \sum_{i=m+1}^\infty 2^{-i} [p_i(J_i(x_\alpha), J_i(y_{\tau_i})) - p_i(J_i(x_\alpha), J_i(y_{\mu_m}))] < r + \varepsilon.$$

Consequently, for the fixed natural member m , we have

$$\begin{aligned} \sum_{i=1}^\infty 2^{-i} p_i(J_i(x_\alpha), J_i(y_{\mu_m})) \\ < r + \varepsilon - \sum_{i=m+1}^\infty 2^{-i} [p_i(J_i(x_\alpha), J_i(y_{\tau_i})) - p_i(J_i(x_\alpha), J_i(y_{\mu_m}))] \\ = r + \varepsilon + \sum_{i=m+1}^\infty 2^{-i} [p_i(J_i(x_\alpha), J_i(y_{\mu_m})) - p_i(J_i(x_\alpha), J_i(y_{\tau_i}))] \\ \leq r + \varepsilon + \sum_{i=m+1}^\infty 2^{-i} p_i(J_i(y_{\tau_i}), J_i(y_{\mu_m})) \\ \leq r + \varepsilon + \sum_{i=m+1}^\infty 2^{-i}. \end{aligned}$$

Hence

$$\bigwedge_{t > \beta} p(x_\alpha, y_t) \leq \bigwedge_{\mu_m > \beta} p(x_\alpha, y_{\mu_m}) = \bigwedge_{\mu_m > \beta} \sum_{i=1}^\infty 2^{-i} p_i(J_i(x_\alpha), J_i(y_{\mu_m})) \leq r + \varepsilon.$$

Because ε is arbitrary, we can obtain $\bigwedge_{\tau > \beta} p(x_\alpha, y_\tau) \leq r = p(x_\alpha, y_\beta)$. Therefore, p satisfies (D3).

(D4) Since it holds that $p_n(J_n(x_\alpha), J_n(y_\beta)) = p_n(J_n(y_{1-\beta}), J_n(x_{1-\alpha}))$ for each $n \in \omega$, we conclude that $p(x_\alpha, y_\beta) = p(y_{1-\beta}, x_{1-\alpha})$.

(3) First, let (\mathcal{X}, δ) be the product space of $\{(X_n, \delta_{p_n}), n \in \omega\}$. For any $V \in \delta_p$ and $x_\alpha \in V$, there is an open neighborhood $U_r(x_\alpha)$ of x_α such that $U_r(x_\alpha) \leq V$, where

$$U_r(x_\alpha) = \bigvee \{y_\beta \mid p(x_\alpha, y_\beta) = \sum_{i=1}^{\infty} 2^{-i} p_n(J_n(x_\alpha), J_n(y_\beta)) < r\}.$$

Taking a natural member q with $\frac{1}{2^q} < r$, we can get $U_{\frac{1}{2^q}}(x_\alpha) \leq V$. Thus, if we define

$$W = \bigvee \{z_\gamma \mid p_n(J_n(x_\alpha), J_n(z_\gamma)) < \frac{1}{2^{q+n+2}}, n \leq q + 2\},$$

then $W \leq U_{\frac{1}{2^q}}(x_\alpha)$. This is because when $z_\gamma \in W$, we always have

$$p(x_\alpha, z_\gamma) < \sum_{n=1}^{n=q+2} \frac{1}{2^{q+n+2}} + \sum_{n=q+3}^{\infty} \frac{1}{2^n} < \frac{1}{2^{q+2}} + \frac{1}{2^{q+2}} = \frac{1}{2^{q+1}} < \frac{1}{2^q}.$$

Clearly, W is an open set in the product topology δ since it can be generated by the subbase of the product space (\mathcal{X}, δ) . Therefore, we conclude that V is an open set in (\mathcal{X}, δ) . Consequently, $\delta_p \subseteq \delta$.

Conversely, let $U = \bigvee \{x_\alpha \mid x_\alpha^n \in V\}$, where V is an open set of some δ_{p_n} . Then U is a member of the subbase of δ . If $x_\alpha \in \{x_\alpha \mid x_\alpha^n \in V\}$, then there is an open set $U_r(x_\alpha^n)$ (see Theorem 1 on U_r) belonging to δ_{p_n} such that $U_r(x_\alpha^n) \leq V$. Therefore, $J_n^{-1}(U_r(x_\alpha^n)) \leq U$. Since $p(x_\alpha, y_\beta) \geq 2^{-n} p_n(x_\alpha^n, y_\beta^n)$, the open set $U_{\frac{r}{2^n}}(x_\alpha)$ of δ_p is a subset of U . In fact, if $p(x_\alpha, y_\beta) < \frac{r}{2^n}$, then $p_n(x_\alpha^n, y_\beta^n) < r$. Hence $U_{\frac{r}{2^n}}(x_\alpha) \leq J_n^{-1}(U_r(x_\alpha^n))$, and then $U_{\frac{r}{2^n}}(x_\alpha) \leq U$. Therefore, U is the union of some open sets in δ_p . Consequently, $\delta \subseteq \delta_p$. To sum up, the proposition is proved. \square

5. σ -Locally Finite Property

In this section, some σ -locally finite properties of Deng pseudo-metric space will be examined based on a defined distance function between fuzzy sets.

Definition 2. Let p be a Deng pseudo-metric on I^X . A distance function $d : I^X \times I^X \rightarrow [0, +\infty)$ is defined by:

$$d[A, B] = \inf\{p(x_\alpha, y_\beta) \mid x_\alpha \in A, y_{1-\beta} \in B\}.$$

Let $A, B \in I^X$, and $x_\alpha \in M_0$. Then, by definition, it is easy to prove that $d[A, x_{1-\alpha}] = \inf\{p(y_\beta, x_\alpha) \mid y_\beta \in A\}$, $d[x_\alpha, B] = \inf\{p(x_\alpha, y_\beta) \mid y_{1-\beta} \in B\}$ and $d[A, B] = d[B, A]$.

Theorem 17. Let p be a Deng pseudo-metric on I^X . If fuzzy sets U and V are quasi-coincident, then $d[U, V] = d[V, U] = 0$.

Proof. Because U and V are quasi-coincident, there is x belonging to X such that $U(x) + V(x) > 1$. Let $U(x) = \gamma$ and $V(x) = \beta$. Given α with $1 - \beta < \alpha < \gamma$, we have $x_\alpha \in U$. Take a number λ satisfying $\alpha > \lambda > 1 - \beta$. Since $x_{1-\lambda} \in V$, $d[U, V] \leq p(x_\alpha, x_\lambda) = 0$. Similarly, $d[V, U] = 0$. \square

Theorem 18. Let p be a Deng pseudo-metric on I^X . Suppose that \mathbb{U} is an open cover of (X, δ_p) , and for each U of \mathbb{U} and each positive integer n , define $U_n = \bigvee \{x_\alpha \mid d[x_\alpha, U'] \geq \frac{1}{2^n}\}$. Then $d[U_n, U'_{n+1}] \geq \frac{1}{2^{n+1}}$.

Proof. Let $x_\alpha \in U_n$ and $y_{1-\beta} \in U'_{n+1}$. For any $z_{1-\gamma} \in U'$, we can obtain $p(x_\alpha, y_\beta) + p(y_\beta, z_\gamma) \geq p(x_\alpha, z_\gamma) \geq d[x_\alpha, U']$. Hence, it is clear that

$$p(x_\alpha, y_\beta) + d[y_\beta, U'] \geq d[x_\alpha, U'], \text{ i.e., } p(x_\alpha, y_\beta) \geq d[x_\alpha, U'] - d[y_\beta, U'].$$

Because of $x_\alpha \in U_n$, there is x_η belonging to U such that $\eta > \alpha$ and $d[x_\alpha, U'] \geq d[x_\eta, U'] \geq \frac{1}{2^\eta}$. Because of $y_{1-\beta} \in U'_{n+1}$, we assert that $d[y_\beta, U'] < \frac{1}{2^{n+1}}$. Thus $p(x_\alpha, y_\beta) \geq d[x_\alpha, U'] - d[y_\beta, U'] > \frac{1}{2^n} - \frac{1}{2^{n+1}} = \frac{1}{2^{n+1}}$. Consequently, $d[U_n, U'_{n+1}] \geq \frac{1}{2^{n+1}}$. \square

Since \mathbb{U} is a nonempty set, we can select a partial order on \mathbb{U} such that \mathbb{U} is well ordered, denoted by “ \prec ” (see Theorem 25 in Chapter 0 of [56]: Every nonempty set can be well ordered).

Theorem 19. Let p be a Deng pseudo-metric on I^X and let the family \mathbb{U} be an open cover of (X, δ_p) . Choose a relation \prec which well orders the family \mathbb{U} and for each $U \in \mathbb{U}$ and each $n \in \omega$ define $U_n^* = U_n \wedge (\vee \{V_{n+1} \mid V \in \mathbb{U} \text{ and } V \prec U\})'$. Then

- (1) Either $V_n^* \leq U'_{n+1}$ or $U_n^* \leq V'_{n+1}$ is true, depending on whether U follows or precedes V in the ordering;
- (2) In either case, $d[U_n^*, V_n^*] \geq \frac{1}{2^{n+1}}$.

Proof. (1) The proof is straightforward.

(2) It is easy to see that $V_n^* \leq V_n$. Furthermore, if $V \prec U$, then $U_n^* \leq V'_{n+1}$. Hence $d[U_n^*, V_n^*] \geq d[V'_{n+1}, V_n] = d[V_n, V'_{n+1}] \geq \frac{1}{2^{n+1}}$. Similarly, when $U \prec V$, $d[U_n^*, V_n^*] \geq \frac{1}{2^{n+1}}$. \square

Theorem 20. Let p be a Deng pseudo-metric on I^X and let the family \mathbb{U} be an open cover of (X, δ_p) . Given $U \in \mathbb{U}$ and given any $n \in \omega$, for each corresponding U_n^* , define $U_n^\sim = \vee \{x_\alpha \mid d[U_n^*, x_{1-\alpha}] < \frac{1}{2^{n+3}}\}$. Then

- (1) U_n^\sim is an open set;
- (2) $d[U_n^\sim, V_n^\sim] \geq \frac{1}{2^{n+2}}$.

Proof. (1) Take a fuzzy point x_α and a number s_α such that they satisfy $d[U_n^*, x_{1-\alpha}] = r < \frac{1}{2^{n+3}}$ and $0 < s_\alpha < \frac{1}{2^{n+3}} - r$, respectively. It follows that $d[U_n^*, z_{1-\gamma}] \leq d[U_n^*, x_{1-\alpha}] + p(x_\alpha, z_\gamma) < r + s_\alpha < \frac{1}{2^{n+3}}$ for any $z_\gamma \in U_{s_\alpha}(x_\alpha)$. Therefore, $U_{s_\alpha}(x_\alpha) \leq U_n^\sim$, so that U_n^\sim is an open set. In addition, if $d[U_n^*, x_{1-\alpha}] < \frac{1}{2^{n+3}} < \frac{1}{2^{n+1}}$, then there is $y_\mu \in U_n^*$ such that $p(y_\mu, x_\alpha) < \frac{1}{2^{n+1}}$. But according to $d[U_n^*, U'_{n+1}] \geq d[U_n, U'_{n+1}] \geq \frac{1}{2^{n+1}}$, it must hold that $x_\alpha \leq U_{n+1}(x)$. Thus $U_n^\sim \leq U_{n+1} \leq U$.

(2) Taking $x_\alpha \in U_n^\sim$ and $y_\beta \in V_n^\sim$ such that $d[U_n^*, x_{1-\alpha}] < \frac{1}{2^{n+3}}$ and $d[V_n^*, y_{1-\beta}] < \frac{1}{2^{n+3}}$, we have $p(z_\gamma, x_\alpha) + p(x_\alpha, y_{1-\beta}) \geq p(z_\gamma, y_{1-\beta}) \geq d[U_n^*, y_\beta]$ for any $z_\gamma \in U_n^*$. Therefore, we have $d[U_n^*, x_\alpha] + p(x_\alpha, y_{1-\beta}) \geq d[U_n^*, y_\beta]$, so that

$$d[U_n^*, x_\alpha] + p(x_\alpha, y_{1-\beta}) + d[y_{1-\beta}, V_n^*] \geq d[U_n^*, y_\beta] + d[y_{1-\beta}, V_n^*].$$

Since for any $z_\gamma \in U_n^*$, $p(z_\gamma, y_{1-\beta}) + p(y_{1-\beta}, w_\alpha) \geq p(z_\gamma, w_\alpha) \geq d[z_\gamma, V_n^*]$ and $w_{1-\alpha} \in V_n^*$, we have $p(z_\gamma, y_{1-\beta}) + d[y_{1-\beta}, V_n^*] \geq d[z_\gamma, V_n^*]$. In addition, in view of $d[z_\gamma, V_n^*] \geq d[U_n^*, V_n^*]$, we can check that $p(z_\gamma, y_{1-\beta}) + d[y_{1-\beta}, V_n^*] \geq d[U_n^*, V_n^*]$. Therefore, we assert that $d[U_n^*, y_\beta] + d[y_{1-\beta}, V_n^*] \geq d[U_n^*, V_n^*]$, so that

$$d[U_n^*, x_{1-\alpha}] + p(x_\alpha, y_{1-\beta}) + d[y_{1-\beta}, V_n^*] \geq d[U_n^*, V_n^*].$$

Additionally, from $x_\alpha \in U_n^\sim$ and $d[U_n^*, V_n^*] \geq \frac{1}{2^{n+1}}$, we have the following inequalities:

$$p(x_\alpha, y_{1-\beta}) \geq d[U_n^*, V_n^*] - d[U_n^*, x_{1-\alpha}] - d[y_{1-\beta}, V_n^*] > \frac{1}{2^{n+1}} - \frac{1}{2^{n+3}} - d[y_{1-\beta}, V_n^*].$$

Note that $y_\beta \in V_n^\sim$ and $d[y_{1-\beta}, V_n^*] = \bigwedge_{x_\omega \in V_n^*} p(y_{1-\beta}, x_{1-\omega}) = \bigwedge_{x_\omega \in V_n^*} p(x_\omega, y_\beta) = d[V_n^*, y_{1-\beta}] \leq \frac{1}{2^{n+3}}$. Therefore, $p(x_\alpha, y_{1-\beta}) > \frac{1}{2^{n+1}} - \frac{1}{2^{n+3}} - \frac{1}{2^{n+3}} = \frac{1}{2^{n+2}}$, so that $d[U_n^\sim, V_n^\sim] \geq \frac{1}{2^{n+2}}$, as desired. \square

Theorem 21. Let \mathbb{V}_n be the family of all sets of the form $U_n^\sim (n \in \omega)$. Given fuzzy point x_α . Then

- (1) If there is a fixed number $r > 0$ such that $d[x_\alpha, V_n^\sim] > r$ for each $V_n^\sim \in \mathbb{V}_n$, then $x_{1-\alpha} \not\leq \bigvee_{U_n^\sim \in \mathbb{V}_n} U_n^\sim$;
- (2) If such a fixed number $r > 0$ is non-existent, then $x_{1-\alpha} \leq \overline{\bigvee_{U_n^\sim \in \mathbb{V}_n} U_n^\sim}$;
- (3) If $\alpha \leq \frac{1}{2}$, then there at most exists a $U_n^\sim \in \mathbb{V}_n$ such that $x_{1-\alpha} \leq \overline{U_n^\sim}$.

Proof. (1) Given $U_n^\sim \in \mathbb{V}_n$, we have $p(x_\alpha, y_{1-\beta}) > r$ for all $y_\beta \in U_n^\sim$. By Theorem 7, $y_{1-\beta} \not\leq B_r(x_\alpha)$, so that $y_{1-\beta} \not\leq U_r(x_\alpha)$, i.e., $U_r(x_\alpha)(y) + \beta < 1$. Therefore, $U_r(x_\alpha)(y) + U_n^\sim(y) \leq 1$ for all $y \in X$. Hence for each $V_n^\sim \in \mathbb{V}_n$, $U_r(x_\alpha)$ and V_n^\sim are non-quasi-coincident, as desired.

(2) Because such a fixed number r is non-existent, for any $\epsilon_k > 0$, there is $V_{n(\epsilon_k)}^\sim$ such that $d[x_\alpha, V_{n(\epsilon_k)}^\sim] < \epsilon_k$. This means that each open neighborhood of x_α is quasi-coincident with $\bigvee_{U_n^\sim \in \mathbb{V}_n} U_n^\sim$. Therefore, $x_{1-\alpha} \leq \overline{\bigvee_{U_n^\sim \in \mathbb{V}_n} U_n^\sim}$, as desired.

(3) Because $\alpha \leq \frac{1}{2}$, we conclude that $p(x_{1-\alpha}, x_\alpha) = 0$. Assume that there exist two $U_n^\sim, V_n^\sim \in \mathbb{V}_n$ such that $x_{1-\alpha} \leq \overline{U_n^\sim}$ and $x_{1-\alpha} \leq \overline{V_n^\sim}$. Then by $d[x_\alpha, U_n^\sim] = 0$ and $d[x_\alpha, V_n^\sim] = 0$ we have

$$d[V_n^\sim, x_\alpha] + p(x_{1-\alpha}, x_\alpha) + d[x_\alpha, U_n^\sim] \geq d[V_n^\sim, U_n^\sim] \geq \frac{1}{2^{n+2}}.$$

Note that $d[V_n^\sim, x_\alpha] = d[x_\alpha, V_n^\sim]$. Therefore, we can obtain that $0 \geq \frac{1}{2^{n+2}}$. But this is a contradiction. \square

6. Two Interrelated Mappings

To solve the metrization problem in $[0, 1]$ -topology in the next section, we shall construct two interrelated mappings in advance based on the normal spaces in this section.

Theorem 22. Let (X, δ) be normal $[0, 1]$ -topological space and let $A \in \delta', B \in \delta$ with $A \leq B$. Therefore, there exists a corresponding family $\{U_r \mid r \in Q_{[0,1]}\}$ relative to A and B satisfying (a) and (b) in Theorem 5.

Define a mapping $f : M_0 \rightarrow [0, 1]$ by

$$f(x_\alpha) = \begin{cases} \inf\{r \in Q_{[0,1]} \mid x_\alpha \in U_r\}, & x_\alpha \in B; \\ 1, & x_\alpha \notin B \end{cases}$$

and for all $x_\alpha, y_\beta \in M_0$ let $g(x_\alpha, y_\beta) = \max\{f(y_\beta) - f(x_\alpha), 0\}$.

(i) Let $V_r = U'_{1-r}$. Then, the family $\{V_r \mid r \in Q_{[0,1]}\}$ satisfies the following properties:

(1) $V_0 = B', V_1 = A'$; (2) if $r < s$, then $V_r \leq V_s$.

(ii) Define a mapping $f^* : M_0 \rightarrow [0, 1]$ by

$$f^*(x_\alpha) = \begin{cases} \inf\{r \in Q_{[0,1]} \mid x_\alpha \leq V_r\}, & x_\alpha \leq A'; \\ 1, & x_\alpha \not\leq A' \end{cases}$$

and for all $x_\alpha, y_\beta \in M_0$ let $g^*(x_\alpha, y_\beta) = \max\{f^*(y_\beta) - f^*(x_\alpha), 0\}$. Then $g^*(y_{1-\beta}, x_{1-\alpha}) = g(x_\alpha, y_\beta)$.

(iii) Both g and g^* satisfy the properties (D1)–(D3) in Definition 1.

Proof. (i) The proof is straightforward.

(ii) Case 1. Assume that $x_\alpha \in B$. Then let us consider two subcases below.

Subcase 1. If $x_\alpha \in A$, then $f(x_\alpha) = \inf\{r \in Q_{[0,1]} \mid x_\alpha \in U_r\} = 0$.

(1) Assume that $y_\beta \in B$. (i) For the case of $y_\beta \in A$, we have $f(y_\beta) = 0$, and then $g(x_\alpha, y_\beta) = 0$. On the other hand, according to $x_\alpha \in A \Leftrightarrow x_{1-\alpha} \notin A'$ and $y_\beta \in A \Leftrightarrow y_{1-\beta} \notin A'$, we have $f^*(x_{1-\alpha}) = 1$ and $f^*(y_{1-\beta}) = 1$. Hence $g^*(y_{1-\beta}, x_{1-\alpha}) = 0$; (ii) For the case of $y_\beta \notin A$, let $f(y_\beta) = \inf\{r \in Q_{[0,1]} \mid y_\beta \in U_r\} = s$. If $s < 1$, then $g(x_\alpha, y_\beta) = s$ and there exists a monotonically decreasing sequence $S = \{s_n \mid s_n \geq s, y_\beta \in U_{s_n}, n \in \omega\} \subseteq Q_{[0,1]}$ such that $\lim_{n \rightarrow \infty} s_n = s$. Therefore, for each $\varepsilon \in Q_{[0,1]}$ with $\varepsilon > 0$ there is a natural number $N(\varepsilon)$ such that $y_{1-\beta} \in U'_{s_n-\varepsilon} = V_{1-s_n+\varepsilon}$ whenever $n > N(\varepsilon)$. Therefore,

$$f^*(y_{1-\beta}) = \inf\{r \in Q_{[0,1]} \mid y_{1-\beta} \leq V_r\} \leq \inf\{1 - s_n + \varepsilon \mid y_{1-\beta} \in V_{1-s_n+\varepsilon}\} \leq 1 - s + \varepsilon,$$

so that $f^*(y_{1-\beta}) \leq 1 - s$ by the arbitrariness of ε . In addition, if $y_{1-\beta} \leq V_r$, then from the equivalence $y_\beta \in U_{s_n} \Leftrightarrow 1 - \beta > V_{1-s_n}(y)$ we have $V_r \geq V_{1-s_n}$, i.e., $r \geq 1 - s_n$ for all $s_n \in S$. Thus $f^*(y_{1-\beta}) = \inf\{r \in Q_{[0,1]} \mid y_{1-\beta} \leq V_r\} \geq 1 - s$. Consequently $f^*(y_{1-\beta}) = 1 - s$, and then $g^*(y_{1-\beta}, x_{1-\alpha}) = s$. If $s = 1$, i.e., $f(y_\beta) = 1$, then $g(x_\alpha, y_\beta) = 1$. In addition, by $f(y_\beta) = 1$, we assert that $y_\beta \in U_1$, but $y_\beta \notin U_r$ for all other $r \in Q_{[0,1]}$. Thus when $r \neq 0$, $y_{1-\beta} \leq V_r$, and then $f^*(y_{1-\beta}) = 0$. Consequently $g^*(y_{1-\beta}, x_{1-\alpha}) = 1$.

(2) If $y_\beta \notin B$, then $f(y_\beta) = 1$, and thus $g(x_\alpha, y_\beta) = 1$. According to $y_\beta \notin B \Leftrightarrow y_{1-\beta} \leq B'$, we assert that $y_{1-\beta} \leq V_r$ for all $r \in Q_{[0,1]}$, so that $f^*(y_{1-\beta}) = 0$. Consequently, $g^*(y_{1-\beta}, x_{1-\alpha}) = 1$.

Subcase 2. Let $x_\alpha \notin A$ and let $f(x_\alpha) = t$.

(1) If $y_\beta \in B$, then (i) For the case of $y_\beta \in A$, we have $f(y_\beta) = 0$. So $g(x_\alpha, y_\beta) = 0$. Moreover, from $y_\beta \in A \Leftrightarrow y_{1-\beta} \notin A'$, we know $f^*(y_{1-\beta}) = 1$, and then $g^*(y_{1-\beta}, x_{1-\alpha}) = 0$; (ii) For the case of $y_\beta \notin A$, let $f(y_\beta) = s$. Then $g(x_\alpha, y_\beta) = \max\{s - t, 0\}$. Similarly, it is true that $f^*(x_{1-\alpha}) = 1 - t$ and $f^*(y_{1-\beta}) = 1 - s$. Thus, $g^*(y_{1-\beta}, x_{1-\alpha}) = \max\{1 - t - (1 - s), 0\} = \max\{s - t, 0\}$.

(2) If $y_\beta \notin B$, then $f(y_\beta) = 1$ and $g(x_\alpha, y_\beta) = \max\{1 - t, 0\} = 1 - t$. In addition, by $f^*(y_{1-\beta}) = 0$ and $f^*(x_{1-\alpha}) = 1 - t$, we have $g^*(y_{1-\beta}, x_{1-\alpha}) = 1 - t$.

Case 2. Assume that $x_\alpha \notin B$. Then $f(x_\alpha) = 1$ and $f^*(x_{1-\alpha}) = 0$.

(1) Let $y_\beta \in B$. (i) If $y_\beta \in A$, then $f(y_\beta) = 0$, and thus $g(x_\alpha, y_\beta) = 0$. From $f^*(x_{1-\alpha}) = 0$ we have $g^*(y_{1-\beta}, x_{1-\alpha}) = 0$. (ii) If $y_\beta \notin A$, then from $f(x_\alpha) = 1$ we know $g(x_\alpha, y_\beta) = 0$. Meanwhile, because of $f^*(x_{1-\alpha}) = 0$, we can obtain $g^*(y_{1-\beta}, x_{1-\alpha}) = 0$.

(2) Let $y_\beta \notin B$. Then, $f(y_\beta) = 1$. Note that $f(x_\alpha) = 1$. So $g(x_\alpha, y_\beta) = 0$. Owing to $f^*(x_{1-\alpha}) = 0$, we know $g^*(y_{1-\beta}, x_{1-\alpha}) = 0$.

(iii) (D1) Let $x_\alpha \geq x_\beta$. If $x_\beta \notin B$, then $f(x_\alpha) = f(x_\beta) = 1$. So $g(x_\alpha, x_\beta) = 0$. If $x_\beta \in B$, then when $x_\alpha \notin B$, we have $f(x_\alpha) = 1$, and then $g(x_\alpha, x_\beta) = 0$; when $x_\alpha \in B$, we have $f(x_\alpha) = \inf\{r \in Q_{[0,1]} \mid x_\alpha \in U_r\} \geq \inf\{r \in Q_{[0,1]} \mid x_\beta \in U_r\} = f(x_\beta)$, and then $g(x_\alpha, x_\beta) = 0$. Besides, by $g^*(y_{1-\beta}, x_{1-\alpha}) = g(x_\alpha, y_\beta)$, it is easy to show that g^* also satisfies (D1).

(D2) To check $g(x_\alpha, y_\beta) + g(y_\beta, z_\gamma) \geq g(x_\alpha, z_\gamma)$ for any $x_\alpha, y_\beta, z_\gamma \in M_0$, we consider the following two cases: (a) when $g(x_\alpha, z_\gamma) = 0$, this conclusion is straightforward; (b) when $g(x_\alpha, z_\gamma) \neq 0$, we have $f(z_\gamma) > f(x_\alpha)$. In this case, (1) if $g(y_\beta, z_\gamma) = 0$, then $f(z_\gamma) \leq f(y_\beta)$, and thus $g(x_\alpha, y_\beta) = f(y_\beta) - f(x_\alpha) \geq f(z_\gamma) - f(x_\alpha) = g(x_\alpha, z_\gamma)$; (2) if $g(x_\alpha, y_\beta) = 0$, then $f(y_\beta) \leq f(x_\alpha)$, and then $f(z_\gamma) - f(y_\beta) \geq f(z_\gamma) - f(x_\alpha)$. Therefore, $g(y_\beta, z_\gamma) \geq g(x_\alpha, z_\gamma)$; (3) if $g(y_\beta, z_\gamma) \neq 0$ and $g(x_\alpha, y_\beta) \neq 0$, then $f(z_\gamma) > f(y_\beta)$ and $f(y_\beta) > f(x_\alpha)$, and then $g(x_\alpha, y_\beta) + g(y_\beta, z_\gamma) = [f(y_\beta) - f(x_\alpha)] + [f(z_\gamma) - f(y_\beta)] = f(z_\gamma) - f(x_\alpha) = g(x_\alpha, z_\gamma)$. Besides, by $g^*(y_{1-\beta}, x_{1-\alpha}) = g(x_\alpha, y_\beta)$, it is easy to see that g^* also satisfies (D2).

(D3) (1) Assume that $f(y_\beta) = 1$. (a) If $y_\beta \in B$, then besides $r = 1, y_\beta \notin U_r$ for all other $r \in Q_{[0,1]}$. Thus, for each η with $\beta < \eta < B(y)$ we assert that besides $r = 1, y_\eta \notin U_r$ for all other $r \in Q_{[0,1]}$. Consequently $f(y_\eta) = 1$. It follows that $g(x_\alpha, y_\eta) = \max\{1 - f(x_\alpha), 0\} = g(x_\alpha, y_\beta)$; (b) if $y_\beta \notin B$, then for each γ with $\beta < \gamma < 1$ we have $y_\gamma \notin B$, and then $f(y_\gamma) = 1$. Therefore, $g(x_\alpha, y_\gamma) = \max\{1 - f(x_\alpha), 0\} = g(x_\alpha, y_\beta)$.

(2) Assume that $0 \leq f(y_\beta) = p < 1$. (a) If there is a fixed number $h > 0$ such that $y_{\beta+h} \leq U_q$ for each q with $q \in (p, 1] \cap Q_{[0,1]}$, and then $f(y_\gamma) = p$ for each γ with $\beta < \gamma < \beta + h$. Therefore, $g(x_\alpha, y_\gamma) = \max\{p - f(x_\alpha), 0\} = g(x_\alpha, y_\beta)$. (b) If such a fixed h is non-existent, then by $f(y_\beta) = \inf\{r \in Q_{[0,1]} \mid y_\beta \in U_r\} = p$ we assert that for any $\varepsilon > 0$ there exists a r belonging to $Q_{[0,1]}$ such that $y_\beta \in U_r$ and $r - p < \varepsilon$. Take a number γ satisfying $\beta < \gamma < \beta + \frac{r-p}{2}$. Then $f(y_\gamma) < p + \varepsilon$. Let $\bigwedge_{\gamma>\beta} f(y_\gamma) = q$. Clearly, $p \leq q$. Thus for any $\varepsilon > 0$

we have $q = \bigwedge_{\gamma>\beta} f(y_\gamma) < p + \varepsilon$. Because ε is arbitrary, we can obtain $q \leq p$. Thus $f(y_\beta) =$

$\bigwedge_{\gamma>\beta} f(y_\gamma)$. Consequently, $g(x_\alpha, y_\beta) = \max\{p - f(x_\alpha), 0\} = \max\{\bigwedge_{\gamma>\beta} f(y_\gamma) - f(x_\alpha), 0\}$.

(1) If $p - f(x_\alpha) < 0$, then $g(x_\alpha, y_\beta) = 0$ and there exists a γ satisfying $\gamma > \beta$ such that $f(y_\gamma) - f(x_\alpha) < 0$. Meanwhile, $\bigwedge_{\gamma>\beta} g(x_\alpha, y_\gamma) = \bigwedge_{\gamma>\beta} \{\max\{f(y_\gamma) - f(x_\alpha), 0\}\} = 0$. Hence

$g(x_\alpha, y_\beta) = \bigwedge_{\gamma>\beta} g(x_\alpha, y_\gamma)$. (2) If $p - f(x_\alpha) = 0$, then this means that for any $\varepsilon > 0$ there

exists a γ satisfying $\gamma > \beta$ such that $f(y_\gamma) - f(x_\alpha) < \varepsilon$. Therefore, $\bigwedge_{\gamma>\beta} g(x_\alpha, y_\gamma) =$

$\bigwedge_{\gamma>\beta} \{\max\{f(y_\gamma) - f(x_\alpha), 0\}\} < \varepsilon$, so that $\bigwedge_{\gamma>\beta} g(x_\alpha, y_\gamma) = 0$. (3) If $p - f(x_\alpha) > 0$, then

$f(y_\gamma) - f(x_\alpha) > 0$ for each $\gamma > \beta$. It follows that $\bigwedge_{\gamma>\beta} g(x_\alpha, y_\gamma) = \bigwedge_{\gamma>\beta} \{\max\{f(y_\gamma) -$

$f(x_\alpha), 0\}\} = \bigwedge_{\gamma>\beta} \{f(y_\gamma) - f(x_\alpha)\} = p - f(x_\alpha)$. In summary, g satisfies (D3). Similarly, so

does g^* . \square

7. Metrization Theorem

For a $[0, 1]$ -topological space (X, δ) , if there is a Deng pseudo-metric (resp., Deng metric) p on I^X such that $\delta = \delta_p$, where δ_p is the pseudo-metric topology, then the space is said to be *Deng pseudo-metrizable* (resp., *Deng-metrizable*). Similar treatment is given to the Erceg metric, Chen metric, and Shi metric.

Theorem 23. *If a $[0, 1]$ -topological space is regular, and the topology has a σ -locally finite base, then it is Deng pseudo-metrizable.*

Proof. The sketch of proof is: Firstly, by Theorem 22 and the property of σ -locally finite base, we will generate a countable family of Deng pseudo-metric spaces $\{(X_n, \delta_{p_n}) \mid n \in \omega\}$. Secondly, by Theorem 16, we will construct a Deng pseudo-metric on I^X and prove that the space (X, δ_p) is exactly the product space of the family $\{(X_n, \delta_{p_n}) \mid n \in \omega\}$. Finally, we will deduce that (X, δ) can be embedded into (X, δ_p) .

Now let us complete the proof step by step. First, for each $n \in \omega$ let $\sigma^n = \{A_i^n \mid i \in \Gamma_n\}$ be locally finite in (X, δ) and let the union $\sigma = \bigcup\{\sigma^n \mid n \in \omega\}$ be a base of (X, δ) .

Arbitrarily select a pair of positive integers m, n . Let $A_i^m \in \sigma^m$ and let it be fixed for the moment. We consider the following open set:

$$A_i^{m,n} = \bigvee\{A^m \mid A^m \in \sigma^m, \overline{A^m} \leq A_i^n\}.$$

For convenience, we denote $A_i^{m,n}$ as A_i . Because δ_m is locally finite, by Theorem 4 we have $\overline{A_i} \leq A_i^n$. Next, the proof shall be divided into the following five steps.

Step 1. By Theorem 22 there exists a family $\{U_r \mid r \in Q_{[0,1]}\}$ corresponding to $\overline{A_i}$ and A_i^n . Therefore, we can define a mapping $f_i : M_0 \rightarrow [0, 1]$ by

$$f_i(x_\alpha) = \begin{cases} \inf\{r \in Q_{[0,1]} \mid x_\alpha \in U_r\}, & x_\alpha \in A_i^n; \\ 1, & x_\alpha \notin A_i^n. \end{cases}$$

For any $x_\alpha, y_\beta \in M_0$ let $g_i(x_\alpha, y_\beta) = \max\{f_i(y_\beta) - f_i(x_\alpha), 0\}$ and for each $r \in Q_{[0,1]}$ let $V_r = U'_{1-r}$. Then there exists a family $\{V_r \mid r \in Q_{[0,1]}\}$. Therefore, we can define a mapping f_i^* by

$$f_i^*(x_\alpha) = \begin{cases} \inf\{r \in Q_{[0,1]} \mid x_\alpha \leq V_r\}, & x_\alpha \leq (\overline{A_i})'; \\ 1, & x_\alpha \not\leq (\overline{A_i})'. \end{cases}$$

Let $g_i^*(x_\alpha, y_\beta) = \max\{f_i^*(y_\beta) - f_i^*(x_\alpha), 0\}$. Then by Theorem 22, we have $g^*(y_{1-\beta}, x_{1-\alpha}) = g(x_\alpha, y_\beta)$.

Step 2. Let $p_i(x_\alpha, y_\beta) = [g_i(x_\alpha, y_\beta) + g_i^*(x_\alpha, y_\beta)]/2$. Then p_i is a Deng pseudo-metric on I^X . This is because both g_i and g_i^* satisfy the properties (D1)–(D3), and so does p_i . Besides, by Theorem 22, we have the following two equalities:

$$p_i(x_\alpha, y_\beta) = \frac{g_i(x_\alpha, y_\beta) + g_i^*(x_\alpha, y_\beta)}{2} = \frac{g_i(x_\alpha, y_\beta) + g_i(y_{1-\beta}, x_{1-\alpha})}{2};$$

$$p_i(y_{1-\beta}, x_{1-\alpha}) = \frac{g_i(y_{1-\beta}, x_{1-\alpha}) + g_i^*(y_{1-\beta}, x_{1-\alpha})}{2} = \frac{g_i(y_{1-\beta}, x_{1-\alpha}) + g_i(x_\alpha, y_\beta)}{2}.$$

Therefore, p_i satisfies (D4).

Step 3. Since σ^n is locally finite, for any $x_{1-\alpha}, y_{1-\beta} \in M_0$ there are two open sets: an open neighborhood $U_{x_{1-\alpha}}$ of $x_{1-\alpha}$ and an open neighborhood $U_{y_{1-\beta}}$ of $y_{1-\beta}$ such that they are quasi-coincident with only a finite family $\{C_p \mid p = p_1, p_2, \dots, p_l\} \subseteq \sigma^n$ and a finite set $\{B_j \mid j = j_1, j_2, \dots, j_m\} \subseteq \sigma^n$, respectively. Let

$$\{A_i^n \mid i = k_1, k_2, \dots, k_q\} = \{C_p \mid p = p_1, p_2, \dots, p_l\} \cup \{B_j \mid j = j_1, j_2, \dots, j_m\}.$$

Therefore, either $U_{x_{1-\alpha}}(z_i) + A_i^n(z_i) > 1$ or $U_{y_{1-\beta}}(z_i) + A_i^n(z_i) > 1$ is true for each i ($i = k_1, k_2, \dots, k_q$) and corresponding $z_i \in X$. This implies that there exists at most a finite family $\{A_i^n \mid i = k_1, k_2, \dots, k_q\}$ such that $x_\alpha \in A_i^n$ or $y_\beta \in A_i^n$. It follows that $f_i(x_\alpha) \leq 1$ or $f_i(y_\beta) \leq 1$ ($i = k_1, k_2, \dots, k_q$). On the other hand, when $k \in \Gamma_n$ and $k \neq i$ ($i = k_1, k_2, \dots, k_q$), it is easy to show that $x_\alpha \notin A_k^n$ and $y_\beta \notin A_k^n$, and then $f_k(x_\alpha) = 1$ and $f_k(y_\beta) = 1$. Hence $g_k(x_\alpha, y_\beta) = 0$. In other words, when $i \in \Gamma_n$, it may be correct that $g_i(x_\alpha, y_\beta) \neq 0$ only if i belongs to the finite index set $\{k_1, k_2, \dots, k_q\}$.

Similarly, for any $x_\alpha, y_\beta \in M_0$, there exists at most a finite family $\{A_i^n \mid i = q_1, q_2, \dots, q_m\}$ such that $x_\alpha \leq (A_i^n)'$ or $y_\beta \leq (A_i^n)'$. It follows that $f_i^*(x_\alpha) \neq 0$ or $f_i^*(y_\beta) \neq 0$ holds ($i = q_1, q_2, \dots, q_m$). When $k \in \Gamma_n$ and $k \neq i$ ($i = q_1, q_2, \dots, q_m$), we have $x_\alpha \leq (A_i^n)'$ and $y_\beta \leq (A_i^n)'$, and then $f_k^*(x_\alpha) = 0$ and $f_k^*(y_\beta) = 0$. Therefore, $g_k^*(x_\alpha, y_\beta) = 0$. In other words, when $i \in \Gamma_n$, it may be correct that $g_i^*(x_\alpha, y_\beta) \neq 0$ only if i belongs to the finite index set $\{q_1, q_2, \dots, q_m\}$.

Let $J = \{k_1, k_2, \dots, k_q\} \cup \{q_1, q_2, \dots, q_m\}$. Then for any $x_\alpha, y_\beta \in M_0$, there exists at most a finite index set J such that when $i \in J$, it may be correct that $p_i(x_\alpha, y_\beta) \neq 0$. Therefore, for the two positive integers m, n , we can define a mapping $p_{m,n} : M_0 \times M_0 \rightarrow [0, +\infty)$ by

$$p_{m,n}(x_\alpha, y_\beta) = \sum \{p_i(x_\alpha, y_\beta) \mid A_i^n \in \sigma^n, i \in J\}.$$

Next, we will prove that each $p_{m,n}$ is also a Deng pseudo-metric on I^X . The proof is as follows:

(D1) Because each $p_i (i \in J)$ satisfies (D1), when $x_\alpha \geq y_\beta$, we have

$$p_{m,n}(x_\alpha, y_\beta) = \sum \{p_i(x_\alpha, y_\beta) \mid A_i^n \in \sigma^n, i \in J\} = 0.$$

(D2) Let $x_\alpha, y_\beta, z_\gamma \in M_0$. Because p_i satisfies (D2) for each i , we have

$$\begin{aligned} & p_{m,n}(x_\alpha, y_\beta) + p_{m,n}(y_\beta, z_\gamma) \\ &= \sum \{p_i(x_\alpha, y_\beta) \mid A_i^n \in \sigma^n, i \in J\} + \sum \{p_i(y_\beta, z_\gamma) \mid A_i^n \in \sigma^n, i \in J\} \\ &= \sum \{p_i(x_\alpha, y_\beta) + p_i(y_\beta, z_\gamma) \mid A_i^n \in \sigma^n, i \in J\} \\ &\geq \sum \{p_i(x_\alpha, z_\gamma) \mid A_i^n \in \sigma^n, i \in J\} \\ &= p_{m,n}(x_\alpha, z_\gamma). \end{aligned}$$

(D3) Because g_i and g_i^* satisfy (D3), i.e., we have the following two equalities:

$$\begin{aligned} \sum \{g_i(x_\alpha, y_\beta) \mid A_i^n \in \sigma^n, i \in J\} &= \sum \{ \bigwedge_{\gamma > \beta} g_i(x_\alpha, y_\gamma) \mid A_i^n \in \sigma^n, i \in J\}, \\ \sum \{g_i^*(x_\alpha, y_\beta) \mid A_i^n \in \sigma^n, i \in J\} &= \sum \{ \bigwedge_{\gamma > \beta} g_i^*(x_\alpha, y_\gamma) \mid A_i^n \in \sigma^n, i \in J\}. \end{aligned}$$

Note that the above two formulas are finite sums. Therefore,

$$\begin{aligned} & p_{m,n}(x_\alpha, y_\beta) \\ &= \sum \{p_i(x_\alpha, y_\beta) \mid A_i^n \in \sigma^n, i \in J\} \\ &= \sum \{[g_i(x_\alpha, y_\beta) + g_i^*(x_\alpha, y_\beta)]/2 \mid A_i^n \in \sigma^n, i \in J\} \\ &= \sum \{g_i(x_\alpha, y_\beta)/2 \mid A_i^n \in \sigma^n, i \in J\} + \sum \{g_i^*(x_\alpha, y_\beta)/2 \mid A_i^n \in \sigma^n, i \in J\} \\ &= \sum \{ \bigwedge_{\gamma > \beta} g_i(x_\alpha, y_\gamma)/2 \mid A_i^n \in \sigma^n, i \in J\} + \sum \{ \bigwedge_{\gamma > \beta} g_i^*(x_\alpha, y_\gamma)/2 \mid A_i^n \in \sigma^n, i \in J\} \\ &= \sum \{ (\bigwedge_{\gamma > \beta} g_i(x_\alpha, y_\gamma) + g_i^*(x_\alpha, y_\gamma)) / 2 \mid A_i^n \in \sigma^n, i \in J\} \\ &= \bigwedge_{\gamma > \beta} \sum \{ (g_i(x_\alpha, y_\gamma) + g_i^*(x_\alpha, y_\gamma)) / 2 \mid A_i^n \in \sigma^n, i \in J\} \\ &= \bigwedge_{\gamma > \beta} p_{m,n}(x_\alpha, y_\gamma). \end{aligned}$$

(D4) Because p_i satisfies (D4) for each $i \in J$, we have the following equalities:

$$\begin{aligned} & p_{m,n}(x_\alpha, y_\beta) \\ &= \sum \{p_i(x_\alpha, y_\beta) \mid A_i^n \in \sigma^n, i \in J\} \\ &= \sum \{p_i(y_{1-\beta}, x_{1-\alpha}) \mid A_i^n \in \sigma^n, i \in J\} \\ &= p_{m,n}(y_{1-\beta}, x_{1-\alpha}). \end{aligned}$$

Therefore, $\{p_{m,n} \mid m \in \omega, n \in \omega\}$ is a countable family of Deng pseudo-metrics. Meanwhile, we denote the topology generated by $p_{m,n}$ as $\delta_{p_{m,n}}$.

Step 4. We will prove that $\bigcup \delta_{p_{m,n}}$ is a base of (X, δ) . For this purpose, we only need to prove the following (a) and (b).

(a) $\delta_{p_{m,n}} \subseteq \delta (m, n \in \omega)$.

By Theorem 1, it is sufficient to find an open set $V_{x_\alpha} \in \delta$ such that $x_\alpha \in V_{x_\alpha} \subseteq U_\varepsilon(x_\alpha)$ for any open set $U_\varepsilon(x_\alpha) \in \delta_{p_{m,n}}$.

Since σ^n is locally finite, there is an open neighborhood $U_{x_{1-\alpha}}$ of $x_{1-\alpha}$ which is only quasi-coincident with finitely many members: $\{A_i^n \mid i = 1, 2, \dots, k\} \subseteq \sigma^n$. Therefore,

$$f_{i_1}(x_\alpha) \leq 1, \dots, f_{i_k}(x_\alpha) \leq 1. \tag{1}$$

Since $f_{i_l}(x_\alpha) = \inf\{r_{i_l} \in Q_{[0,1]} \mid x_\alpha \in U_{r_{i_l}}\} = t_{i_l}$ ($l = 1, 2, \dots, k$), we may select an open set $U_{r_{i_l}}$ with $x_\alpha \in U_{r_{i_l}}$ such that $r_{i_l} - t_{i_l} < \frac{\varepsilon}{2k}$. Therefore, when $y_\beta \in U_{r_{i_l}}$, we have $f_{i_l}(y_\beta) \leq r_{i_l}$ ($l = 1, 2, \dots, k$). Thus

$$g_{i_l}(x_\alpha, y_\beta) = \max\{f_{i_l}(y_\beta) - f_{i_l}(x_\alpha), 0\} \leq \max\{r_{i_l} - t_{i_l}, 0\} < \frac{\varepsilon}{2k}, l = 1, \dots, k.$$

when $A_{i_m}^n \in \sigma^n$ with $m \neq 1, 2, \dots, k$, it must hold $f_{i_m}(x_\alpha) = 1$, and then $g_{i_m}(x_\alpha, y_\beta) = \max\{f_{i_m}(y_\beta) - f_{i_m}(x_\alpha), 0\} = 0$.

Similarly, there is an open neighborhood U_{x_α} of x_α which is only quasi-coincident with finitely many members: $\{A_{j_t}^n \mid t = 1, 2, \dots, p\} \subseteq \sigma^n$. This implies that $x_\alpha \not\leq (A_{j_t}^n)'$ for each j_t ($t = 1, 2, \dots, p$), so that

$$f_{j_1}^*(x_\alpha) > 0, \dots, f_{j_p}^*(x_\alpha) > 0. \tag{2}$$

Let $f_{j_t}^*(x_\alpha) = \inf\{r_{j_t} \in Q_{[0,1]} \mid x_\alpha \leq V_{j_t}\} = s_{j_t}$ ($t = 1, 2, \dots, p$). Then, we can select an open set $V_{r_{j_t}}$ with $x_\alpha \leq V_{r_{j_t}}$ ($t = 1, 2, \dots, p$) such that $r_{j_t} - s_{j_t} < \frac{\varepsilon}{4p}$. Take a number h_{j_t} of $Q_{[0,1]}$ satisfying $r_{j_t} < h_{j_t} < r_{j_t} + \frac{\varepsilon}{4p}$ such that $x_\alpha \leq V_{h_{j_t}}$ ($t = 1, 2, \dots, p$). Hence $h_{j_t} - s_{j_t} < \frac{\varepsilon}{2p}$ ($t = 1, 2, \dots, p$). Because $r_{j_t} < h_{j_t}$, it is true that $U_{1-h_{j_t}} \leq U_{1-r_{j_t}}$. Therefore, by the property of $\{U_r \mid r \in Q_{[0,1]}\}$ we have $\bar{U}_{1-h_{j_t}} \leq U_{1-r_{j_t}}$, i.e., $U'_{1-r_{j_t}} \leq (\bar{U}_{1-h_{j_t}})'$. Furthermore, because $U_{1-h_{j_t}} \leq \bar{U}_{1-h_{j_t}}$, it is true that $(\bar{U}_{1-h_{j_t}})' \leq U'_{1-h_{j_t}}$. Hence $x_\alpha \leq V_{r_{j_t}} \leq (\bar{U}_{1-h_{j_t}})' \leq V_{h_{j_t}}$. Let $O_{h_{j_t}} = V_{h_{j_t}}^\circ, t = 1, 2, \dots, p$. Then $x_\alpha \leq (\bar{U}_{1-h_{j_t}})' \leq O_{h_{j_t}}$. If $z_\gamma \in O_{h_{j_t}} \leq V_{h_{j_t}}, t = 1, 2, \dots, p$, then $f_{j_t}^*(z_\gamma) \leq h_{j_t}$. Hence

$$g_{j_t}^*(x_\alpha, z_\gamma) = \max\{f_{j_t}^*(z_\gamma) - f_{j_t}^*(x_\alpha), 0\} \leq \max\{h_{j_t} - s_{j_t}, 0\} < \frac{\varepsilon}{2p}, t = 1, \dots, p.$$

Now, let

$$V_{x_\alpha} = U_{x_\alpha} \wedge U_{r_{i_1}} \wedge \dots \wedge U_{r_{i_k}} \wedge O_{h_{j_1}} \wedge \dots \wedge O_{h_{j_p}}. \tag{3}$$

Since U_{x_α} and $A_{j_s}^n$ are not quasi-coincident for each j_s ($s \neq 1, 2, \dots, p$), when $y_\beta \in V_{x_\alpha}$, we have $y_\beta \leq (A_{j_s}^n)'$ and then $f_{j_s}^*(y_\beta) = 0$ ($s \neq 1, 2, \dots, p$). And, consequently, $g_{j_s}^*(x_\alpha, y_\beta) = 0$ ($s \neq 1, 2, \dots, p$). If $k \leq p$, then

$$p_{m,n}(x_\alpha, y_\beta) \leq \left[\frac{(\frac{\varepsilon}{2k} + \frac{\varepsilon}{2p})}{2}\right] \times k + \frac{0 + \frac{\varepsilon}{2p}}{2} \times (p - k) = \frac{\varepsilon}{4} + \frac{\varepsilon}{4} < \varepsilon.$$

If $k > p$, then

$$p_{m,n}(x_\alpha, y_\beta) \leq \left[\frac{(\frac{\varepsilon}{2k} + \frac{\varepsilon}{2p})}{2}\right] \times p + \frac{\varepsilon}{2k} \times (k - p) = \frac{\varepsilon}{4} + \frac{\varepsilon}{4k}(2k - p) < \frac{\varepsilon}{4} + \frac{\varepsilon}{2} < \varepsilon.$$

In either case, $x_\alpha \in V_{x_\alpha} \leq U_\varepsilon(x_\alpha)$. Therefore, (a) is proved.

Incidentally, to make the above proof more perfect, we add the following two points. If $f_{i_l}(x_\alpha) = 1$ for all i_l ($l = 1, \dots, k$), then $g_{i_l}(x_\alpha, y_\beta) = 0$. Meanwhile, let us consider two more special cases below.

(i) If there exists a nonempty set $\Lambda = \{j_{w_1}, \dots, j_{w_q}\} \subseteq \{j_1, \dots, j_p\}$ such that each element $j_{w_i} \in \Lambda$ satisfies $0 < f_{j_{w_i}}^*(x_\alpha) < 1$, then

$$V_{x_\alpha} = U_{x_\alpha} \wedge O_{j_{w_1}} \wedge \dots \wedge O_{j_{w_q}}.$$

(ii) If $f_{j_t}^*(x_\alpha) = 1$ for all j_t ($t = 1, \dots, p$), then $g_{j_s}^*(x_\alpha, y_\beta) = 0$. Let $V_{x_\alpha} = U_{x_\alpha}$. Then for any $y_\beta \in U_{x_\alpha}$ we have $p_{m,n}(x_\alpha, y_\beta) = 0$, and thus $y_\beta \in U_\varepsilon(x_\alpha)$, i.e., $x_\alpha \in V_{x_\alpha} \leq U_\varepsilon(x_\alpha)$.

(b) Each member in δ is the union of some members in $\cup \delta_{p_{m,n}}$.

Let $x_\alpha \in B \in \delta$. Because (X, δ) is regular, there exists an open set v belonging to σ such that $x_\alpha \in v \leq \bar{v} \leq B$. Therefore, it is easy to show that there is a natural member n such that $v \in \sigma^n \subseteq \sigma$. For convenience, we denote v as A_i^n . Similarly, for $x_\alpha \in A_i^n$, there are another natural member m and an open set A_j^m belonging to σ^m such that

$$x_\alpha \in A_j^m \leq \overline{A_j^m} \leq A_i^n \leq \overline{A_i^n} \leq B.$$

Let $A_i = \bigvee \{A^m \mid A^m \in \sigma^m, \overline{A^m} \leq A_i^n\}$. Clearly, $x_\alpha \in A_j^m \leq A_i \leq \overline{A_i} \leq A_i^n \leq \overline{A_i^n} \leq B$. Therefore, by Theorem 22 there exists a corresponding family $\{U_r \mid r \in Q_{[0,1]}\}$ relative to $\overline{A_i}$ and B such that $x_\alpha \in U_r$ for all U_r ($r \in Q_{[0,1]}$). And consequently $f_i(x_\alpha) = 0$. If $y_\beta \notin B$, then $y_\beta \notin A_i^n$. Thus $f_i(y_\beta) = 1$, and then $g_i(x_\alpha, y_\beta) = \max\{f_i(y_\beta) - f_i(x_\alpha), 0\} = 1$. Therefore, we assert that $p_{m,n}(x_\alpha, y_\beta) \geq \frac{1}{2}$. In other words, as long as $p_{m,n}(x_\alpha, y_\beta) < \frac{1}{2}$, it must hold that $y_\beta \in B$. This implies that for each $x_\alpha \in B$ there exists $U_{\frac{1}{2}}(x_\alpha)$ belonging to $\delta_{p_{m,n}}$ such that $U_{\frac{1}{2}}(x_\alpha) \leq B$. Thus $B = \bigvee_{x_\alpha \in B} U_{\frac{1}{2}}(x_\alpha)$. That is to say, if $B \in \delta$, then there is $D \subseteq \bigcup_{n,m \in \omega} \delta_{p_{m,n}}$ such that $B = \bigvee D$. Therefore, (b) is proved.

Step 5. Based on the discussions above, we renumber the countable set $\{p_{m,n} \mid m = 1, 2, \dots, n = 1, 2, \dots\}$ as $\{p_n \mid n \in \omega\}$. Let $\mathcal{X} = \prod_{n \in \omega, X_n = X} X_n$. By Theorem 4, we define a mapping $p : M_0 \times M_0 \rightarrow [0, 1]$ by

$$p(x_\alpha, y_\beta) = \sum_{n \in \omega} 2^{-n} p_n(J_n(x_\alpha), J_n(y_\beta)),$$

where $J_n : I^X \rightarrow I^X$ is the n -th projection, and affirm that p is a Deng pseudo-metric on I^X and (\mathcal{X}, δ_p) is the product space of $\{(X, \delta_{p_n}) \mid n \in \omega\}$, where (\mathcal{X}, δ_p) is generated by $\Gamma_p = \{J_n^{-1}(U) \mid U \in \delta_{p_n}, n \in \omega\}$ as a subbase. Now let us prove that (X, δ) can be embedded into (\mathcal{X}, δ_p) .

Let $x^\omega = (x, x, \dots, x, \dots)$ and denote x_α^ω as the fuzzy point whose support and value are x^ω and $\alpha \in (0, 1)$, respectively. All these fuzzy points are denoted by $M_1 = \{x_\alpha^\omega \mid x \in X, \alpha \in (0, 1)\}$. Let $\mathbb{X} = \{x^\omega \mid x \in X\}$. A mapping $e : M_1 \rightarrow M_0$ is defined by $e(x_\alpha^\omega) = x_\alpha$. Obviously, e is a bijection and its inverse mapping e^{-1} embeds M_0 into I^X . Let $p_e = p \mid M_1 \times M_1$. Consequently, we regenerate a new mapping $p_e : M_1 \times M_1 \rightarrow [0, 1]$. It is easy to prove that p_e is a Deng pseudo-metric on $I^{\mathbb{X}}$, and $(\mathbb{X}, \delta_{p_e})$ is a subspace of (\mathcal{X}, δ_p) . Because Γ_p is a subbase of (\mathcal{X}, δ_p) , $\Gamma_p \mid \mathbb{X}$ is certainly a subbase of $(\mathbb{X}, \delta_{p_e})$. Moreover, because of Step 4, $\Gamma_p \mid \mathbb{X}$ is exactly a base of $(\mathbb{X}, \delta_{p_e})$. Hence (X, δ) and $(\mathbb{X}, \delta_{p_e})$ are homeomorphic. In fact, let $p_{e_1}(x_\alpha, y_\beta) = p_e(x_\alpha^\omega, y_\beta^\omega)$ for any $x_\alpha^\omega, y_\beta^\omega \in M_1$. Then (X, δ) can be embedded into (\mathcal{X}, δ_p) and the mapping p_{e_1} is a Deng pseudo-metric on I^X , which metricizes the $[0, 1]$ -topological space (X, δ) . Consequently, $\delta = \delta_{p_{e_1}}$. In summary, the proof has been completed. \square

Theorem 24. A $[0, 1]$ -topological space is Deng-metrizable if and only if it is T_1 and Deng pseudo-metrizable.

Proof. (Sufficiency). Let p be a Deng metric on I^X and let $y_{\lambda_2} \leq \bigwedge_{r>0} B_r(x_{\lambda_1})$ (see Theorem 7 on B_r). Then for any $r > 0$ we have $y_{\lambda_2} \leq B_r(x_{\lambda_1})$ and then $p(x_{\lambda_1}, y_{\lambda_2}) \leq r$, so that $p(x_{\lambda_1}, y_{\lambda_2}) = 0$. By (D5) we can obtain $x = y$ and $\lambda_1 \geq \lambda_2$. Hence $\bigwedge_{r>0} B_r(x_{\lambda_1}) = x_{\lambda_1}$. In addition, by Theorem 7 we can assert that $B_r(x_{\lambda_1})$ ($r > 0$) is a closed set. Thus, $\bigwedge_{r>0} \overline{B_r(x_{\lambda_1})} = \bigwedge_{r>0} B_r(x_{\lambda_1})$. Consequently, $\bar{x}_{\lambda_1} = x_{\lambda_1}$, as desired.

(Necessity). If $p(x_{\lambda_1}, y_{\lambda_2}) = 0$, then $p(y_{1-\lambda_2}, x_{1-\lambda_1}) = 0$. For any $r > 0$, by (D3) we can take a number $1 - \lambda_r > 1 - \lambda_1$ such that $p(y_{1-\lambda_2}, x_{1-\lambda_r}) < r$, and then $x_{1-\lambda_1} < x_{1-\lambda_r} \leq$

$U_r(y_{1-\lambda_2})$, i.e., $x_{\lambda_1} + U_r(y_{1-\lambda_2}) > 1$. Consequently $y_{\lambda_2} \leq \bar{x}_{\lambda_1} = x_{\lambda_1}$. Therefore, $x = y$ and $\lambda_1 \geq \lambda_2$, so that p satisfies (D5). \square

Because of the conclusion $D = C \cap Y \cap E$ in Theorem 8, Theorem 22 and Theorem 23, we can obtain the main result in this paper as follow

Corollary 1. *If a $[0, 1]$ -topological space (X, δ) is T_1 and regular, and δ has a σ -locally finite base, then it is Deng-, Erceg-, Chen-, and Shi-metrizable.*

8. Conclusions

In this paper, we study the metrization problem: whether there is a metric such that a given $[0, 1]$ -topology coincides with the metric topology. Eventually, we obtain a desired result:

If a $[0, 1]$ -topological space (X, δ) is T_1 and regular, and δ has a σ -locally finite base, then it is Deng-, Erceg-, Chen-, and Shi-metrizable.

Based on the result, we can conclude that Deng's, Liang's, and Yang's metric results, which appeared in the Introduction (refer to [13,24,30] for details), are all special cases of our proposition. This is because if (X, δ) is C_{II} , then δ must have a σ -locally finite base, but the converse is not true. Therefore, Corollary 1 proved by us is the most satisfactory solution to the metrization problem in $[0, 1]$ -topology so far.

In the future, we will consider whether or not our results can be generalized to L -topology [9,15]. In addition, we will further investigate Erceg metric, Shi metric, Deng metric, and Chen metric. Additionally, we will continue to research the kind of lattice-valued topological spaces each of whose topologies has a σ -locally finite base. Beyond that, we also intend to inquire into some questions on the fuzzifying metric topology (see [16,33,39,45]).

Funding: This research received no external funding.

Acknowledgments: The author wishes to express deep gratitude to Fu-Gui Shi from Beijing Institute of Technology, Wei Yao from Nanjing University of Information Science and Technology, Yue-Li Yue from the Ocean University of China, and Yue Huang from the Department of Foreign Languages, Guangdong University of Technology, for a number of very valuable suggestions and improvements.

Conflicts of Interest: The author declares no conflict of interest.

References

- Bing, R.H. Metrization of topological spaces. *Can. J. Math.* **1951**, *3*, 175–186. [\[CrossRef\]](#)
- Dowker, C.H. An embedding theorem for paracompact metric spaces. *Duke Math. J.* **1947**, *14*, 639–645. [\[CrossRef\]](#)
- Nagata, J. On a necessary and sufficient condition of metrizability. *J. Inst. Polytech.* **1950**, *1*, 93–100.
- Stone, A.H. Paracompactness and product spaces. *Bull. Am. Math. Soc.* **1948**, *54*, 977–982. [\[CrossRef\]](#)
- Chang, C.L. Fuzzy topological spaces. *J. Math. Anal. Appl.* **1968**, *24*, 182–190. [\[CrossRef\]](#)
- Zadeh, L.A. Fuzzy sets. *Inform. Control* **1965**, *8*, 338–353. [\[CrossRef\]](#)
- Goguen, J.A. The fuzzy Tychonoff Theorem. *J. Math. Anal. Appl.* **1973**, *18*, 734–742. [\[CrossRef\]](#)
- Chen, P.; Meng, B.; Ba, X. L-Quasi (Pseudo)-Metric in L-Fuzzy Set Theory. *Mathematics* **2023**, *11*, 3152. [\[CrossRef\]](#)
- Chen, P. *Metrics in L-Fuzzy Topology*; Published by Science Press: Beijing, China, 2017; Volume 12. (In Chinese)
- Chen, P.; Duan, P. Research of Deng metric and its related problems. *Fuzzy Syst. Math.* **2015**, *29*, 28–35. (In Chinese)
- Chen, P.; Duan, P. Research on a kind of pointwise parametric in L lattices. *Fuzzy Syst. Math.* **2016**, *30*, 23–30. (In Chinese)
- Chen, P.; Shi, F.G. Further simplification of Erceg metric and its properties. *Adv. Math.* **2007**, *36*, 586–592. (In Chinese)
- Deng, Z.K. Fuzzy pseudo-metric spaces. *J. Math. Anal. Appl.* **1982**, *86*, 74–95. [\[CrossRef\]](#)
- Deng, Z.K. M-uniformization and metrization of fuzzy topological spaces. *J. Math. Anal. Appl.* **1985**, *112*, 471–486. [\[CrossRef\]](#)
- Erceg, M.A. Metric spaces in fuzzy set theory. *J. Math. Anal. Appl.* **1979**, *69*, 205–230. [\[CrossRef\]](#)
- George, A.; Veeramani, P. On some results in fuzzy metric spaces. *Fuzzy Sets Syst.* **1994**, *64*, 395–399. [\[CrossRef\]](#)
- Gregori, V.; Morillas, S.; Sapena, A. Examples of fuzzy metrics and applications. *Fuzzy Sets Syst.* **2011**, *170*, 95–111. [\[CrossRef\]](#)
- Gregori, V.; Sapena, A. On fixed point theorems in fuzzy metric spaces. *Fuzzy Sets Syst.* **2002**, *125*, 245–252. [\[CrossRef\]](#)
- Diamond, P.; Kloeden, P. Metric spaces of fuzzy sets. *Fuzzy Sets Syst.* **1990**, *35*, 241–249. [\[CrossRef\]](#)
- Smirnov, Y.M. A necessary and sufficient condition for metrizability of a topological space. *Dokl. Akad. Nauk* **1951**, *77*, 197–200.
- Hutton, B. Uniformities on fuzzy topological spaces. *J. Math. Anal. Appl.* **1977**, *58*, 559–571. [\[CrossRef\]](#)

22. Kim, D.S.; Kim, Y.K. Some properties of a new metric on the space of fuzzy numbers. *Fuzzy Sets Syst.* **2004**, *145*, 395–410. [[CrossRef](#)]
23. Liang, J.H. A few problems in fuzzy metric spaces. *Ann. Math.* **1984**, *6A*, 59–67. (In Chinese)
24. Liang, J.H. Pointwise characterizations of fuzzy metrics and its applications. *Acta Math. Sin.* **1987**, *30*, 733–741. (In Chinese)
25. Luo, M.K. A note on fuzzy paracompact and fuzzy metric. *J. Sichuan Univ.* **1985**, *4*, 141–150. (In Chinese)
26. Meng, B.; Chen, P.; Ba, X. Expansion Theory of Deng's Metric in $[0,1]$ -Topology. *Mathematics* **2023**, *11*, 3414. [[CrossRef](#)]
27. Morillas, S.; Gregori, V.; Peris-Fajarnés, G. A fast impulsive noise color image filter using fuzzy metrics. *Real Time Imaging* **2005**, *11*, 417–428. [[CrossRef](#)]
28. Shi, F.G. Pointwise pseudo-metrics in L -fuzzy set theory. *Fuzzy Sets Syst.* **2001**, *121*, 209–216. [[CrossRef](#)]
29. Šostak, A.P. Basic structures of fuzzy topology. *J. Math. Sci.* **1996**, *78*, 662–701. [[CrossRef](#)]
30. Yang, L.C. Theory of p.q. metrics on completely distributive lattices. *Chin. Sci. Bull.* **1988**, *33*, 247–250. (In Chinese)
31. Yue, Y.; Shi, F.G. On fuzzy pseudo-metric spaces. *Fuzzy Sets Syst.* **2010**, *161*, 1105–1116. [[CrossRef](#)]
32. Eklund, P.; Gäbler, W. Basic notions for topology I/II. *Fuzzy Sets Syst.* **1988**, *26–27*, 333–356/171–195.
33. Kramosil, I.; Michalek, J. Fuzzy metric statistical metric spaces. *Kybernetika* **1975**, *11*, 336–344.
34. Morsi, N.N. On fuzzy pseudo-normed vector spaces. *Fuzzy Sets Syst.* **1988**, *27*, 351–372. [[CrossRef](#)]
35. Adibi, H.; Cho, Y.; O'regan, D.; Saadati, R. Common fixed point theorems in L -fuzzy metric spaces. *Appl. Math. Comput.* **2006**, *182*, 820–828. [[CrossRef](#)]
36. Al-Mayahi, N.F.; Ibrahim, L.S. Some properties of two-fuzzy metric spaces. *Gen. Math. Notes* **2013**, *17*, 41–52.
37. Çayh, G.D. On the structure of uninorms on bounded lattices. *Fuzzy Sets Syst.* **2019**, *357*, 2–26.
38. George, A.; Veeramani, P. On some results of analysis for fuzzy metric spaces. *Fuzzy Sets Syst.* **1997**, *90*, 365–368. [[CrossRef](#)]
39. Gregori, V.; Romaguera, S. Some properties of fuzzy metric spaces. *Fuzzy Sets Syst.* **2000**, *115*, 485–489. [[CrossRef](#)]
40. Hua, X.J.; Ji, W. Uninorms on bounded lattices constructed by t-norms and t-subconorms. *Fuzzy Sets Syst.* **2022**, *427*, 109–131. [[CrossRef](#)]
41. Grabiec, M. Fixed points in fuzzy metric spaces. *Fuzzy Sets Syst.* **1988**, *27*, 385–389. [[CrossRef](#)]
42. Morillas, S.; Gregori, V.; Peris-Fajarnés, G.; Sapena, A. New adaptive vector filter using fuzzy metrics. *J. Electron. Imaging* **2007**, *16*, 033007. [[CrossRef](#)]
43. Sharma, S. Common fixed point theorems in fuzzy metric spaces. *Fuzzy Sets Syst.* **2002**, *127*, 345–352. [[CrossRef](#)]
44. Shi, F.G. (L, M) -fuzzy metric spaces. *Indian J. Math.* **2010**, *52*, 231–250.
45. Shi, F.G. L -metric on the space of L -fuzzy numbers. *Fuzzy Sets Syst.* **2020**, *399*, 95–109. [[CrossRef](#)]
46. Yager, R.R. Defending against strategic manipulation in uninorm-based multi-agent decision making. *Fuzzy Sets Syst.* **2003**, *140*, 331–339. [[CrossRef](#)]
47. Peng, Y.W. Simplification of Erceg fuzzy metric function and its application. *Fuzzy Sets Syst.* **1993**, *54*, 181–189.
48. Chen, P.; Shi, F.G. A note on Erceg pseudo-metric and pointwise pseudo-metric. *J. Math. Res. Exp.* **2008**, *28*, 339–443.
49. Gierz, G.; Hofmann, K.H.; Keimel, K.; Lawson, J.D.; Mislove, M.; Scott, D.S. *A Compendium of Continuous Lattices*; Springer: Berlin/Heidelberg, Germany, 1980.
50. Wang, G.J. *Theory of L-Fuzzy Topological Spaces*; Shaanxi Normal University Press: Xi'an, China, 2008. (In Chinese)
51. Shi, F.G. Pointwise quasi-uniformities and p.q. metrics on completely distributive lattices. *Acta Math. Sin.* **1996**, *39*, 701–706. (In Chinese)
52. Shi, F.G.; Zheng, C.Y. Metrization theorems on L -topological spaces. *Fuzzy Sets Syst.* **2005**, *149*, 455–471. [[CrossRef](#)]
53. Lan, Y.Y.; Long, F. The sufficiency and necessity conditions for metrization of the fuzzy topological spaces. *J. Hunan City Univ.* **2006**, *15*, 37–39. (In Chinese)
54. Chen, P.; Qiu, X. Expansion theorem of Deng metric. *Fuzzy Syst. Math.* **2019**, *33*, 54–65. (In Chinese)
55. Pu, P.M.; Liu, Y.M. Fuzzy topology I. neighborhood structure of a fuzzy point and Moore-Smith convergence. *J. Math. Anal. Appl.* **1980**, *76*, 571–599.
56. Kelley, J.L. *General Topology*, 1955 ed.; Van Nostrand: New York, NY, USA, 1955.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.