



Article

# On Properties of Karamata Slowly Varying Functions with Remainder and Their Applications

Azam A. Imomov <sup>1,\*</sup>, Erkin E. Tukhtaev <sup>1,†</sup> and János Sztrik <sup>2,†</sup>

<sup>1</sup> Department of Algebra and Geometry, Karshi State University, Karshi City 180100, Uzbekistan; tukhtaev\_erkin@mail.ru

<sup>2</sup> Faculty of Informatics, University of Debrecen, 4032 Debrecen, Hungary; sztrik.janos@inf.unideb.hu

\* Correspondence: imomov\_azam@mail.ru

† These authors contributed equally to this work.

**Abstract:** In this paper, we study the asymptotic properties of slowly varying functions of one real variable in the sense of Karamata. We establish analogs of fundamental theorems on uniform convergence and integral representation for slowly varying functions with a remainder depending on the types of remainder. We also prove several important theorems on the asymptotic representation of integrals of Karamata functions. Under certain conditions, we observe a “narrowing” of classes of slowly varying functions concerning the types of remainder. At the end of the paper, we discuss the possibilities of the application of slowly varying functions in the theory of stochastic branching systems. In particular, under the condition of the finiteness of the moment of the type  $\mathbb{E}[x \ln x]$  for the particle transformation intensity, it is established that the property of slow variation with a remainder is implicitly present in the asymptotic structure of a non-critical Markov branching random system.

**Keywords:** slowly varying function; integral representation; remainder; Landau symbols; stochastic branching systems; criticality; invariant distributions

**MSC:** 26A12; 60J80



**Citation:** Imomov, A.A.; Tukhtaev, E.E.; Sztrik, J. On Properties of Karamata Slowly Varying Functions with Remainder and Their Applications. *Mathematics* **2024**, *12*, 3266. <https://doi.org/10.3390/math12203266>

Academic Editor: Ioannis K. Argyros

Received: 29 August 2024

Revised: 10 October 2024

Accepted: 12 October 2024

Published: 18 October 2024



**Copyright:** © 2024 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

## 1. Introduction

The concept of regular variation, initiated by the famous Serbian mathematician Jovan Karamata in the early 1930s, is a special one-sided locally asymptotic property of functions of a real variable. It arose from the desire to logically extend the class of functions with power-law asymptotic monotonicity near some point to a class of functions with behavior akin to a power function multiplied by a coefficient changing “more slowly” than the power function. The first book in the world of mathematical literature specifically devoted to the systematic study of Karamata functions was published by the Australian mathematician E. Seneta [1] in 1976. Detailed materials related to the application of the theory of regularly varying functions in various areas of mathematics can be found in the monographs [2,3]. The possibilities of applying regularly varying functions in the theory of Markov branching random systems were first discussed in the work of V.Zolotarev [4].

A real-valued, positive, and measurable function  $L(x)$  is said to be slowly varying (SV) at infinity (in the sense of Karamata) if  $L(\lambda x)/L(x) \rightarrow 1$  as  $x \rightarrow \infty$  for an arbitrary  $\lambda \in \mathbb{R}_+$ , where  $\mathbb{R}_+$  denotes the positive semiaxis of real numbers. In what follows, we use the symbol  $\mathcal{SV}_\infty$  to denote the class of SV-functions at infinity. It is easy to see that if  $L(x) \in \mathcal{SV}_\infty$ , then  $L(1/x)$  is slowly varying at zero. Thus, one can define the slowly varying conception at any finite point by shifting the origin to that point. In this regard, we limit ourselves to considering functions from the class  $\mathcal{SV}_\infty$ . The well-known fundamental theorem on integral representation states that any function  $L(x) \in \mathcal{SV}_\infty$  can be represented in the form

$$L(x) = M(x) \exp \int_b^x \frac{\varepsilon(u)}{u} du$$

for some  $b \in \mathbb{R}_+$ , where  $M(x)$  is a bounded measurable function defined on the set  $[b, \infty]$ , so that  $|M(x)| \rightarrow M \in \mathbb{R}_+$ . The function  $\varepsilon(x)$  is continuous and infinitesimal as  $x \rightarrow \infty$ , called the index of variation of  $L(x)$ . In the special case when  $M(x) \equiv const$ , the function  $L(x)$  is called normalized. One of the important characteristics of normalized functions from the class  $\mathcal{SV}_\infty$  is the fact that if it is differentiable, then

$$\frac{xL'(x)}{L(x)} = \varepsilon(x) \rightarrow 0 \quad \text{as } x \rightarrow \infty;$$

see [3] (Ch.1.3, §2).

A function  $R(x)$  is called a regularly varying (RV) function at infinity with regularity index  $\rho$  if it can be represented as  $R(x) = x^\rho L(x)$  for some  $L(x) \in \mathcal{SV}_\infty$ . It follows from the RV-function definition that

$$\frac{R(\lambda x)}{R(x)} \rightarrow \lambda^\rho \quad \text{as } x \rightarrow \infty$$

for an arbitrary  $\lambda \in \mathbb{R}_+$ ; see [1]. Denote by  $\mathcal{R}_\infty^\rho$  the class of RV-functions at infinity. Then, it is obvious that  $\mathcal{SV}_\infty \equiv \mathcal{R}_\infty^0$ .

In this report, we are interested in one important characteristic of SV-functions, indicating the degree of their variation in the class  $\mathcal{SV}_\infty$ . Thus, according to the definition, the function  $\omega_\lambda(x) := L(\lambda x)/L(x) - 1$  has an infinitesimal value at infinity. Research in recent years has shown that the use of Karamata functions with a certain degree of smallness  $\omega_\lambda(x)$  allows us to improve known theorems of probability theory, in particular, to obtain facts about deep properties of stochastic branching systems of various types. In this regard, we turn to the works [5,6], in which practically unimprovable asymptotic estimates of the survival probabilities of a population of individuals in stochastic branching systems with a reproductive law having a finite moment of order  $1 + \nu$  for all  $\nu \in (0, 1)$  were found.

Write

$$\frac{L(\lambda x)}{L(x)} - 1 = \omega_\lambda(x), \tag{1}$$

where  $\omega_\lambda(x) \rightarrow 0$  as  $x \rightarrow \infty$ . The relation (1) points to the fact that the asymptotic properties of the function  $L(x) \in \mathcal{SV}_\infty$  depend on the decreasing rate of  $\omega_\lambda(x)$ .

In what follows, we use Landau symbols  $o, \mathcal{O}, \mathcal{O}^*$  to compare two functions  $f(*)$  and  $h(*)$  at the point  $x_0$  (finite or infinite):

$$\begin{aligned} f(x) = o(h(x)) & \quad \text{if} \quad \frac{|f(x)|}{|h(x)|} \rightarrow 0, \\ f(x) = \mathcal{O}(h(x)) & \quad \text{if} \quad \frac{|f(x)|}{|h(x)|} \leq A, \\ f(x) = \mathcal{O}^*(h(x)) & \quad \text{if} \quad \frac{|f(x)|}{|h(x)|} \rightarrow A, \end{aligned}$$

as  $x \rightarrow x_0$ , where  $A$  is a finite positive constant. Also,  $f(x) \sim h(x)$  means  $f(x)/h(x) \rightarrow 1$ .

In the monograph [3] (Ch.2.3, §1), the functions  $L(x) \in \mathcal{SV}_\infty$  with a remainder term of the form  $\omega_\lambda(x) = o(1/\varphi(x))$ , where  $\varphi(x)$  is an infinitely large function, are considered. It is also proved there that if the function  $L(x)$  is continuously differentiable and the function  $\tau(x) := \exp\{\varphi(x)\}$  does not decrease and  $\tau(x) > 1$ , then the characteristic representation

$$\frac{xL'(x)}{L(x)} = o\left(\frac{1}{\ln \tau(x)}\right) \quad \text{as } x \rightarrow \infty \tag{2}$$

entails the following relation:

$$\omega_\lambda(x) \ln \tau(x) \rightarrow 0 \quad \text{as } x \rightarrow \infty.$$

Further, suppose that  $r(x)$  is defined for  $x \in \mathbb{R}_+$  such that  $r(x) \in \mathbb{R}_+$  and  $r(x) \downarrow 0$  as  $x \rightarrow \infty$ . A function  $L(x) \in \mathcal{SV}_\infty$  is called an SV-function with a remainder if it satisfies for all  $\lambda \in \mathbb{R}_+$  one of the following conditions as  $x \rightarrow \infty$  [3] (Ch.3.12, §1):

- (SR1)  $\omega_\lambda(x) = \mathcal{O}(r(x));$
- (SR2)  $\omega_\lambda(x) \sim \text{const}(\lambda)r(x);$
- (SR3)  $\omega_\lambda(x) = o(r(x)).$

By introducing the notation  $\psi(x) := \ln L(x)$ , the above conditions can be replaced by the following ones:

- (SR1)  $\psi(\lambda x) - \psi(x) = \mathcal{O}(r(x));$
- (SR2)  $\psi(\lambda x) - \psi(x) \sim \text{const}(\lambda)r(x);$
- (SR3)  $\psi(\lambda x) - \psi(x) = o(r(x)).$

The class of SV-functions with a remainder  $\omega_\lambda(x)$  defined by conditions SR1–SR3 is denoted throughout by  $\mathcal{SV}_\infty(\omega)$ .

In Section 2, we study the asymptotic properties of functions from the class of  $\mathcal{SV}_\infty(\omega)$ . We establish analogs of fundamental theorems on uniform convergence and integral representation. We also establish some important asymptotic formulas for improper integrals of RV-functions. In Section 3, we briefly discuss the presence of the slow-varying property in the asymptotic structure of continuous-time stochastic Markov branching systems.

### 2. Main Theorems

We begin by presenting the following theorems, which are important generalizations of the fundamental theorems from [1] (Ch.1, §2) on uniform convergence and the integral representation for functions  $L(x) \in \mathcal{SV}_\infty(\omega)$ .

From now on, we use Landau symbols only concerning the asymptotics as  $x \rightarrow \infty$ , unless otherwise stated.

**Theorem 1.** *Let  $L(x) \in \mathcal{SV}_\infty(\omega)$ . Then, for any  $[a, b] \subset \mathbb{R}_+$ , the conditions SR1–SR3 are satisfied uniformly for all  $\lambda \in [a, b]$*

$$\sup \left\{ |\omega_\lambda(x)| : \lambda \in [a, b] \right\} = \begin{cases} \mathcal{O}(r(x)) & \text{for SR1,} \\ \mathcal{O}^*(r(x)) & \text{for SR2,} \\ o(r(x)) & \text{for SR3.} \end{cases}$$

**Theorem 2.** *The function  $L(x)$  defined for  $x \in [a, \infty) \subset \mathbb{R}_+$  belongs to the class  $\mathcal{SV}_\infty(\omega)$  if and only if for some  $b \in [a, \infty)$  the following integral representation is allowed:*

$$\ln L(x) = \eta(x) + \int_b^x \frac{\varepsilon(u)}{u} du \tag{3}$$

where  $\eta(x)$  is a measurable function on the set  $[b, \infty)$  such that  $\eta(x) \rightarrow C_\eta$ ,  $|C_\eta| < \infty$ , and  $\varepsilon(x)$  is a continuous function on  $[b, \infty)$  such that  $\varepsilon(x) \rightarrow 0$  as  $x \rightarrow \infty$ . Moreover,

$$\left. \begin{matrix} \eta(x) - C_\eta \\ \varepsilon(x) \end{matrix} \right\} = \begin{cases} \mathcal{O}(r(x)) & \text{for SR1,} \\ \mathcal{O}^*(r(x)) & \text{for SR2,} \\ o(r(x)) & \text{for SR3.} \end{cases} \tag{4}$$

**The proof** of the last two theorems is based on the proof scheme of similar theorems given in [1] (*Trans. Rus. Ch. Annex., §A*) for the case  $\omega_\lambda(x) = \mathcal{O}(1/\varphi(x))$  with infinitely large  $\varphi(x)$  as  $x \rightarrow \infty$ . Therefore, we do not dwell on the details of the discussion proof. The superiority of these results over similar theorems for functions from the class  $\mathcal{SV}_\infty$  lies in the fact that the first theorem estimates the degrees of uniform smallness of the remainder  $\omega_\lambda(x)$ , and the second theorem estimates, in particular, the varying index of  $\varepsilon(x)$ . Thus, the nature of the varying of the functions  $\eta(x)$  and  $\varepsilon(x)$  depends on the degree of smallness of the remainder  $\omega_\lambda(x)$  as  $x \rightarrow \infty$ .

Next, we prove the following theorem on the asymptotic behavior of functions from the class  $\mathcal{SV}_\infty(\omega)$ .

**Theorem 3.** *Each function  $L(x) \in \mathcal{SV}_\infty(\omega)$  has a finite limit  $C_L := \lim_{x \rightarrow \infty} L(x)$ . If the condition  $\mathcal{SR}1$  is satisfied with  $r(x) = \mathcal{O}(L(x)/x^\sigma)$ , where  $\sigma \in \mathbb{R}_+$ , then*

$$\eta(x) = C_\eta + \mathcal{O}^*(1/x^\sigma) \quad \text{and} \quad \varepsilon(x) = \mathcal{O}^*(1/x^\sigma).$$

Moreover, the following asymptotic representation is valid:

$$L(x) = C_L + \mathcal{O}^*(1/x^\sigma). \tag{5}$$

Conversely, if for the function  $L(x)$  defined on some interval  $[a, \infty) \subset \mathbb{R}_+$  (5) holds, then  $L(x) \in \mathcal{SV}_\infty(\omega)$  with a remainder term  $\omega_\lambda(x) = \mathcal{O}^*(1/x^\sigma)$ .

**Proof.** In the condition of the theorem, from the corresponding statement in (4) it follows that  $\varepsilon(x) = \mathcal{O}(L(x)/x^\sigma)$ . Since the number  $\sigma \in \mathbb{R}_+$  is positive, then by the properties of SV-functions, we obtain that the improper integral  $\int_b^\infty [\varepsilon(u)/u]du$  converges. Again, from (4), we find  $\eta(x) = C_\eta + \mathcal{O}(L(x)/x^\sigma)$  as  $x \rightarrow \infty$ . Collecting these facts in (3), we have

$$\ln L(\infty) = C_\eta + \int_b^\infty \frac{\varepsilon(u)}{u} du,$$

which is equivalent to  $\lim_{x \rightarrow \infty} L(x) =: C_L < \infty$ . It follows that the remainder term

$$\omega_\lambda(x) = \mathcal{O}^*(1/x^\sigma) \quad \text{as} \quad x \rightarrow \infty.$$

Now, we prove Formula (5). To do this, we write (3) in the form

$$L(x) = \exp\left\{C_\eta + \mathcal{O}^*(1/x^\sigma)\right\} L_0(x), \tag{6}$$

where  $L_0(x)$  is the normalized SV-function associated with  $L(x)$ . Next, we have

$$L_0(\infty) = \int_b^\infty \frac{\varepsilon(u)}{u} du =: C_0 < \infty,$$

which follows from the fact that  $L(\infty) =: C_L < \infty$ . Now, we obtain

$$C_0 - L_0(x) = C_0 \left(1 - \frac{L_0(x)}{C_0}\right) = C_0 \left(1 - \exp\left(-\int_x^\infty \frac{\varepsilon(u)}{u} du\right)\right).$$

The integral in the last formula tends to zero as the tail of a convergent integral. Then, due to the fact that  $1 - e^{-u} \sim u$  as  $u \rightarrow 0$ , we obtain the following equalities:

$$C_0 - L_0(x) = \mathcal{O}^*\left(\int_x^\infty \frac{1}{u^{1+\sigma}} du\right) = \mathcal{O}^*(1/x^\sigma).$$

Combining the last relation with Formula (6) gives

$$\begin{aligned} L(x) &= \exp\{C_\eta + \mathcal{O}^*(1/x^\sigma)\} \left( C_0 + \mathcal{O}^*(1/x^\sigma) \right) \\ &= C_0 \exp\{C_\eta\} \left( 1 + \mathcal{O}^*(1/x^\sigma) \right). \end{aligned}$$

From here, denoting  $C_L := C_0 \exp\{C_\eta\}$ , we get to (5).

The second part of the theorem follows from Formula (5). Indeed,

$$\frac{L(\lambda x)}{L(x)} = \frac{C_L + \mathcal{O}^*(1/x^\sigma)}{C_L + \mathcal{O}^*(1/x^\sigma)} = 1 + \mathcal{O}^*(1/x^\sigma),$$

which stands for  $L(x) \in \mathcal{SV}_\infty(\omega)$  with a remainder  $\omega_\lambda(x) = \mathcal{O}^*(1/x^\sigma)$ .

The theorem is proved completely.  $\square$

**Remark 1.** The assertion of Theorem 3 entails the following conclusion. If the function  $r(x)$  has the order of decreasing  $\varepsilon(x) = \mathcal{O}(L(x)/x^\sigma)$  for some  $\sigma \in \mathbb{R}_+$ , then the class of functions with remainders determined by the condition SR1 coincides with the class of functions with the condition SR2. In this case, the function  $x^\sigma \omega_\lambda(x)$  has a finite limit in an explicit form as  $x \rightarrow \infty$ .

We now turn to the consideration of functions from the class  $\mathcal{R}_\infty^\rho$  with the remainder. The following theorem describes their important asymptotic character.

**Theorem 4.** Let  $L(x) \in \mathcal{SV}_\infty(\omega)$  be a function with a remainder term  $\omega_\lambda(x) = \mathcal{O}^*(L(x)/x^\sigma)$  for some  $\sigma \in \mathbb{R}_+$ . If it is differentiable, then for functions  $R(x) = x^\rho L(x)$  from the class  $\mathcal{R}_\infty^\rho$ , the following asymptotic relation holds:

$$\frac{xR'(x)}{R(x)} = \rho + \mathcal{O}^*(1/x^\sigma). \tag{7}$$

**Proof.** Due to the differentiability of the function  $L(x)$ , we have

$$R'(x) = \rho \frac{R(x)}{x} + x^\rho L'(x) \tag{8}$$

In view of (3), we write the integral representation

$$R(x) = x^\rho \exp\left\{ \eta(x) + \int_b^\infty \frac{\varepsilon(u)}{u} du \right\}, \tag{9}$$

where  $\eta(x) = C_\eta + \mathcal{O}^*(1/x^\sigma)$  and  $\varepsilon(x) = \mathcal{O}^*(1/x^\sigma)$ , which is proved in Theorem 3. At the same time, from the same representation (3) we easily obtain the following relation:

$$L'(x) = L(x) \left[ \mathcal{O}^*(1/x^{\sigma+1}) + \frac{\varepsilon(x)}{x} \right] = \frac{L(x)}{x} \mathcal{O}^*(1/x^\sigma). \tag{10}$$

Now, combining the relations (8)–(10), we obtain

$$R'(x) = \frac{R(x)}{x} \left[ \rho + \mathcal{O}^*(1/x^\sigma) \right],$$

which is equivalent to relation (7).  $\square$

The following statement is an analog of Karamata’s theorem for functions from the class  $L(x) \in \mathcal{SV}_\infty(\omega)$  on the asymptotes of the integral, in which the estimate of the next term of the integral’s value is established.

**Theorem 5.** Let  $L(x) \in \mathcal{SV}_\infty(\omega)$  be a locally bounded function on the set  $[a, \infty) \subset \mathbb{R}_+$  with a remainder  $\omega_\lambda(x) = \mathcal{O}^*(L(x)/x^\sigma)$  for some  $\sigma \in \mathbb{R}_+$ . Then, for all  $\alpha \in (-1, \infty)$  the following relation holds:

$$\int_a^x t^\alpha L(t) dt = \frac{1}{\alpha + 1} x^{\alpha+1} L(x) \left( 1 + \mathcal{O}\left(1/x^\beta\right) \right), \tag{11}$$

where  $\beta = \min(\sigma, \alpha + 1)$ .

**Proof.** Denote

$$I(x) := \int_a^x t^\alpha L(t) dt.$$

Replacing the variable  $t = ux$ , we have

$$\begin{aligned} I(x) &= x^{\alpha+1} \int_{a/x}^1 u^\alpha L(ux) du \\ &= x^{\alpha+1} L(x) \int_{a/x}^1 u^\alpha du - x^{\alpha+1} L(x) \int_{a/x}^1 \lambda^\alpha \left[ 1 - \frac{L(\lambda x)}{L(x)} \right] d\lambda. \end{aligned}$$

Obviously,  $\int_{a/x}^1 u^\alpha du = (x^{\alpha+1} - a^{\alpha+1}) / (1 + \alpha)x^{\alpha+1}$ . To estimate the second integral, we can easily verify that the function  $L(x)$  satisfies the assertions of Theorem 3 with the modulus of the remainder  $|\omega_\lambda(x)| = |1 - L(\lambda x)/L(x)|$  and

$$\frac{L(\lambda x)}{L(x)} - 1 = \mathcal{O}^*(1/x^\sigma)$$

for all  $\lambda \in (0, 1]$ . Then, we obtain the following equality:

$$I(x) = \frac{1}{\alpha + 1} x^{\alpha+1} L(x) (1 + \tau(x)), \tag{12}$$

where

$$\tau(x) = \mathcal{O}\left(\frac{1}{x^{\alpha+1}}\right) + \mathcal{O}\left(\frac{1}{x^\sigma}\right).$$

To complete the proof of the theorem, it suffices to choose the degree of decrease of the tail of  $\tau(x)$  in the order of  $\mathcal{O}(1/x^\beta)$ , where  $\beta = \min(\sigma, \alpha + 1)$ . Then, the representation (12) can be written as (11).

The theorem is proved.  $\square$

Now consider the integral  $\int_a^x t^\alpha L(t) dt$  for  $\alpha \in (-\infty, -1)$ . Take some normalized function  $L_0(x) \in \mathcal{SV}_\infty(\omega)$  defined on  $[a, \infty) \subset \mathbb{R}_+$ . It is known that such functions admit the integral representation  $L_0(x) = \exp \int_b^x [\varepsilon(u)/u] du$  for some  $b \in \mathbb{R}_+$ . We introduce the integral

$$J(x) := \int_a^x t^\alpha L_0(t) dt$$

and using the formula of integration by parts, we write it in the following form:

$$J(x) = \frac{1}{\alpha + 1} L_0(t) t^{\alpha+1} \Big|_a^x + \frac{1}{\alpha + 1} \int_a^x t^{\alpha+1} dL_0(t). \tag{13}$$

It is obvious that  $d[L_0(t)] = L_0(t)\varepsilon(t)dt/t$ . Taking this equality into account, from the relation (13), we write out

$$\int_a^x t^\alpha L_0(t) \left[ 1 - \frac{1}{\alpha + 1} \varepsilon(t) \right] dt = \frac{1}{\alpha + 1} L_0(t) t^{\alpha+1} \Big|_a^x. \tag{14}$$

Now, define the function

$$L(x) := L_0(x) \left[ 1 - \frac{1}{\alpha + 1} \varepsilon(x) \right].$$

Then,

$$\frac{L_0(x)}{L(x)} = 1 + \mathcal{O}^*(\varepsilon(x)) \tag{15}$$

and  $L_0(x) \in \mathcal{SV}_\infty$ . Thus, relations (14) and (15) lead us to the following conclusion: for every normalized function  $L_0(x)$ , there exists a function  $L(x)$  from the class  $\mathcal{SV}_\infty$  with property (15) such that the following relation holds:

$$\int_a^x t^\alpha L(t) dt = \frac{1}{|\alpha + 1|} \left[ \frac{1}{a^{|\alpha+1|}} L_0(a) - \frac{1}{x^{|\alpha+1|}} L_0(x) \right]. \tag{16}$$

On the other hand, as Theorem 3 states, for any function  $\ell(x) \in \mathcal{SV}_\infty$  there exists a function  $v(x)$  such that  $|v(x)| \rightarrow v \in \mathbb{R}_+$  as  $x \rightarrow \infty$  and the following integral representation is allowed:

$$\ell(x) = v(x) \exp \int_b^x \frac{\varepsilon(u)}{u} du.$$

Therefore, there exists a normalized function  $\ell_0(x) \in \mathcal{SV}_\infty$  such that  $\ell(x)/\ell_0(x) \rightarrow v$  as  $x \rightarrow \infty$ . Therefore, Formula (16) is true for all functions in the class of  $\mathcal{SV}_\infty$ .

We have, thus, proved the following theorem.

**Theorem 6.** For any function  $L(x) \in \mathcal{SV}_\infty$  defined on the set  $[a, \infty) \subset \mathbb{R}_+$ , the integral

$$I(x) := \int_a^x t^\alpha L(t) dt$$

converges for all  $\alpha \in (-\infty, -1)$ , and there exists a normalized function  $L_0(x) \in \mathcal{SV}_\infty$  with the property (15) such that the value of the integral  $I(x)$  is calculated by the Formula (16).

The next statement follows from Theorem 6.

**Corollary 1.** Let the function  $L(x) \in \mathcal{SV}_\infty(\omega)$  from Theorem 6 satisfy the condition  $\mathcal{SR}1$  with a remainder  $\omega_\lambda(x) = \mathcal{O}(L(x)/x^\sigma)$  for  $\sigma \in \mathbb{R}_+$ . Then, for the tail of the integral  $\int_a^x t^\alpha L(t) dt$ , the following asymptotic estimation is true:

$$\int_x^\infty t^\alpha L(t) dt = \frac{1}{|\alpha + 1|} \frac{1}{x^{|\alpha+1|}} L(x) \left( 1 + \mathcal{O}^*(1/x^\sigma) \right).$$

The proof is based on the proof scheme of Theorem 5.

### 3. On Applications of SV-Functions

In conclusion, we state one important application of SV-functions in the theory of branching random systems.

Let  $\mathbb{N}$  be the set of natural numbers and  $\mathbb{N}_0 = \{0\} \cup \mathbb{N}$ . Denote by  $Z(t)$  the population size at time  $t \in \mathcal{T} := [0, +\infty)$  in a homogeneous continuous-time Markov branching system with branching law intensity  $\{a_j, j \in \mathbb{N}_0\}$ . The family of random variables  $\{Z(t), t \in \mathcal{T}\}$  forms a homogeneously continuous-time decomposable Markov chain with state space  $\mathcal{S}_0 = \{0\} \cup \mathcal{S}$ , where  $\{0\}$  is the only absorbing state and  $\mathcal{S} \subset \mathbb{N}$  is the class of all communicating states. We introduce the conditional probability  $\mathbb{P}_i\{*\} := \mathbb{P}\{*|Z(0) = i\}$  given that at the initial moment there are  $i \in \mathcal{S}$  particles in the system. The transition probabilities of this chain  $p_{ij}(t) = \mathbb{P}_i\{Z(t) = j\}$  for any  $i, j \in \mathcal{S}_0$  are determined by the  $i$ -fold convolution of the probability  $p_j(t) := p_{1j}(t)$ . The probability generating function (GF)

$$F(t; s) := \sum_{j \in \mathcal{S}_0} p_j(t) s^j$$

admits the following local representation:

$$F(\tau; s) = s + f(s) \cdot \tau + o(\tau) \quad \text{as } \tau \downarrow 0$$

for all  $s \in [0, 1)$ , where

$$f(s) = \sum_{j \in \mathcal{S}_0} a_j s^j.$$

The parameter

$$m := \sum_{j \in \mathcal{S}} j a_j = f'(1-)$$

denotes the average intensity of the law of transformation of one particle, which essentially regulates the asymptotic behavior of the trajectories of the system  $\{Z(t)\}$ . The probability of degeneration  $q$  of a Markov branching system, as the smallest positive root of the equation  $f(s) = 0$  on the set  $[0, 1]$ , is equal to 1 when  $m \leq 0$  and less than 1 when  $m > 0$ . In this regard, three types of the system  $\{Z(t)\}$  are distinguished in accordance with its asymptotic behavior. It is called subcritical, critical, and supercritical if  $m < 0$ ,  $m = 0$ , and  $m > 0$ , respectively. A detailed description of Markov branching random systems and classical results on the structural and asymptotic properties of these systems are contained in the monographs [2,7–11]. Special tasks are discussed in papers [12–16]

In the papers [6,17–22], deeper properties of branching system models are studied using elements of Karamata SV-functions. The main advantage of using SV-functions is that in this context, one can bypass the conditions of the finiteness of the factorial moments of an integer order of the particle transformation intensity laws. For example, if in the critical case, we assume the condition that the infinitesimal GF  $f(s)$  admits for all  $s \in [0, 1)$  the representation

$$f(s) = (1 - s)^{1+\nu} \mathcal{M}\left(\frac{1}{1 - s}\right) \tag{17}$$

for  $\nu \in [0, 1)$ , where  $\mathcal{M}(\ast) \in \mathcal{SV}_\infty$ , then  $f''(1-) = \infty$ . The arguments in the works [5,17,23,24], based on the condition (17), contributed to the refinement of several classical theorems established under the condition of a finite dispersion of the law of intensities of particle transformation in critical branching systems.

In what follows, we consider a non-critical branching system with the probability of degeneration  $q > 0$  and in the case when  $p_0(t) + p_1(t) > 0$  for any  $t \in \mathcal{T}$ , which corresponds to the case of branching random systems of the Schröder type [24]. In accordance with the class  $\mathcal{SV}_\infty$ , we denote by  $\mathcal{SV}_0$  the class of SV-functions at zero.

The following result is established in the paper [6].

**Lemma 1.** *Let  $\beta := \exp\{f'(q)\}$  and  $R_q(t; s) := 1 - F(t; qs)/q$ . There exist functions  $\ell(\ast)$  and  $\mathcal{L}(\ast)$  from the class  $\mathcal{SV}_0$  such that for all  $s \in [0, 1)$  the following representation holds:*

$$R_q(t; s) = (1 - s)L_\beta(t; 1 - s)\beta^t \tag{18}$$

for any  $t \in \mathcal{T}$ , where

$$L_\beta(t; x) = \ell(x) \cdot \mathcal{L}(x\ell(x)\beta^t)$$

for all  $x \in (0, 1]$ . At that  $\ell(1) = 1$  and  $\lim_{x \downarrow 0} \ell(x)\mathcal{L}(x) = 1$ .

Now, we demonstrate some important consequences of this lemma. For  $s = 0$ , from the relation (18), we find the probability of the population size being positive in the system  $\{Z(t), t \in \mathcal{T}\}$  at the final moment of time:

$$\mathbb{P}\{t < \mathcal{H} < \infty\} = qR_q(t; 0) = qL_\beta(t; 1)\beta^t = q\mathcal{L}(\beta^t)\beta^t,$$



Here, the variable  $\mathcal{H} := \inf\{t \in \mathcal{T} : Z(t) = 0\}$  denotes the moment of degeneration of the branching system initiated by the single founder. If we additionally require that condition

$$\sum_{j \in \mathcal{S}} a_j j \ln j < \infty, \tag{19}$$

is satisfied, then  $\mathcal{L}(u) \rightarrow 1/\mu$  as  $u \downarrow 0$ , where  $\mu$  is the mathematical expectation of the invariant distribution of the non-critical Markov branching system. Therefore, we have the following deeper information for the desired probability:

$$\beta^{-t} \cdot \mathbb{P}\{t < \mathcal{H} < \infty\} \rightarrow \frac{q}{\mu} \quad \text{as } t \rightarrow \infty.$$

Since in the subcritical case  $q = 1$  and  $m \equiv \ln \beta$ , then  $\mathbb{P}\{\mathcal{H} < \infty\} = 1$  and the probability of survival of the system at the current moment  $\mathbb{P}\{Z(t) > 0\} = R_1(t; 0)$ . Therefore, it is easy to calculate that the mathematical expectation of the number of particles on positive trajectories of the subcritical system varies slowly at zero, i.e.,

$$\mathbb{E}[Z(t) \mid Z(t) > 0] = \frac{\mathbb{E}Z(t)}{\mathbb{P}\{Z(t) > 0\}} = \mathcal{L}_m(e^{mt}),$$

where  $\mathcal{L}_m(*)\mathcal{L}(*) = 1$ . If we again require the condition (19) to be satisfied, we obtain that  $\mathcal{L}_m(u) \rightarrow \mu$  as  $u \downarrow 0$ . Then, the conditional mathematical expectation  $\mathbb{E}[Z(t) \mid Z(t) > 0]$  slowly stabilizes with increasing  $t$ , approaching  $\mu$ . For a known value of  $\mu$ , one can find an expression for the famous Kolmogorov constant  $\mathcal{K}$ , from the theory of subcritical Markov branching systems, in the form  $\mathcal{K} = 1/\mu$ . Note that the explicit form of the Kolmogorov constant in the case of branching systems with discrete time and finite dispersion of the particle transformation law was calculated for the first time in a recent work [25].

According to the definition, we can write that for any  $\lambda \in \mathbb{R}_+$

$$\frac{\mathcal{L}_m(\lambda u)}{\mathcal{L}_m(u)} = 1 + \omega_\lambda(u),$$

where  $\omega_\lambda(u)$  is infinitesimal at  $u \downarrow 0$ . Then, according to Theorem 3,

$$\mathcal{L}_m(u) = \mu + \rho(u),$$

where  $\rho(u) \rightarrow 0$  as  $u \downarrow 0$ . Thus, it is not difficult to establish an analog of Theorem 3 for the class of MM functions at zero. Consequently, by the definition of the class of functions  $\mathcal{SV}_\infty$ , the function  $\mathcal{L}_m(u)$  belongs to the class  $\mathcal{SV}_0$  with a remainder term  $\omega_\lambda(u)$  that satisfies one of the following conditions as  $u \downarrow 0$ :

- (SR<sub>0</sub>1)  $\omega_\lambda(u) = \mathcal{O}(\rho(u))$ ;
- (SR<sub>0</sub>2)  $\omega_\lambda(u) \sim \text{const}(\lambda)\rho(u)$ ;
- (SR<sub>0</sub>3)  $\omega_\lambda(u) = o(\rho(u))$ .

In this case, we obtain the following asymptote for the conditional mathematical expectation of the population size:

$$\mathbb{E}[Z(t) \mid Z(t) > 0] = \frac{1}{\mathcal{K}} + \tau(t),$$

where  $\tau(t) \rightarrow 0$  as  $t \rightarrow \infty$ . At the same time, we note that if we additionally assume the finiteness of the second infinitesimal moment  $f''(1-)$ , then following the arguments from the paper [25], we can find an explicit expression for the constant  $\mathcal{K}$  and determine the rate of decrease of the remainder  $\tau(t)$ .

The above Lemma 1 and its corollaries indicate that the property of regular variation is implicitly present in the asymptotic structure of a non-critical Markov branching system. Similar situations are observed in many other mathematical structures; see [3].

#### 4. Conclusions

The main characteristic properties of slowly varying functions in the sense of Karamata are presented in the main theorems: the uniform convergence theorem and the integral representation theorem; see [1] (Ch.1). In this paper, we study the class of slowly varying functions with a remainder. The concept of slow varying with a remainder reveals deeper properties of Karamata functions. The main reason for the appearance of this paper is the need for the concept of slow varying with a remainder in the theory of branching random systems. Namely, in the work of Imomov and Tukhtaev [5], Karamata functions with a remainder were used for the first time, which made it possible to estimate the tails of asymptotic expansions in limit theorems of the theory of discrete branching systems with immigration. In connection with the above, it is of great importance to establish analogs of a number of the main theorems of the theory of Karamata functions for the case of slow varying with a remainder.

Overall, this paper proves six theorems, which are new in the sense that the results obtained in them improve classical results, giving explicit estimates of tails in asymptotic formulas. Promisingly, these theorems will certainly contribute to the establishing of new theorems and improving existing theorems in areas of research where methods of asymptotic analysis are used in combination with the slow variation concept.

**Author Contributions:** Conceptualization, A.A.I.; Methodology, A.A.I. and E.E.T.; Validation, A.A.I., E.E.T. and J.S.; Formal Analysis, A.A.I., E.E.T. and J.S.; Investigation, A.A.I. and E.E.T.; Resources, A.A.I. and E.E.T.; Writing-Original Draft Preparation, A.A.I.; Writing-Review and Editing, A.A.I.; Visualization, A.A.I., E.E.T. and J.S.; Supervision, A.A.I.; Project Administration, A.A.I. and J.S. All authors have read and agreed to the published version of the manuscript.

**Funding:** This research received no external funding.

**Data Availability Statement:** All data generated or analyzed during this study are included in this published article.

**Acknowledgments:** The authors thank the reviewers for their careful reading of the manuscript and for their kind comments and useful suggestions that contributed to the improvement of the paper.

**Conflicts of Interest:** The authors declare no conflict of interest.

#### References

1. Seneta, E. *Regularly Varying Functions*; Translated Russian, Nauka: Moscow, Russia, 1985; Springer: Berlin/Heidelberg, Germany, 1976.
2. Asmussen, S.; Hering, H. *Branching Processes*; Birkhäuser: Boston, MA, USA, 1983.
3. Bingham, N.H.; Goldie, C.M.; Teugels, J.L. *Regular Variation*; Cambridge University Press: Cambridge, UK, 1987.
4. Zolotarev, V.M. More Exact Statements of Several Theorems in the Theory of Branching Processes. *Theory Probab. Appl.* **1957**, *2*, 245–253. [[CrossRef](#)]
5. Imomov, A.A.; Tukhtaev, E.E. On asymptotic structure of critical Galton-Watson branching processes allowing immigration with infinite variance. *Stoch. Model.* **2023**, *39*, 118–140. [[CrossRef](#)]
6. Imomov, A.A.; Meyliev, A.K. On the asymptotic structure of non-critical Markov stochastic branching processes with continuous time. *Vestn. Tomsk. Gos. Univ. Mat. Mekh.* **2021**, *69*, 22–36.
7. Athreya, K.B.; Ney, P.E. *Branching Processes*; Springer: New York, NY, USA, 1972.
8. Haccou, P.; Jagers, P.; Vatutin, V. *Branching Processes: Variation, Growth, and Extinction of Populations*; Cambridge University Press: Cambridge, UK, 2007.
9. Harris, T.E. *Theory of Branching Stochastic Process*; Springer: Berlin/Heidelberg, Germany, 1963.
10. Jagers, P. *Branching Processes with Biological Applications*; John Wiley & Sons Inc.: Hoboken, NJ, USA; Pitman Press: London, UK, 1975.
11. Sevastyanov, B.A. *Branching Process*; Nauka: Moscow, Russia, 1971. (In Russian)
12. Chen, X.; He, H. Lower deviation and moderate deviation probabilities for maximum of a branching random walk. *arXiv* **2018**, arXiv:1807.08263. [[CrossRef](#)]
13. Nagaev, S.V.; Wachtel, V. The Critical Galton-Watson Process without Further Power Moments. *J. Appl. Prob.* **2007**, *44*, 753–769. [[CrossRef](#)]
14. Pakes, A.G. Revisiting conditional limit theorems for the mortal simple branching process. *Bernoulli* **1999**, *5*, 969–998. [[CrossRef](#)]

15. Pakes, A.G. Limit theorems for the simple branching process allowing immigration, I. The case of finite offspring mean. *Adv. Appl. Prob.* **1979**, *11*, 31–62. [[CrossRef](#)]
16. Pakes, A.G. Some new limit theorems for critical branching processes allowing immigration. *Stoch. Proc. Appl.* **1975**, *3*, 175–185. [[CrossRef](#)]
17. Formanov, S.K.; Azimov, Z.B. Markov branching processes with regularly varying generating function and immigration of a special form. *Theory Prob. Math. Stat.* **2002**, *65*, 181–188.
18. Hudson, I.L.; Seneta, E. A note on simple branching processes with infinite mean. *J. Appl. Prob.* **1977**, *14*, 836–842. [[CrossRef](#)]
19. Li, J.; Chen, A.; Pakes, A.G. Asymptotic properties of the Markov Branching Process with Immigration. *J. Theor. Probab.* **2012**, *25*, 122–143. [[CrossRef](#)]
20. Seneta, E. Regularly Varying Functions in the Theory of Simple Branching Processes. *Adv. Appl. Prob.* **1974**, *6*, 408–420. [[CrossRef](#)]
21. Schuh, H.J.; Barbour, A.D. On the Asymptotic Behaviour of Branching Processes with Infinite Mean. *Adv. Appl. Prob.* **1977**, *9*, 681–723. [[CrossRef](#)]
22. Slack, R.S. A branching process with mean one and possible infinite variance. *Wahrscheinlichkeitstheor. Verw. Geb.* **1968**, *9*, 139–145. [[CrossRef](#)]
23. Imomov, A.A. On long-term behavior of continuous-time Markov Branching Processes allowing immigration. *J. Sib. Fed. Univ. Math. Phys.* **2014**, *7*, 443–454. [[CrossRef](#)]
24. Pakes, A.G. Critical Markov branching process limit theorems allowing infinite variance. *Adv. Appl. Prob.* **2010**, *42*, 460–488. [[CrossRef](#)]
25. Imomov, A.A.; Murtazaev, M. On the Kolmogorov constant explicit form in the theory of Discrete-time Stochastic Branching Systems. *J. Appl. Prob.* **2024**, *61*, 927–941. [[CrossRef](#)]

**Disclaimer/Publisher’s Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.