

Article

# Extension of Extragradient Techniques for Variational Inequalities

**Yonghong Yao** <sup>1,2,3,\*</sup>, **Ke Wang** <sup>1</sup>, **Xiaowei Qin** <sup>1</sup> and **Li-Jun Zhu** <sup>2,3</sup>

<sup>1</sup> School of Mathematical Sciences, Tianjin Polytechnic University, Tianjin 300387, China; wangke@aliyun.com (K.W.); qinxiaowei@aliyun.com (X.Q.)

<sup>2</sup> The Key Laboratory of Intelligent Information and Data Processing of NingXia Province, North Minzu University, Yinchuan 750021, China; zhulijun1995@sohu.com

<sup>3</sup> Health Big Data Research Institute of North Minzu University, Yinchuan 750021, China

\* Correspondence: yaoyonghong@aliyun.com or yaoyonghong@tjpu.edu.cn

Received: 4 December 2018; Accepted: 18 January 2019; Published: 22 January 2019



**Abstract:** An extragradient type method for finding the common solutions of two variational inequalities has been proposed. The convergence result of the algorithm is given under mild conditions on the algorithm parameters.

**Keywords:** variational inequality; extragradient-type method; inverse-strongly-monotone; projection; strong convergence

**MSC:** 47H05; 47J05; 47J25.

## 1. Introduction

Let  $H$  be a real Hilbert space equipped with inner product  $\langle \cdot, \cdot \rangle$  and norm  $\|\cdot\|$ . Let  $\emptyset \neq C \subset H$  be a closed and convex set. Let  $A: C \rightarrow H$  be a mapping. Recall that the variational inequality (VI) seeks an element  $x^* \in C$  such that

$$\langle Az^*, z - z^* \rangle \geq 0, \quad \forall z \in C. \quad (1)$$

The solution set of (1) is denoted by  $VI(C, A)$ .

The problem (1) introduced and studied by Stampacchia [1] is being applied as a useful tool and model to refine a multitude of problems. A large number of methods for solving VI (1) are projection methods that implement projections onto the feasible set of the VI (1), or onto another set in order to achieve a solution. Several iterative methods for solving the VI (1) have been proposed. See, e.g., [2–30]. A basic one is the natural extension of the gradient projection algorithm for solving the optimization problem  $\min_{x \in C} f(x)$ . For  $x_0 \in C$ , calculate iteratively the sequence  $\{x_n\}$  through

$$x_{n+1} = proj_C[x_n - \alpha_n \nabla f(x_n)], \quad n \geq 0, \quad (2)$$

where  $proj$  is the metric projection and  $\alpha_n > 0$  is the step-size.

Korpelevich [31] introduced an iterative method for solving the VI (1), known as the extragradient method ([7]). In Korpelevich's method, two projections are used for computing the next iteration. For the current iteration  $x_n$ , compute

$$\begin{cases} y_n = proj_C[x_n - \xi Ax_n], \\ x_{n+1} = proj_C[x_n - \xi Ay_n], \end{cases} \quad n \geq 0, \quad (3)$$

where  $\xi > 0$  is a fixed number.

Korpelevich's method has received so much attention by a range of scholars, who improved it in several ways; see, e.g., [32–36]. Now, we know that Korpelevich's method (3) can only achieve weak convergence in a large dimensional spaces ([37,38]). In order to reach strong convergence, Korpelevich's method was adapted by many mathematicians. For example, in [32], it is shown that several extragradient-type methods converge strongly to an element in  $VI(C, A)$ .

Very recently, Censor, Gibali and Reich [39] presented an alternating method for finding common solutions of two variational inequalities. In [40], Zaslavski studied an extragradient method for finding common solutions of a finite family of variational inequalities.

Inspired by the work given above, in this article, we present an extragradient type method for finding the common solutions of two variational inequalities. We prove the strong convergence of the proposed method under the mild assumptions on the parameters.

## 2. Preliminaries

Let  $H$  be a real Hilbert space. Let  $C \subset H$  be a nonempty, closed and convex set.

**Definition 1.** An operator  $A: C \rightarrow H$  is called Lipschitz if

$$\|Au^\dagger - Av^\dagger\| \leq L\|u^\dagger - v^\dagger\|, \forall u^\dagger, v^\dagger \in C,$$

where  $L > 0$  is a constant.

If  $L = 1$ , we call  $A$  is nonexpansive.

**Definition 2.** An operator  $A: C \rightarrow H$  is called inverse strongly monotone if

$$\alpha\|Au^\dagger - Av^\dagger\|^2 \leq \langle Au^\dagger - Av^\dagger, u^\dagger - v^\dagger \rangle, \forall u^\dagger, v^\dagger \in C,$$

where  $\alpha \geq 0$  is a constant.

In this case, we call  $A$  is  $\alpha$ -inverse-strongly-monotone.

**Proposition 1** ([41]). If  $C$  is a bounded closed convex subset of a real Hilbert space  $H$  and  $A: C \rightarrow H$  is an inverse strongly monotone operator, then  $VI(C, A) \neq \emptyset$ .

For fixed  $z \in H$ , there exists a unique  $z^\dagger \in C$  satisfying

$$\|z - z^\dagger\| = \inf\{\|z - \tilde{z}\| : \tilde{z} \in C\}.$$

We denote  $z^\dagger$  by  $proj_C z$ . The following inequality is an important property of projection  $proj_C$ : for given  $x \in H$ ,

$$\langle x - proj_C x, y - proj_C x \rangle \leq 0, \forall y \in C, \quad (4)$$

which is equivalent to

$$\langle x - y, proj_C x - proj_C y \rangle \geq \|proj_C x - proj_C y\|^2, \forall x, y \in H.$$

It follows that  $proj_C$  is nonexpansive. We also know that  $2proj_C - I$  is nonexpansive.

**Lemma 1 ([41]).** If  $\emptyset \neq C$  is a closed convex subset of a real Hilbert space  $H$  and  $A : C \rightarrow H$  is an  $\alpha$ -inverse strongly monotone operator, then

$$\|(I - \mu A)u - (I - \mu A)v\|^2 \leq \|u - v\|^2 + \mu(\mu - 2\alpha)\|Au - Av\|^2, \forall u, v \in C.$$

Especially,  $I - \mu A$  is nonexpansive provided  $0 \leq \mu \leq 2\alpha$ .

**Lemma 2 ([42]).** Suppose that  $\{u_n\}$  and  $\{v_n\}$  are two bounded sequences in Banach spaces. Let  $\{\lambda_n\} \subset [0, 1]$  be a sequence satisfying  $0 < \liminf_{n \rightarrow \infty} \lambda_n \leq \limsup_{n \rightarrow \infty} \lambda_n < 1$ . Suppose  $u_{n+1} = (1 - \lambda_n)v_n + \lambda_n u_n, \forall n \geq 0$  and  $\limsup_{n \rightarrow \infty} (\|v_{n+1} - v_n\| - \|u_{n+1} - u_n\|) \leq 0$ . Then  $\lim_{n \rightarrow \infty} \|u_n - v_n\| = 0$ .

**Lemma 3 ([43]).** Let  $\{\mu_n\} \subset (0, \infty)$ ,  $\{\gamma_n\} \subset (0, 1)$  and  $\{\delta_n\}$  be three real number sequences. If  $\mu_{n+1} \leq (1 - \gamma_n)\mu_n + \delta_n$  for all  $n \geq 0$  with  $\sum_{n=1}^{\infty} \gamma_n = \infty$  and  $\limsup_{n \rightarrow \infty} \delta_n/\gamma_n \leq 0$  or  $\sum_{n=1}^{\infty} |\delta_n| < \infty$ , then  $\lim_{n \rightarrow \infty} \mu_n = 0$ .

### 3. Main Results

Let  $\emptyset \neq C$  be a convex and closed subset of a real Hilbert space  $H$ . Let the operators  $A, B : C \rightarrow H$  be  $\alpha$ -inverse strongly monotone and  $\beta$ -inverse strongly monotone, respectively. Let  $\{\alpha_n\} \subset [0, 1]$ ,  $\{\beta_n\} \subset [0, 1]$ ,  $\{\lambda_n\} \subset [0, 2\alpha]$  and  $\{\mu_n\} \subset [0, 2\beta]$  be four sequences. In the sequel, assume  $\Omega = VI(C, A) \cap VI(C, B) \neq \emptyset$ .

Motivated by the algorithms presented in [31,39,40], we present the following iterative Algorithm 1 for finding the common solution of two variational inequalities.

---

#### Algorithm 1:

For  $u, x_0 \in C$ . Assume the sequence  $\{x_n\}$  has been constructed. Compute the next iteration  $\{x_{n+1}\}$  by the following manner

$$\begin{cases} y_n = proj_C[x_n - (1 - \alpha_n)\lambda_n Ax_n + \alpha_n(u - x_n)], \\ x_{n+1} = proj_C[x_n - \mu_n \beta_n B y_n + \beta_n(y_n - x_n)], \end{cases} n \geq 0. \quad (5)$$

Suppose that the control parameters  $\{\alpha_n\}$ ,  $\{\beta_n\}$ ,  $\{\lambda_n\}$  and  $\{\mu_n\}$  satisfy the following assumptions:

- (C1):  $\lim_{n \rightarrow \infty} \alpha_n = 0$  and  $\sum_{n=1}^{\infty} \alpha_n = \infty$ ;
  - (C2):  $\liminf_{n \rightarrow \infty} \beta_n > 0$  and  $\lim_{n \rightarrow \infty} (\beta_{n+1} - \beta_n) = 0$ ;
  - (C3):  $\lambda_n \in [a, b] \subset (0, 2\alpha)$  and  $\lim_{n \rightarrow \infty} (\lambda_{n+1} - \lambda_n) = 0$ ;
  - (C4):  $\mu_n \in [c, d] \subset (0, 2\beta)$  and  $\lim_{n \rightarrow \infty} (\mu_{n+1} - \mu_n) = 0$ .
- 

We will divide our main result into several propositions.

**Proposition 2.** The sequence  $\{x_n\}$  generated by (5) is bounded.

**Proof.** Choose any  $x^* \in \Omega$ . Note that  $x^* = proj_C[(I - \delta A)x^*]$  for any  $\delta > 0$ . Hence,

$$x^* = proj_C[\alpha_n x^* + (1 - \alpha_n)(x^* - \lambda_n Ax^*)], \forall n \geq 0. \quad (6)$$

Thus, by (5) and (6), we have

$$\begin{aligned} \|y_n - x^*\| &= \|proj_C[\alpha_n u + (1 - \alpha_n)(I - \lambda_n A)x_n] - proj_C[\alpha_n x^* + (1 - \alpha_n)(I - \lambda_n A)x^*]\| \\ &\leq \|[\alpha_n u + (1 - \alpha_n)(I - \lambda_n A)x_n] - [\alpha_n x^* + (1 - \alpha_n)(I - \lambda_n A)x^*]\| \\ &\leq \|\alpha_n(u - x^*) + (1 - \alpha_n)[(I - \lambda_n A)x_n - (I - \lambda_n A)x^*]\| \\ &\leq \alpha_n\|u - x^*\| + (1 - \alpha_n)\|(I - \lambda_n A)x_n - (I - \lambda_n A)x^*\|. \end{aligned} \quad (7)$$

From Lemma 1, we know that  $I - \lambda_n A$  and  $I - \mu_n B$  are nonexpansive. Thus, from (7), we get

$$\|y_n - x^*\| \leq \alpha_n \|u - x^*\| + (1 - \alpha_n) \|x_n - x^*\|.$$

So,

$$\begin{aligned} \|x_{n+1} - x^*\| &= \|proj_C[(1 - \beta_n)x_n + \beta_n(y_n - \mu_n B y_n)] - proj_C[(1 - \beta_n)x^* + \beta_n(x^* - \mu_n B x^*)]\| \\ &\leq \|(1 - \beta_n)x_n + \beta_n(y_n - \mu_n B y_n)\| - \|(1 - \beta_n)x^* + \beta_n(x^* - \mu_n B x^*)\| \\ &\leq \beta_n \|y_n - \mu_n B y_n\| + (1 - \beta_n) \|x_n - x^*\| \\ &\leq \beta_n \|y_n - x^*\| + (1 - \beta_n) \|x_n - x^*\| \\ &\leq (1 - \beta_n \alpha_n) \|x_n - x^*\| + \beta_n \alpha_n \|u - x^*\|. \end{aligned} \quad (8)$$

It follows that

$$\|x_{n+1} - x^*\| \leq \max\{\|x_0 - x^*\|, \|u - x^*\|\}.$$

Then  $\{x_n\}$  is bounded, and hence the sequences  $\{y_n\}$ ,  $\{Ax_n\}$  and  $\{By_n\}$  are all bounded.  $\square$

**Proposition 3.** *The following two conclusions hold*

$$\lim_{n \rightarrow \infty} \|x_{n+1} - x_n\| = 0 \text{ and } \lim_{n \rightarrow \infty} \|x_n - y_n\| = 0.$$

**Proof.** Let  $S = 2proj_C - I$ . It is clear that  $S$  is nonexpansive. Set  $v_n = (1 - \beta_n)x_n + \beta_n(I - \mu_n B)y_n$  for all  $n \geq 0$ . Then, we can rewrite  $x_{n+1}$  in (5) as

$$\begin{aligned} x_{n+1} &= \frac{1 - \beta_n}{2}x_n + \frac{\beta_n}{2}(I - \mu_n B)y_n + \frac{1}{2}Sv_n \\ &= \frac{1 + \beta_n}{2}z_n + \frac{1 - \beta_n}{2}x_n, \end{aligned}$$

where  $z_n = \frac{\beta_n}{1 + \beta_n}(I - \mu_n B)y_n + \frac{1}{1 + \beta_n}Sv_n$  for all  $n \geq 0$ .

Hence,

$$z_{n+1} - z_n = \frac{1}{1 + \beta_{n+1}}Sv_{n+1} + \frac{\beta_{n+1}}{1 + \beta_{n+1}}(I - \mu_{n+1}B)y_{n+1} - \frac{\beta_n}{1 + \beta_n}(I - \mu_n B)y_n - \frac{1}{1 + \beta_n}Sv_n.$$

Hence,

$$\begin{aligned} \|z_{n+1} - z_n\| &\leq \frac{\beta_{n+1}}{1 + \beta_{n+1}}\|(I - \mu_{n+1}B)y_{n+1} - (I - \mu_n B)y_n\| + \frac{1}{1 + \beta_{n+1}}\|Sv_{n+1} - Sv_n\| \\ &\quad + \left| \frac{\beta_{n+1}}{1 + \beta_{n+1}} - \frac{\beta_n}{1 + \beta_n} \right| \|(I - \mu_n B)y_n\| + \left| \frac{1}{1 + \beta_{n+1}} - \frac{1}{1 + \beta_n} \right| \|Sv_n\| \\ &\leq \frac{\beta_{n+1}}{1 + \beta_{n+1}}\|(I - \mu_{n+1}B)y_{n+1} - (I - \mu_{n+1}B)y_n\| \\ &\quad + \frac{\beta_{n+1}}{1 + \beta_{n+1}}|\mu_{n+1} - \mu_n| \|By_n\| + \left| \frac{\beta_{n+1}}{1 + \beta_{n+1}} - \frac{\beta_n}{1 + \beta_n} \right| \|y_n - \mu_n B y_n\| \\ &\quad + \frac{1}{1 + \beta_{n+1}}\|Sv_{n+1} - Sv_n\| + \left| \frac{1}{1 + \beta_{n+1}} - \frac{1}{1 + \beta_n} \right| \|Sv_n\|. \end{aligned}$$

By using the nonexpansivity of  $I - \mu_n B$  and  $S$  to deduce

$$\begin{aligned}
\|z_{n+1} - z_n\| &\leq \frac{\beta_{n+1}}{1 + \beta_{n+1}} \|y_{n+1} - y_n\| + \left| \frac{\beta_{n+1}}{1 + \beta_{n+1}} - \frac{\beta_n}{1 + \beta_n} \right| \|y_n - \mu_n B y_n\| \\
&\quad + \beta_{n+1} [(I - \mu_{n+1} B) y_{n+1} - (I - \mu_{n+1} B) y_n] + \left| \frac{1}{1 + \beta_n} - \frac{1}{1 + \beta_n} \right| \|S v_n\| \\
&\quad + \frac{\beta_{n+1}}{1 + \beta_{n+1}} |\mu_{n+1} - \mu_n| \|B y_n\| + \frac{1}{1 + \beta_{n+1}} \|(1 - \beta_{n+1})(x_{n+1} - x_n)\| \\
&\quad + (\beta_{n+1} - \beta_n)(y_n - x_n) + (\beta_n \mu_n - \beta_{n+1} \mu_{n+1}) B y_n \| \\
&\leq \frac{|\beta_{n+1} - \beta_n|}{1 + \beta_{n+1}} (\|x_n\| + \|y_n\|) + \frac{1 - \beta_{n+1}}{1 + \beta_{n+1}} \|x_{n+1} - x_n\| + \left| \frac{1}{1 + \beta_n} - \frac{1}{1 + \beta_n} \right| \|S v_n\| \\
&\quad + \frac{2\beta_{n+1}}{1 + \beta_{n+1}} \|y_{n+1} - y_n\| + \left| \frac{\beta_{n+1}}{1 + \beta_{n+1}} - \frac{\beta_n}{1 + \beta_n} \right| \|y_n - \mu_n B y_n\| \\
&\quad + \frac{\beta_{n+1}}{1 + \beta_{n+1}} |\mu_{n+1} - \mu_n| \|B y_n\| + \frac{|\beta_{n+1} \mu_{n+1} - \beta_n \mu_n|}{1 + \beta_{n+1}} \|B y_n\|. \tag{9}
\end{aligned}$$

Next, we estimate  $\|y_{n+1} - y_n\|$ . By (5), we get

$$\begin{aligned}
\|y_{n+1} - y_n\| &= \|proj_C[\alpha_{n+1} u + (1 - \alpha_{n+1})(I - \lambda_{n+1} A)x_{n+1}] - proj_C[\alpha_n u + (1 - \alpha_n)(I - \lambda_n A)x_n]\| \\
&\leq |\alpha_{n+1} - \alpha_n| \|x_n\| + |\lambda_{n+1} - \lambda_n| \|Ax_n\| + \alpha_{n+1} \|u\| + \alpha_n \|u\| + \|x_{n+1} - x_n\| \\
&\quad + |\lambda_{n+1} - \lambda_n| \|Ax_n\| + |\lambda_{n+1} \alpha_{n+1} - \lambda_n \alpha_n| \|Ax_n\|. \tag{10}
\end{aligned}$$

Substituting (10) into (9) to get

$$\begin{aligned}
\|z_{n+1} - z_n\| &\leq \left| \frac{\beta_{n+1}}{1 + \beta_{n+1}} - \frac{\beta_n}{1 + \beta_n} \right| \|y_n - \mu_n B y_n\| + \frac{\beta_{n+1}}{1 + \beta_{n+1}} |\mu_{n+1} - \mu_n| \|B y_n\| \\
&\quad + \frac{|\beta_{n+1} - \beta_n|}{1 + \beta_{n+1}} (\|x_n\| + \|y_n\|) + \frac{|\beta_{n+1} \mu_{n+1} - \beta_n \mu_n|}{1 + \beta_{n+1}} \|B y_n\| + \|x_{n+1} - x_n\| \\
&\quad + \left| \frac{1}{1 + \beta_n} - \frac{1}{1 + \beta_n} \right| \|S v_n\| + 2|\lambda_{n+1} - \lambda_n| \|Ax_n\| + 2\alpha_{n+1} \|u\| + 2\alpha_n \|u\| \\
&\quad + 2|\alpha_{n+1} - \alpha_n| \|x_n\| + 2|\lambda_{n+1} - \lambda_n| \|Ax_n\| + 2|\lambda_{n+1} \alpha_{n+1} - \lambda_n \alpha_n| \|Ax_n\|.
\end{aligned}$$

Since  $\lim_{n \rightarrow \infty} (\beta_{n+1} - \beta_n) = 0$  and  $\lim_{n \rightarrow \infty} (\mu_{n+1} - \mu_n) = 0$ , we derive that  $\lim_{n \rightarrow \infty} \left| \frac{\beta_{n+1}}{1 + \beta_{n+1}} - \frac{\beta_n}{1 + \beta_n} \right| = 0$ ,  $\lim_{n \rightarrow \infty} |\beta_{n+1} \mu_{n+1} - \beta_n \mu_n| = 0$  and  $\lim_{n \rightarrow \infty} \left| \frac{1}{1 + \beta_n} - \frac{1}{1 + \beta_n} \right| = 0$ . At the same time, note that  $\{x_n\}$ ,  $\{Ax_n\}$ ,  $\{y_n\}$  and  $\{B y_n\}$  are bounded. Therefore,

$$\limsup_{n \rightarrow \infty} (\|z_{n+1} - z_n\| - \|x_{n+1} - x_n\|) \leq 0.$$

Applying Lemma 2 to derive

$$\lim_{n \rightarrow \infty} \|z_n - x_n\| = 0.$$

Hence,

$$\lim_{n \rightarrow \infty} \|x_{n+1} - x_n\| = \lim_{n \rightarrow \infty} \frac{1 + \beta_n}{2} \|z_n - x_n\| = 0.$$

By virtue of (7), (8) and Lemma 1, we deduce

$$\begin{aligned}
\|x_{n+1} - x^*\|^2 &\leq (1 - \beta_n)\|x_n - x^*\|^2 + \beta_n\|(y_n - \mu_n B y_n) - (x^* - \mu_n B x^*)\|^2 \\
&\leq (1 - \beta_n)\|x_n - x^*\|^2 + \beta_n\|y_n - x^*\|^2 + \beta_n\mu_n(\mu_n - 2\beta)\|B y_n - B x^*\|^2 \\
&\leq \beta_n[\alpha_n(u - x^*) + (1 - \alpha_n)[(x_n - \lambda_n A x_n) - (x^* - \lambda_n A x^*)]]\|^2 \\
&\quad + (1 - \beta_n)\|x_n - x^*\|^2 + \beta_n\mu_n(\mu_n - 2\beta)\|B y_n - B x^*\|^2 \\
&\leq \beta_n[\alpha_n\|u - x^*\|^2 + (1 - \alpha_n)\|(I - \lambda_n A)x_n - (I - \lambda_n A)x^*\|^2] \\
&\quad + \beta_n\mu_n(\mu_n - 2\beta)\|B y_n - B x^*\|^2 + (1 - \beta_n)\|x_n - x^*\|^2 \\
&\leq (1 - \beta_n)\|x_n - x^*\|^2 + \alpha_n\beta_n\|u - x^*\|^2 + \beta_n\mu_n(\mu_n - 2\beta)\|B y_n - B x^*\|^2 \\
&\quad + (1 - \alpha_n)\beta_n[\|x_n - x^*\|^2 + \lambda_n(\lambda_n - 2\alpha)\|A x_n - A x^*\|^2] \\
&\leq \alpha_n\beta_n\|u - x^*\|^2 + \|x_n - x^*\|^2 + \beta_n\mu_n(\mu_n - 2\beta)\|B y_n - B x^*\|^2 \\
&\quad + (1 - \alpha_n)\beta_n\lambda_n(\lambda_n - 2\alpha)\|A x_n - A x^*\|^2.
\end{aligned}$$

It follows that

$$\begin{aligned}
&(1 - \alpha_n)\beta_n\lambda_n(2\alpha - \lambda_n)\|A x_n - A x^*\|^2 + \beta_n\mu_n(2\beta - \mu_n)\|B y_n - B x^*\|^2 \\
&\leq \alpha_n\beta_n\|u - x^*\|^2 + \|x_n - x^*\|^2 - \|x_{n+1} - x^*\|^2 \\
&\leq \alpha_n\beta_n\|u - x^*\|^2 + (\|x_n - x^*\| + \|x_{n+1} - x^*\|) \times \|x_n - x_{n+1}\|.
\end{aligned}$$

This implies that

$$\lim_{n \rightarrow \infty} \|A x_n - A x^*\| = 0 \text{ and } \lim_{n \rightarrow \infty} \|B y_n - B x^*\| = 0$$

According to (4) and (5), we get

$$\begin{aligned}
\|y_n - x^*\|^2 &= \|proj_C[\alpha_n u + (1 - \alpha_n)(I - \lambda_n A)x_n] - proj_C(I - \lambda_n A)x^*\|^2 \\
&\leq \langle \alpha_n u + (1 - \alpha_n)(I - \lambda_n A)x_n - (I - \lambda_n A)x^*, y_n - x^* \rangle \\
&= \frac{1}{2} \left\{ \|(I - \lambda_n A)x_n - (I - \lambda_n A)x^* + \alpha_n(u - x_n + \lambda_n A x^*)\|^2 + \|y_n - x^*\|^2 \right. \\
&\quad \left. - \|\alpha_n u + (1 - \alpha_n)(I - \lambda_n A)x_n - (I - \lambda_n A)x^* - (y_n - x^*)\|^2 \right\} \\
&\leq \frac{1}{2} \left\{ \|(I - \lambda_n A)x_n - (I - \lambda_n A)x^*\|^2 \right. \\
&\quad + 2\alpha_n\|u - x_n + \lambda_n A x^*\|\|(I - \lambda_n A)x_n - (I - \lambda_n A)x^* + \alpha_n(u - x_n + \lambda_n A x^*)\| \\
&\quad \left. + \|y_n - x^*\|^2 - \|(x_n - y_n) - \lambda_n(A x_n - A x^*) + \alpha_n(u - x_n + \lambda_n A x^*)\|^2 \right\} \\
&\leq \frac{1}{2} \left\{ \|(I - \lambda_n A)x_n - (I - \lambda_n A)x^*\|^2 + \alpha_n M + \|y_n - x^*\|^2 \right. \\
&\quad \left. - \|(x_n - y_n) - \lambda_n(A x_n - A x^*) + \alpha_n(u - x_n + \lambda_n A x^*)\|^2 \right\} \\
&\leq \frac{1}{2} \left\{ \|x_n - x^*\|^2 - \|x_n - y_n\|^2 + \alpha_n M + \|y_n - x^*\|^2 + 2\lambda_n \langle x_n - y_n, A x_n - A x^* \rangle \right. \\
&\quad \left. - 2\alpha_n \langle u - x_n + \lambda_n A x^*, x_n - y_n \rangle - \|\lambda_n(A x_n - A x^*) - \alpha_n(u - x_n + \lambda_n A x^*)\|^2 \right\} \\
&\leq \frac{1}{2} \left\{ \|x_n - x^*\|^2 + \alpha_n M + \|y_n - x^*\|^2 - \|x_n - y_n\|^2 \right. \\
&\quad \left. + 2\alpha_n\|u - x_n + \lambda_n A x^*\|\|x_n - y_n\| + 2\lambda_n\|x_n - y_n\|\|A x_n - A x^*\| \right\},
\end{aligned}$$

where  $M > 0$  is some constant such that

$$\sup_n \{2\|u - x_n + \lambda_n Ax^*\|\|(I - \lambda_n A)x_n - (I - \lambda_n A)x^* + \alpha_n(u - x_n + \lambda_n Ax^*)\|\} \leq M.$$

Thus

$$\begin{aligned} \|y_n - x^*\|^2 &\leq \|x_n - x^*\|^2 + \alpha_n M - \|x_n - y_n\|^2 + 2\lambda_n\|x_n - y_n\|\|Ax_n - Ax^*\| \\ &\quad + 2\alpha_n\|u - x_n + \lambda_n Ax^*\|\|x_n - y_n\|, \end{aligned}$$

and it follows that

$$\begin{aligned} \|x_{n+1} - x^*\|^2 &\leq \|x_n - x^*\|^2 + \alpha_n M - \beta_n\|x_n - y_n\|^2 + 2\lambda_n\|x_n - y_n\|\|Ax_n - Ax^*\| \\ &\quad + 2\alpha_n\|u - x_n + \lambda_n Ax^*\|\|x_n - y_n\|. \end{aligned}$$

Therefore,

$$\begin{aligned} \beta_n\|x_n - y_n\|^2 &\leq 2\lambda_n\|x_n - y_n\|\|Ax_n - Ax^*\| + (\|x_n - x^*\| + \|x_{n+1} - x^*\|)\|x_{n+1} - x_n\| \\ &\quad + \alpha_n M + 2\alpha_n\|u - x_n + \lambda_n Ax^*\|\|x_n - y_n\|. \end{aligned}$$

Since  $\lim_{n \rightarrow \infty} \alpha_n = 0$ ,  $\lim_{n \rightarrow \infty} \|x_n - x_{n+1}\| = 0$  and  $\lim_{n \rightarrow \infty} \|Ax_n - Ax^*\| = 0$ , we derive

$$\lim_{n \rightarrow \infty} \|x_n - y_n\| = 0.$$

This concludes the proof.  $\square$

**Proposition 4.**  $\limsup_{n \rightarrow \infty} \langle \tilde{x} - u, \tilde{x} - y_n \rangle \leq 0$ , where  $\tilde{x} = P_\Omega u$ .

**Proof.** Let  $\{y_{n_i}\}$  be a subsequence of  $\{y_n\}$  satisfying

$$\limsup_{n \rightarrow \infty} \langle \tilde{x} - u, \tilde{x} - y_n \rangle = \lim_{i \rightarrow \infty} \langle \tilde{x} - u, \tilde{x} - y_{n_i} \rangle.$$

By the boundedness of  $\{y_{n_i}\}$ , we can choose a subsequence  $\{y_{n_{ij}}\}$  of  $\{y_{n_i}\}$  such that  $y_{n_{ij}} \rightharpoonup z$ .

Next, we demonstrate that  $z \in \Omega$ . First, we prove that  $z \in \text{VI}(C, A)$ . Let  $N_C v$  be the normal cone of  $C$  at  $v \in C$ ; i.e.,  $N_C v = \{w \in H : 0 \leq \langle v - u, w \rangle, \forall u \in C\}$ . Define a mapping  $T$  by the formula

$$Tv = \begin{cases} Av + N_C v, & v \in C, \\ \emptyset, & v \notin C. \end{cases}$$

Let  $(v, w) \in G(T)$ . Since  $y_n \in C$  and  $w - Av \in N_C v$ , we deduce  $0 \leq \langle v - y_n, w - Av \rangle$ . According to  $y_n = \text{proj}_C[\alpha_n u + (1 - \alpha_n)(I - \lambda_n A)x_n]$ , we obtain

$$0 \leq \langle v - y_n, y_n - \alpha_n u - (1 - \alpha_n)(x_n - \lambda_n Ax_n) \rangle,$$

that is,

$$0 \leq \langle v - y_n, \frac{y_n - x_n}{\lambda_n} + Ax_n - \frac{\alpha_n}{\lambda_n}(u - x_n + \lambda_n Ax_n) \rangle.$$

Thus,

$$\begin{aligned}
\langle v - y_{n_i}, w \rangle &\geq \langle v - y_{n_i}, Av \rangle \\
&\geq \langle v - y_{n_i}, Av \rangle - \langle v - y_{n_i}, \frac{y_{n_i} - x_{n_i}}{\lambda_{n_i}} + Ax_{n_i} - \frac{\alpha_{n_i}}{\lambda_{n_i}}(u - x_{n_i} + \lambda_{n_i}Ax_{n_i}) \rangle \\
&= \langle v - y_{n_i}, Av - Ax_{n_i} - \frac{y_{n_i} - x_{n_i}}{\lambda_{n_i}} + \frac{\alpha_{n_i}}{\lambda_{n_i}}(u - x_{n_i} + \lambda_{n_i}Ax_{n_i}) \rangle \\
&= \langle v - y_{n_i}, Av - Ay_{n_i} \rangle + \langle v - y_{n_i}, Ay_{n_i} - Ax_{n_i} \rangle \\
&\quad - \langle v - y_{n_i}, \frac{y_{n_i} - x_{n_i}}{\lambda_{n_i}} - \frac{\alpha_{n_i}}{\lambda_{n_i}}(u - x_{n_i} + \lambda_{n_i}Ax_{n_i}) \rangle \\
&\geq \langle v - y_{n_i}, Ay_{n_i} - Ax_{n_i} \rangle - \langle v - y_{n_i}, \frac{y_{n_i} - x_{n_i}}{\lambda_{n_i}} - \frac{\alpha_{n_i}}{\lambda_{n_i}}(u - x_{n_i} + \lambda_{n_i}Ax_{n_i}) \rangle.
\end{aligned}$$

Noting that  $\alpha_{n_i} \rightarrow 0$  and  $\|y_{n_i} - x_{n_i}\| \rightarrow 0$ , we deduce  $0 \leq \langle v - z, w \rangle$ . Hence,  $z \in T^{-1}(0)$  and thus  $z \in \text{VI}(C, A)$ .

Next, we show that  $z \in \text{VI}(C, B)$ . Define a mapping  $T^\dagger$  as follows

$$T^\dagger v = \begin{cases} Bv^\dagger + N_C v^\dagger, & v^\dagger \in C, \\ \emptyset, & v^\dagger \notin C. \end{cases}$$

Let  $(v^\dagger, w^\dagger) \in G(T^\dagger)$ . Since  $w^\dagger - Bv^\dagger \in N_C v$  and  $x_{n+1} \in C$ , we obtain  $0 \leq \langle v^\dagger - x_{n+1}, w^\dagger - Bv^\dagger \rangle$ . By virtue of  $x_{n+1} = \text{proj}_C[x_n - \mu_n \beta_n B y_n + \beta_n(y_n - x_n)]$ , we obtain

$$0 \leq \langle v^\dagger - x_{n+1}, \mu_n \beta_n B y_n + x_{n+1} - x_n - \beta_n(y_n - x_n) \rangle,$$

that is,

$$0 \leq \langle v^\dagger - x_{n+1}, B y_n + \frac{x_{n+1} - x_n}{\beta_n \mu_n} - \frac{y_n - x_n}{\mu_n} \rangle.$$

Therefore, we have

$$\begin{aligned}
\langle v^\dagger - x_{n_i+1}, w^\dagger \rangle &\geq \langle v^\dagger - x_{n_i+1}, Bv^\dagger \rangle \\
&\geq \langle v^\dagger - x_{n_i+1}, Bv^\dagger \rangle - \langle v^\dagger - x_{n_i+1}, \frac{x_{n_i+1} - x_{n_i}}{\beta_{n_i} \mu_{n_i}} + B y_{n_i} - \frac{y_{n_i} - x_{n_i}}{\mu_{n_i}} \rangle \\
&= \langle v^\dagger - x_{n_i+1}, Bv^\dagger - B y_{n_i} - \frac{x_{n_i+1} - x_{n_i}}{\beta_{n_i} \mu_{n_i}} + \frac{y_{n_i} - x_{n_i}}{\mu_{n_i}} \rangle \\
&= \langle v^\dagger - x_{n_i+1}, Bv^\dagger - B y_{n_i} \rangle + \langle v^\dagger - x_{n_i+1}, Bx_{n_i+1} - B y_{n_i} \rangle \\
&\quad - \langle v^\dagger - x_{n_i+1}, \frac{x_{n_i+1} - x_{n_i}}{\beta_{n_i} \mu_{n_i}} - \frac{y_{n_i} - x_{n_i}}{\mu_{n_i}} \rangle \\
&\geq \langle v^\dagger - x_{n_i+1}, Bx_{n_i+1} - B y_{n_i} \rangle - \langle v - x_{n_i+1}, \frac{x_{n_i+1} - x_{n_i}}{\beta_{n_i} \mu_{n_i}} - \frac{y_{n_i} - x_{n_i}}{\mu_{n_i}} \rangle.
\end{aligned}$$

Noting that  $\|y_{n_i} - x_{n_i}\| \rightarrow 0$  and  $\|x_{n_i+1} - x_{n_i}\| \rightarrow 0$ , we get  $0 \leq \langle v^\dagger - z, w^\dagger \rangle$ . Hence,  $z \in T^{\dagger-1}(0)$  and  $z \in \text{VI}(C, B)$ . Thus,  $z \in \Omega$  and we have

$$\limsup_{n \rightarrow \infty} \langle \tilde{x} - u, \tilde{x} - y_n \rangle = \lim_{n \rightarrow \infty} \langle \tilde{x} - u, \tilde{x} - y_{n_i} \rangle = \langle \tilde{x} - u, \tilde{x} - z \rangle \leq 0.$$

□

Finally, we prove our main result.

**Theorem 1.** Suppose that  $\Omega = VI(C, A) \cap VI(C, B) \neq \emptyset$ . Assume that  $\{\alpha_n\}$ ,  $\{\beta_n\}$ ,  $\{\lambda_n\}$  and  $\{\mu_n\}$  satisfy the following restrictions (C1)–(C4). Then  $\{x_n\}$  defined by (5) converges strongly to  $\tilde{x} = proj_{\Omega}(u)$ .

**Proof.** First, we have Propositions 2–4 in hand. In terms of (4), we have

$$\begin{aligned}\|y_n - \tilde{x}\|^2 &= \|proj_C[\alpha_n u + (1 - \alpha_n)(x_n - \lambda_n A x_n)] - proj_C[x^* - (1 - \alpha_n)\lambda_n A x^*]\|^2 \\ &\leq \langle \alpha_n(u - x^*) + (1 - \alpha_n)[(x_n - \lambda_n A x_n) - (x^* - \lambda_n A x^*)], y_n - x^* \rangle \\ &\leq \alpha_n \langle \tilde{x} - u, \tilde{x} - y_n \rangle + (1 - \alpha_n) \|(x_n - \lambda_n A x_n) - (\tilde{x} - \lambda_n A \tilde{x})\| \|y_n - \tilde{x}\| \\ &\leq \alpha_n \langle \tilde{x} - u, \tilde{x} - y_n \rangle + (1 - \alpha_n) \|x_n - \tilde{x}\| \|y_n - \tilde{x}\| \\ &\leq \alpha_n \langle \tilde{x} - u, \tilde{x} - y_n \rangle + \frac{1 - \alpha_n}{2} \|x_n - \tilde{x}\|^2 + \frac{1}{2} \|y_n - \tilde{x}\|^2.\end{aligned}$$

It follows that

$$\|y_n - \tilde{x}\|^2 \leq (1 - \alpha_n) \|x_n - \tilde{x}\|^2 + 2\alpha_n \langle \tilde{x} - u, \tilde{x} - y_n \rangle.$$

Therefore,

$$\begin{aligned}\|x_{n+1} - \tilde{x}\|^2 &\leq (1 - \beta_n) \|x_n - \tilde{x}\|^2 + \beta_n \|y_n - \tilde{x}\|^2 \\ &\leq (1 - \alpha_n \beta_n) \|x_n - \tilde{x}\|^2 + 2\alpha_n \beta_n \langle \tilde{x} - u, \tilde{x} - y_n \rangle.\end{aligned}$$

By Lemma 3 and the above inequality, we deduce  $x_n \rightarrow \tilde{x}$ . This completes the proof.  $\square$

If we take  $u = 0$ , then we have the following Algorithm 2.

---

#### Algorithm 2:

For initial value  $x_0 \in C$ . Assume the sequence  $\{x_n\}$  has been constructed. Compute the next iteration  $\{x_{n+1}\}$  by the following manner

$$\begin{cases} y_n = proj_C[(1 - \alpha_n)(x_n - \lambda_n A x_n)], \\ x_{n+1} = proj_C[x_n - \mu_n \beta_n B y_n + \beta_n(y_n - x_n)], n \geq 0. \end{cases} \quad (11)$$


---

**Corollary 1.** Suppose that  $\Omega = VI(C, A) \cap VI(C, B) \neq \emptyset$ . Assume that  $\{\alpha_n\}$ ,  $\{\beta_n\}$ ,  $\{\lambda_n\}$  and  $\{\mu_n\}$  satisfy the following restrictions (C1)–(C4). Then  $\{x_n\}$  defined by (11) converges strongly to the minimum norm element  $\tilde{x}$  in  $\Omega$ .

#### 4. Conclusions

In this paper, we investigated the variational inequality problem. We suggest an extragradient type method for finding the common solutions of two variational inequalities. We prove the strong convergence of the method under the mild conditions. Noting that in our suggested iterative sequence (Equation (5)), the involved operators  $A$  and  $B$  require some form of strong monotonicity. A natural question arises, i.e., how to weaken these assumptions?

**Author Contributions:** All the authors have contributed equally to this paper. All the authors have read and approved the final manuscript.

**Funding:** This research was partially supported by the grants NSFC61362033 and NZ17015.

**Acknowledgments:** The authors are thankful to the anonymous referees for their careful corrections and valuable comments on the original version of this paper.

**Conflicts of Interest:** The authors declare no conflict of interest.

## References

1. Stampacchi, G. Formes bilinéaires coercitives sur les ensembles convexes. *C. R. Acad. Sci.* **1964**, *258*, 4413–4416.
2. Alber, Y.-I.; Iusem, A.-N. Extension of subgradient techniques for nonsmooth optimization in Banach spaces. *Set Valued Anal.* **2001**, *9*, 315–335. [[CrossRef](#)]
3. Bello Cruz, J.-Y.; Iusem, A.-N. A strongly convergent direct method for monotone variational inequalities in Hilbert space. *Numer. Funct. Anal. Optim.* **2009**, *30*, 23–36. [[CrossRef](#)]
4. Cho, S.-Y.; Qin, X.; Yao, J.-C.; Yao, Y. Viscosity approximation splitting methods for monotone and nonexpansive operators in Hilbert spaces. *J. Nonlinear Convex Anal.* **2018**, *19*, 251–264.
5. Dong, Q.-L.; Cho, Y.-J.; Zhong, L.-L.; Rassias, T.-M. Inertial projection and contraction algorithms for variational inequalities. *J. Glob. Optim.* **2018**, *7*, 687–704. [[CrossRef](#)]
6. Dong, Q.-L.; Cho, Y.-J.; Rassias, T.-M. The projection and contraction methods for finding common solutions to variational inequality problems. *Optim. Lett.* **2018**, *12*, 1871–1896. [[CrossRef](#)]
7. Facchinei, F.; Pang, J.-S. *Finite-Dimensional Variational Inequalities and Complementarity Problems*; Springer: New York, NY, USA, 2003; Volumes 1 and 2.
8. He, B.-S.; Yang, Z.-H.; Yuan, X.-M. An approximate proximal-extragradient type method for monotone variational inequalities. *J. Math. Anal. Appl.* **2004**, *300*, 362–374. [[CrossRef](#)]
9. Bello Cruz, J.-Y.; Iusem, A.-N. Convergence of direct methods for paramonotone variational inequalities. *Comput. Optim. Appl.* **2010**, *46*, 247–263. [[CrossRef](#)]
10. Iiduka, H.; Takahashi, W. Weak convergence of a projection algorithm for variational inequalities in a Banach space. *J. Math. Anal. Appl.* **2008**, *339*, 668–679. [[CrossRef](#)]
11. Li, C.-L.; Jia, Z.-F.; Postolache, M. New convergence methods for nonlinear uncertain variational inequality problems. *J. Nonlinear Convex Anal.* **2018**, *19*, 2153–2164.
12. Lions, J.-L.; Stampacchia, G. Variational inequalities. *Comm. Pure Appl. Math.* **1967**, *20*, 493–517. [[CrossRef](#)]
13. Lu, X.; Xu, H.-K.; Yin, X. Hybrid methods for a class of monotone variational inequalities. *Nonlinear Anal.* **2009**, *71*, 1032–1041. [[CrossRef](#)]
14. Solodov, M.-V.; Svaiter, B.-F. A new projection method for variational inequality problems. *SIAM J. Control Optim.* **1999**, *37*, 765–776. [[CrossRef](#)]
15. Solodov, M.-V.; Tseng, P. Modified projection-type methods for monotone variational inequalities. *SIAM J. Control Optim.* **1996**, *34*, 1814–1830. [[CrossRef](#)]
16. Thakur, B.-S.; Postolache, M. Existence and approximation of solutions for generalized extended nonlinear variational inequalities. *J. Inequal. Appl.* **2013**, *2013*, 590. [[CrossRef](#)]
17. Wang, S.-H.; Zhao, M.-L.; Kumam, P.; Cho, Y.-J. A viscosity extragradient method for an equilibrium problem and fixed point problem in Hilbert space. *J. Fixed Point Theory Appl.* **2018**, *20*, 19. [[CrossRef](#)]
18. Xu, H.-K.; Kim, T.-H. Convergence of hybrid steepest-descent methods for variational inequalities. *J. Optim. Theory Appl.* **2003**, *119*, 185–201. [[CrossRef](#)]
19. Yao, Y.; Chen, R.; Xu, H.-K. Schemes for finding minimum-norm solutions of variational inequalities. *Nonlinear Anal.* **2010**, *72*, 3447–3456. [[CrossRef](#)]
20. Yao, Y.; Liou, Y.-C.; Kang, S.-M. Approach to common elements of variational inequality problems and fixed point problems via a relaxed extragradient method. *Comput. Math. Appl.* **2010**, *59*, 3472–3480. [[CrossRef](#)]
21. Yao, Y.; Liou, Y.-C.; Postolache, M. Self-adaptive algorithms for the split problem of the demicontractive operators. *Optimization* **2018**, *67*, 1309–1319. [[CrossRef](#)]
22. Yao, Y.; Postolache, M.; Liou, Y.-C. Variant extragradient-type method for monotone variational inequalities. *Fixed Point Theory Appl.* **2013**, *2013*, 185. [[CrossRef](#)]
23. Yao, Y.; Postolache, M.; Liou, Y.-C.; Yao, Z.-S. Construction algorithms for a class of monotone variational inequalities. *Optim. Lett.* **2016**, *10*, 1519–1528. [[CrossRef](#)]
24. Yao, Y.; Qin, X.; Yao, J.-C. Projection methods for firmly type nonexpansive operators. *J. Nonlinear Convex Anal.* **2018**, *19*, 407–415.
25. Yao, Y.; Shahzad, N. Strong convergence of a proximal point algorithm with general errors. *Optim. Lett.* **2012**, *6*, 621–628. [[CrossRef](#)]
26. Yao, Y.; Yao, J.-C.; Liou, Y.-C.; Postolache, M. Iterative algorithms for split common fixed points of demicontractive operators without priori knowledge of operator norms. *Carpathian J. Math.* **2018**, *34*, 459–466.

27. Zegeye, H.; Yao, Y. Minimum-norm solution of variational inequality and fixed point problem in Banach spaces. *Optimization* **2015**, *64*, 453–471. [[CrossRef](#)]
28. Zhao, J.; Liang, Y.-S.; Liu, Y.-L.; Cho, Y.-J. Split equilibrium, variational inequality and fixed point problems for multi-valued mappings in Hilbert spaces. *Appl. Comput. Math.* **2018**, *17*, 271–283.
29. Yao, Y.; Postolache, M.; Yao, J.-C. An iterative algorithm for solving the generalized variational inequalities and fixed points problems. *Mathematics* **2019**, *7*, 61. [[CrossRef](#)]
30. Yao, Y.; Postolache, M.; Yao, J.-C. Iterative algorithms for generalized variational inequalities. *U. Politeh. Buch. Ser. A* **2019**, in press.
31. Korpelevich, G.-M. An extragradient method for finding saddle points and for other problems. *Ekon. Matemat. Metod.* **1976**, *12*, 747–756.
32. Bnouhachem, A.; Noor, M.-A.; Hao, Z. Some new extragradient iterative methods for variational inequalities. *Nonlinear Anal.* **2009**, *70*, 1321–1329. [[CrossRef](#)]
33. Censor, Y.; Gibali, A.; Reich, S. Strong convergence of subgradient extragradient methods for the variational inequality problem in Hilbert space. *Optim. Methods Softw.* **2011**, *26*, 827–845. [[CrossRef](#)]
34. Iusem, A.-N.; Svaiter, B.-F. A variant of Korpelevich’s method for variational inequalities with a new search strategy. *Optimization* **1997**, *42*, 309–321. [[CrossRef](#)]
35. Iusem, A.-N.; Lucambio Pérez, L.-R. An extragradient-type algorithm for non-smooth variational inequalities. *Optimization* **2000**, *48*, 309–332. [[CrossRef](#)]
36. Khobotov, E.-N. Modification of the extra-gradient method for solving variational inequalities and certain optimization problems. *USSR Comput. Math. Math. Phys.* **1989**, *27*, 120–127. [[CrossRef](#)]
37. Censor, Y.; Gibali, A.; Reich, S. Extensions of Korpelevich’s extragradient method for the variational inequality problem in Euclidean space. *Optimization* **2012**, *61*, 1119–1132. [[CrossRef](#)]
38. Censor, Y.; Gibali, A.; Reich, S. The subgradient extragradient method for solving variational inequalities in Hilbert space. *J. Optim. Theory Appl.* **2011**, *148*, 318–335. [[CrossRef](#)]
39. Censor, Y.; Gibali, A.; Reich, S. A von Neumann alternating method for finding common solutions to variational inequalities. *Nonlinear Anal.* **2012**, *75*, 4596–4603. [[CrossRef](#)]
40. Zaslavski, A.-J. The extragradient method for finding a common solution of a finite family of variational inequalities and a finite family of fixed point problems in the presence of computational errors. *J. Math. Anal. Appl.* **2013**, *400*, 651–663. [[CrossRef](#)]
41. Takahashi, W.; Toyoda, M. Weak convergence theorems for nonexpansive mappings and monotone mappings. *J. Optim. Theory Appl.* **2003**, *118*, 417–428. [[CrossRef](#)]
42. Suzuki, T. Strong convergence theorems for infinite families of nonexpansive mappings in general Banach spaces. *Fixed Point Theory Appl.* **2005**, *2005*, 103–123. [[CrossRef](#)]
43. Xu, H.-K. Iterative algorithms for nonlinear operators. *J. Lond. Math. Soc.* **2002**, *66*, 240–256. [[CrossRef](#)]



© 2019 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).