



Article

Universality in Short Intervals of the Riemann Zeta-Function Twisted by Non-Trivial Zeros

Antanas Laurinčikas ^{1,†}  and Darius Šiaučiūnas ^{2,*,†} 

¹ Institute of Mathematics, Faculty of Mathematics and Informatics, Vilnius University, Naugarduko str. 24, LT-03225 Vilnius, Lithuania; antanas.laurincikas@mif.vu.lt

² Regional Development Institute, Šiauliai University, P. Višinskio str. 25, LT-76351 Šiauliai, Lithuania

* Correspondence: darius.siauciunas@su.lt

† These authors contributed equally to this work.

Received: 3 October 2020; Accepted: 20 October 2020; Published: 3 November 2020



Abstract: Let $0 < \gamma_1 < \gamma_2 < \dots \leq \gamma_k \leq \dots$ be the sequence of imaginary parts of non-trivial zeros of the Riemann zeta-function $\zeta(s)$. Using a certain estimate on the pair correlation of the sequence $\{\gamma_k\}$ in the intervals $[N, N + M]$ with $N^{1/2+\varepsilon} \leq M \leq N$, we prove that the set of shifts $\zeta(s + ih\gamma_k)$, $h > 0$, approximating any non-vanishing analytic function defined in the strip $\{s \in \mathbb{C} : 1/2 < \text{Res} < 1\}$ with accuracy $\varepsilon > 0$ has a positive lower density in $[N, N + M]$ as $N \rightarrow \infty$. Moreover, this set has a positive density for all but at most countably $\varepsilon > 0$. The above approximation property remains valid for certain compositions $F(\zeta(s))$.

Keywords: Montgomery pair correlation conjecture; non-trivial zeros; Riemann zeta-function; universality

MSC: 11M06; 11M26

1. Introduction

The Riemann zeta-function $\zeta(s)$, $s = \sigma + it$, is defined, for $\sigma > 1$, by

$$\zeta(s) = \sum_{m=1}^{\infty} \frac{1}{m^s} = \prod_p \left(1 - \frac{1}{p^s}\right)^{-1},$$

where the infinite product is taken over all prime numbers, and has analytic continuation over the whole complex plane, except for the point $s = 1$ which is a simple pole with residue 1. The function $\zeta(s)$ and its value distribution play an important role not only in analytic number theory but in mathematics in general.

It is well known by a Bohr and Courant work [1] that the set of values of $\zeta(\sigma + it)$ with any fixed $\sigma \in (1/2, 1]$ is dense in \mathbb{C} . Voronin obtained [2] the infinite-dimensional version of the Bohr–Courant theorem, proving the so-called universality of $\zeta(s)$. This means that every non-vanishing analytic function in the strip $D = \{s \in \mathbb{C} : 1/2 < \sigma < 1\}$ can be approximated by shifts $\zeta(s + i\tau)$. We recall the modern version of the Voronin theorem. Denote by \mathcal{K} the class of compact subsets of the strip D with connected complements, and by $H_0(K)$ with $K \in \mathcal{K}$ the class of continuous non-vanishing functions on K that are analytic in the interior of K . Then, for $K \in \mathcal{K}$, $f(s) \in H_0(K)$ and every $\varepsilon > 0$, the inequality

$$\liminf_{T \rightarrow \infty} \frac{1}{T} \text{meas} \left\{ \tau \in [0, T] : \sup_{s \in K} |\zeta(s + i\tau) - f(s)| < \varepsilon \right\} > 0$$

is true; see, for example, [3–6]. Thus, we have that there are infinitely many shifts $\zeta(s + i\tau)$ approximating a given function $f(s) \in H_0(K)$.

The above theorem is of continuous type because τ in shifts $\zeta(s + i\tau)$ can take arbitrary real values. If τ runs over a certain discrete set, then we have the discrete universality that was proposed in [7]. Denote by $\#A$ the cardinality of a set A , and suppose that N runs over the set of non-negative integers. If K and $f(s)$ are as above, then we have, for $h > 0$ and $\varepsilon > 0$,

$$\liminf_{N \rightarrow \infty} \frac{1}{N+1} \# \left\{ 0 \leq k \leq N : \sup_{s \in K} |\zeta(s + ikh) - f(s)| < \varepsilon \right\} > 0.$$

Approximations of analytic functions by more general discrete shifts were considered in [8–10].

Denote by $\gamma_1 < \gamma_2 < \dots \leq \gamma_k \leq \dots$ the positive imaginary parts of non-trivial zeros $\rho_k = \beta_k + i\gamma_k$ of the function $\zeta(s)$. Discrete universality theorems with shifts $\zeta(s + ih\gamma_k)$ were obtained in [11,12]. In [11], for this the Riemann hypothesis was used, while in [12], the weak form of the Montgomery pair correlation conjecture [13] was involved. More precisely, the estimate, for $c > 0$,

$$\sum_{\substack{0 < \gamma_k, \gamma_l \leq T \\ |\gamma_k - \gamma_l| < c / \log T}} 1 \ll_c T \log T, \quad T \rightarrow \infty, \tag{1}$$

was required. Analogical results for more general functions were given in [14,15].

On the other hand, all above theorems are non-effective in the sense that any concrete shift approximating a given analytic function is not known. This shortcoming leads to the idea of universality in intervals as short as possible containing τ with approximating property. The first result in this direction was obtained in [16].

Theorem 1. *Suppose that $T^{1/3}(\log T)^{26/15} \leq H \leq T, K \in \mathcal{K}$ and $f(s) \in H_0(K)$. Then, for every $\varepsilon > 0$,*

$$\liminf_{T \rightarrow \infty} \frac{1}{H} \text{meas} \left\{ \tau \in [T, T + H] : \sup_{s \in K} |\zeta(s + i\tau) - f(s)| < \varepsilon \right\} > 0.$$

The aim of this paper is the universality of the function $\zeta(s)$ in short intervals with shifts $\zeta(s + ih\gamma_k)$. In this case, the estimate (1) is not sufficient. Therefore, for $N^{1/2+\varepsilon} \leq M \leq N$ with $\varepsilon > 0$, we use the following hypothesis:

$$\sum_{\substack{k=N \\ \gamma_k - \gamma_l < c / \log N}}^{N+M} \sum_{l=N}^{N+M} 1 \ll_c M, \tag{2}$$

which, as estimate (1), also gives a certain information on the pair correlation of non-trivial zeros, differently from estimate (1), however, in short intervals.

Theorem 2. *Suppose that $N^{1/2+\varepsilon} \leq M \leq N$, and estimate (2) are true. Let $K \in \mathcal{K}$ and $f(s) \in H_0(K)$. Then, for every $\varepsilon > 0$ and $h > 0$,*

$$\liminf_{N \rightarrow \infty} \frac{1}{M+1} \# \left\{ N \leq k \leq N + M : \sup_{s \in K} |\zeta(s + ih\gamma_k) - f(s)| < \varepsilon \right\} > 0.$$

Moreover, “lim inf” can be replaced by “lim” for all but at most countably many $\varepsilon > 0$.

Theorem 2 has a generalization for certain compositions $F(\zeta(s))$. Denote by $H(D)$ the space of analytic functions on the strip D endowed with the topology of uniform convergence on compacta. Moreover, let

$$S = \{g \in H(D) : \text{either } g(s) \neq 0 \text{ for all } s \in D, \text{ or } g(s) \equiv 0\},$$

and, for the operator $F : H(D) \rightarrow H(D)$ and distinct complex numbers a_1, \dots, a_r ,

$$H_{a_1, \dots, a_r; F}(D) = \{g \in H(D) : g(s) \neq a_j \text{ for all } s \in D, j = 1, \dots, r\} \cup \{F(0)\}.$$

Then we have

Theorem 3. Suppose that estimate (2) is true, $N^{1/2+\varepsilon} \leq M \leq N$, and $F : H(D) \rightarrow H(D)$ is a continuous operator such that $F(S) \supset H_{a_1, \dots, a_r; F}(D)$. For $r = 1$, let $K \in \mathcal{K}$ and $f(s)$ be a continuous $\neq a_1$ function on K , and analytic in the interior of K . For $r \geq 2$, let K be an arbitrary compact subset of D , and $f(s) \in H_{a_1, \dots, a_r; F}(D)$. Then, for every $\varepsilon > 0$ and $h > 0$,

$$\liminf_{N \rightarrow \infty} \frac{1}{M+1} \# \left\{ N \leq k \leq N+M : \sup_{s \in K} |F(\zeta(s+ih\gamma_k)) - f(s)| < \varepsilon \right\} > 0.$$

Moreover “lim inf” can be replaced by “lim” for all but at most countably many $\varepsilon > 0$.

For example, the operators $F(g) = \sin g$ and $F(g) = \sinh g$ satisfy the hypotheses of Theorem 3 with $a_1 = -1$ and $a_2 = 1$.

The proofs of Theorems 2 and 3 use probabilistic limit theorems for measures in the space $H(D)$. Denote by $\mathcal{B}(\mathbb{X})$ the Borel σ -field of the space \mathbb{X} . The main limit theorem will be proved for

$$P_{N,M,h}(A) = \frac{1}{M+1} \# \{N \leq k \leq N+M : \zeta(s+ih\gamma_k) \in A\}, A \in \mathcal{B}(H(D)),$$

as $N \rightarrow \infty$. We divide its proof into four sections.

2. A Limit Theorem on the Torus

Denote by γ the unit circle on the complex plane, by \mathbb{P} the set of all prime numbers, and define the set

$$\Omega = \prod_{p \in \mathbb{P}} \gamma_p,$$

where $\gamma_p = \gamma$ for all $p \in \mathbb{P}$. With the product topology and pointwise multiplication, the torus Ω is a compact topological Abelian group. Therefore, on $(\Omega, \mathcal{B}(\Omega))$, the probability Haar measure m_H can be defined, and we have the probability space $(\Omega, \mathcal{B}(\Omega), m_H)$. Denote by $\omega(p)$ the p th component of an element $\omega \in \Omega$, $p \in \mathbb{P}$.

In this section, we will prove a limit theorem for

$$Q_{N,M,h}(A) = \frac{1}{M+1} \# \left\{ N \leq k \leq N+M : \left(p^{-ih\gamma_k} : p \in \mathbb{P} \right) \in A \right\}, A \in \mathcal{B}(\Omega),$$

as $N \rightarrow \infty$.

Before the statement of a limit theorem for $Q_{N,M,h}$ as $N \rightarrow \infty$, we will recall some useful results that will be used in its proof. Denote by $N(T)$ the number of non-trivial zeros of $\zeta(s)$ in the region $\{s \in \mathbb{C} : 0 < t < T\}$.

Lemma 1 (von Mangoldt formula). For $T \rightarrow \infty$,

$$N(T) = \frac{T}{2\pi} \log \frac{T}{2\pi e} + O(\log T).$$

For the proof, see, for example, [17].

Denote by $N(\sigma, T)$ the number of zeros $\rho = \beta + i\gamma$ of $\zeta(s)$ with $\beta > \sigma$ and $|\gamma| < T$.

Lemma 2. Suppose that $H \geq T^\alpha$ with $\alpha > 27/82$. Then, for $1/2 < \sigma < 1$, uniformly in σ ,

$$N(\sigma, T + H) - N(\sigma, T) = O\left(\frac{H}{\sigma - 1/2}\right).$$

Proof of the lemma can be found in [18].

For positive $u \neq 1$, denote by $\Lambda(u)$ the von Mangoldt function if $u \in \mathbb{N} \setminus \{1\}$, and zero, otherwise.

Lemma 3. For positive $x \neq 1$ and $T \rightarrow \infty$,

$$\sum_{0 < \gamma_k < T} x^{\rho_k} = \left(\Lambda(x) - x\Lambda\left(\frac{1}{x}\right)\right) \frac{T}{2\pi} + O\left(T^{(1/2)+\varepsilon}\right)$$

with every $\varepsilon > 0$.

Proof. The lemma is Theorem 2 of [19] with $a = 0$. \square

Lemma 4. Suppose that $N^{1/2+\varepsilon} \leq M \leq N$ with $\varepsilon > 0$. Then, for positive $x \neq 1$, as $N \rightarrow \infty$,

$$\sum_{k=N}^{N+M} x^{\rho_k} \ll_x \frac{M}{\sqrt{\log M}}.$$

Proof. Since

$$\frac{N}{\log N} \ll \gamma_N \ll \frac{N}{\log N'}$$

in view of Lemma 3,

$$\sum_{\gamma_N < \gamma \leq \gamma_{N+M}} x^\rho = \left(\Lambda(x) - x\Lambda\left(\frac{1}{x}\right)\right) \frac{\gamma_{N+M} - \gamma_N}{2\pi} + O\left(\frac{N^{1/2+\varepsilon}}{\sqrt{\log N}}\right). \tag{3}$$

An application of Lemma 1 gives

$$N + M = \sum_{\gamma \leq \gamma_{N+M}} 1 = \frac{\gamma_{N+M}}{2\pi} \log \frac{\gamma_{N+M}}{2\pi e} + O(\log N)$$

and

$$N = \sum_{\gamma \leq \gamma_N} 1 = \frac{\gamma_N}{2\pi} \log \frac{\gamma_N}{2\pi e} + O(\log N).$$

Therefore,

$$\gamma_{N+M} = \frac{2\pi(N + M)}{\log(\gamma_{N+M}/(2\pi e))} + O(1)$$

and

$$\gamma_N = \frac{2\pi N}{\log(\gamma_N/(2\pi e))} + O(1).$$

Hence,

$$\gamma_{N+M} - \gamma_N \leq \frac{2\pi(N + M)}{\log(\gamma_N/(2\pi e))} - \frac{2\pi N}{\log(\gamma_N/(2\pi e))} + O(1) \ll \frac{M}{\log N} + O(1) \ll \frac{M}{\log M}. \tag{4}$$

This together with Equation (3) proves the lemma. \square

Now, we state the limit theorem for $Q_{N,M,h}$.

Theorem 4. Suppose that, for any $\varepsilon > 0$, $N^{1/2+\varepsilon} \leq M \leq N$. Then $Q_{N,M,h}$ converges weakly to the Haar measure m_H as $N \rightarrow \infty$.

Proof. Denote by $g_{N,M,h}(\underline{k})$, $\underline{k} = (k_p : k_p \in \mathbb{Z}, p \in \mathbb{P})$, the Fourier transform of $Q_{N,M,h}$, i.e.,

$$g_{N,M,h}(\underline{k}) = \int_{\Omega} \left(\prod_{p \in \mathbb{P}}^* \omega^{k_p}(p) \right) dQ_{N,M,h},$$

where the star “*” means that only a finite number of integers k_p are distinct from zero. Thus, by the definition of $Q_{N,M,h}$,

$$g_{N,M,h}(\underline{k}) = \frac{1}{M+1} \sum_{k=N}^{N+M} \exp \left\{ -ih \gamma_k \sum_{p \in \mathbb{P}}^* k_p \log p \right\}. \tag{5}$$

Clearly,

$$g_{N,M,h}(\underline{0}) = 1. \tag{6}$$

Now, suppose that $\underline{k} \neq \underline{0}$. Since the set $\{\log p : p \in \mathbb{P}\}$ is linearly independent within the field of rational numbers \mathbb{Q} , in that case we have

$$a \stackrel{def}{=} \sum_{p \in \mathbb{P}}^* k_p \log p \neq 0.$$

Thus, we will estimate the sum

$$\sum_{k=N}^{N+M} \exp\{iha \gamma_k\}.$$

It is easily seen that

$$\begin{aligned} \sum_{k=N}^{N+M} \left(\exp\{ha\beta_k\} - \exp\left\{\frac{1}{2}ha\right\} \right) &\ll_{h,a} \sum_{k=N}^{N+M} \left| \exp\left\{ha\left(\beta_k - \frac{1}{2}\right)\right\} - 1 \right| \\ &\ll_{h,a} \sum_{k=N}^{N+M} \left| \beta_k - \frac{1}{2} \right| = \sum'_{k=N}^{N+M} \left| \beta_k - \frac{1}{2} \right| + \sum''_{k=N}^{N+M} \left| \beta_k - \frac{1}{2} \right|, \end{aligned} \tag{7}$$

where $|\beta_k - 1/2| \leq 1/\log \log M$ in Σ' , and $|\beta_k - 1/2| > 1/\log \log M$ in Σ'' . Obviously,

$$\sum'_{k=N}^{N+M} \left| \beta_k - \frac{1}{2} \right| \leq \frac{M}{\log \log M}. \tag{8}$$

Therefore, by Lemma 2 and estimate (2),

$$\sum''_{k=N}^{N+M} \left| \beta_k - \frac{1}{2} \right| \ll \sum''_{\gamma_N < \gamma \leq \gamma_{N+M}} 1 \ll \frac{M \log \log M}{\log M}.$$

This, and estimates (7) and (8) show that

$$\sum_{k=N}^{N+M} \exp\{(\beta_k + i\gamma_k)ha\} - \sum_{k=N}^{N+M} \exp\left\{\left(\frac{1}{2} + i\gamma_k\right)ha\right\} \ll_{h,a} \frac{M}{\log \log M}. \tag{9}$$

Lemma 4 with $x = \exp\{ha\}$ implies

$$\sum_{k=N}^{N+M} \exp\{(\beta_k + i\gamma_k)ha\} \ll_{h,a} \frac{M}{\sqrt{\log M}}.$$

Therefore, in view of estimate (9),

$$\sum_{k=N}^{N+M} \exp\{iha\gamma_k\} \ll_{h,a} \sum_{k=N}^{N+M} \exp\left\{\left(\frac{1}{2} + i\gamma_k\right) ha\right\} \ll_{h,a} \frac{M}{\log \log M}.$$

Thus, by Equation (5),

$$g_{N,M,h}(k) \ll_{h,a} \frac{1}{\log \log M}.$$

This together with Equation (6) shows that

$$\lim_{N \rightarrow \infty} g_{N,M,h}(k) = \begin{cases} 1 & \text{if } k = 0, \\ 0 & \text{if } k \neq 0, \end{cases}$$

and the lemma is proved because the right-hand side of the latter equality is the Fourier transform of the measure m_H . \square

3. A Limit Theorem for Absolutely Convergent Series

Let $\theta > 1/2$ be a fixed number, and $v_n(m) = \exp\{-(m/n)^\theta\}$ for $m, n \in \mathbb{N}$. Extend the function $\omega(p)$ to the set \mathbb{N} by setting

$$\omega(m) = \prod_{\substack{p^l | m \\ p^{l+1} \nmid m}} \omega^l(p),$$

and define

$$\zeta_n(s) = \sum_{m=1}^{\infty} \frac{v_n(m)}{m^s}$$

and

$$\zeta_n(s, \omega) = \sum_{m=1}^{\infty} \frac{\omega(m)v_n(m)}{m^s}.$$

Then the latter series are absolutely convergent for $\sigma > 1/2$ [5]. Consider the function $u_n : \Omega \rightarrow H(D)$ defined by

$$u_n(\omega) = \zeta_n(s, \omega).$$

The absolute convergence of the series $\zeta_n(s, \omega)$ implies the continuity of u_n . For $A \in \mathcal{B}(H(D))$, define

$$P_{N,M,n,h}(A) = \frac{1}{M+1} \#\{N \leq k \leq N+M : \zeta_n(s + ih\gamma_k) \in A\}.$$

Theorem 5. Suppose that $N^{1/2+\epsilon} \leq M \leq N$. Then $P_{N,M,n,h}$ converges weakly to the measure $m_H u_n^{-1} \stackrel{\text{def}}{=} V_n$.

Proof. The theorem follows from the equality

$$P_{N,M,n,h}(A) = Q_{N,M,h}(u_n^{-1}A) = Q_{N,M,h}u_n^{-1}(A), \quad A \in \mathcal{B}(H(D)),$$

continuity of the function u_n , Theorem 4 and Theorem 5.1 of [20]. \square

The weak convergence of $P_{N,M,h}$ is closely connected to that of V_n as $n \rightarrow \infty$. Define

$$\zeta(s, \omega) = \prod_{p \in \mathbb{P}} \left(1 - \frac{\omega(p)}{p^s}\right)^{-1}.$$

Then $\zeta(s, \omega)$ is an $H(D)$ -valued random element on the probability space $(\Omega, \mathcal{B}(\Omega), m_H)$ [5]. We recall that the latter infinite product, for almost all ω , is uniformly convergent on compact subsets $K \subset D$. Denote by P_ζ the distribution of the random element $\zeta(s, \omega)$, i.e.,

$$P_\zeta(A) = m_H\{\omega \in \Omega : \zeta(s, \omega) \in A\}, \quad A \in \mathcal{B}(H(D)).$$

The following statement is very important.

Proposition 1. *The probability measure V_n converges weakly to measure P_ζ as $n \rightarrow \infty$.*

Proof. For $A \in \mathcal{B}(H(D))$, define

$$R_T(A) = \frac{1}{T} \text{meas}\{\tau \in [0, T] : \zeta(s + i\tau) \in A\}.$$

It is known that R_T , as $T \rightarrow \infty$, converges weakly to P_ζ [5]. Moreover, R_T , as $T \rightarrow \infty$, and V_n , as $n \rightarrow \infty$, converge weakly to the same probability measure on $(H(D), \mathcal{B}(H(D)))$. Thus, V_n converges weakly to P_ζ as $n \rightarrow \infty$. \square

4. Mean Square Estimates in Short Intervals

To derive the weak convergence of $P_{N,M,h}$ from that of $P_{N,M,n,h}$ as $N \rightarrow \infty$, the estimate for

$$\sum_{k=N}^{N+M} |\zeta(\sigma + ih\gamma_k + it)|^2$$

with $t \in \mathbb{R}$ is needed.

We will use the following mean square estimate in short intervals.

Lemma 5. *Suppose that $\sigma, 1/2 < \sigma < 1$, is fixed and $T^{1/3}(\log T)^{26/15} \leq H \leq T$. Then, uniformly in H ,*

$$\int_{T-H}^{T+H} |\zeta(\sigma + it)|^2 \ll_\sigma H.$$

The lemma follows from Theorem 7.1 of [21], and was used in [16].

Lemma 6. *Suppose that $N^{1/2+\epsilon} \leq M \leq N$ and estimate (2) is true. Then, for every fixed $\sigma, 1/2 < \sigma < 1$, $h > 0$ and $t \in \mathbb{R}$,*

$$\sum_{k=N}^{N+M} |\zeta(\sigma + ih\gamma_k + it)| \ll_{\sigma,h} M(1 + |t|).$$

Proof. We will apply the Gallagher lemma connecting discrete mean squares with those continuous of some functions; for the proof, see Lemma 1.4 of [22]. Let $T_0, T \geq \delta > 0$ be real numbers, $\mathcal{T} \neq \emptyset$ be a finite set in the interval $[T_0 + \delta/2, T_0 + T - \delta/2]$,

$$N_\delta(x) = \sum_{\substack{t \in \mathcal{T} \\ |t-x| < \delta}} 1$$

and let $S(x)$ be a complex-valued continuous function on $[T_0, T + T_0]$ having a continuous derivative on $(T_0, T + T_0)$. Then the Gallagher lemma asserts that

$$\sum_{t \in \mathcal{T}} N_\delta^{-1}(t) |S(t)|^2 \leq \frac{1}{\delta} \int_{T_0}^{T_0+T} |S(x)|^2 dx + \left(\int_{T_0}^{T_0+T} |S(x)|^2 dx \int_{T_0}^{T_0+T} |S'(x)|^2 dx \right)^{1/2}. \tag{10}$$

We apply the Gallagher lemma for the function $\zeta(s + ikh\gamma_k + it)$. In our case $\delta = c/\log N$, $T_0 = h\gamma_N - \delta/2$, $T = h\gamma_{N+M} - h\gamma_N + \delta/2$ and $\mathcal{T} = \{h\gamma_N, h\gamma_{N+1}, \dots, h\gamma_{N+M}\}$. By estimate (2), we have

$$\sum_{k=N}^{N+M} N_\delta(h\gamma_k) = \sum_{k=N}^{N+M} \sum_{\substack{l=N \\ |\gamma_k - \gamma_l| < c/(h \log N)}}^{N+M} 1 \ll_h M. \tag{11}$$

Now, an application of the Gallagher lemma gives

$$\begin{aligned} \sum_{k=N}^{N+M} |\zeta(\sigma + ih\gamma_k + it)| &= \sum_{k=N}^{N+M} \sqrt{N_\delta(h\gamma_k)N^{-1}(h\gamma_k)} |\zeta(\sigma + ih\gamma_k + it)| \\ &\leq \left(\sum_{k=N}^{N+M} N_\delta(h\gamma_k) \sum_{k=N}^{N+M} N^{-1}(h\gamma_k) |\zeta(\sigma + ih\gamma_k + it)|^2 \right)^{1/2} \\ &\ll_h \sqrt{M} \sqrt{\log N} \left(\int_{h\gamma_N - \delta/2}^{h\gamma_{N+M}} |\zeta(\sigma + i\tau + it)|^2 d\tau \right. \\ &\quad \left. + \left(\int_{h\gamma_N - \delta}^{h\gamma_{N+M}} |\zeta(\sigma + i\tau + it)|^2 d\tau \int_{h\gamma_N - \delta}^{h\gamma_{N+M}} |\zeta'(\sigma + i\tau + it)|^2 d\tau \right)^{1/2} \right)^{1/2}. \tag{12} \end{aligned}$$

The estimate (4) gives with certain $c_h > 0$

$$\int_{h\gamma_N - \delta}^{h\gamma_{N+M}} |\zeta(\sigma + i\tau + it)|^2 d\tau \ll \int_{h\gamma_N - \delta - |t|}^{h\gamma_N + c_h(M/\log M) + |t|} |\zeta(\sigma + i\tau)|^2 d\tau. \tag{13}$$

If $c_h(M/\log M) + |t| \leq h\gamma_N$, then, in view of Lemma 5, the right-hand side of (13) is

$$\ll_{\sigma, h} \frac{M}{\log M} + |t| \ll_{\sigma, h} \frac{M}{\log M} (1 + |t|).$$

If $c_h(M/\log M) + |t| > h\gamma_N$, then

$$h\gamma_N + c_h \frac{M}{\log M} + |t| < 2 \left(c_h \frac{M}{\log M} + |t| \right)$$

and

$$h\gamma_N - \delta > h\gamma_N - 2c_h \frac{M}{\log M} - 2|t| > -h\gamma_N c_h \frac{M}{\log M} - |t| > -2 \left(c_h \frac{M}{\log M} + |t| \right).$$

Thus, in this case,

$$\int_{h\gamma_N - \delta}^{h\gamma_{N+M}} |\zeta(\sigma + i\tau + it)|^2 d\tau \ll_h \int_{-2(c_h(M/\log M) + |t|)}^{2(c_h(M/\log M) + |t|)} |\zeta(\sigma + i\tau)|^2 d\tau \ll_{\sigma, h} \frac{M}{\log M} (1 + |t|).$$

This together with estimate (13) shows that

$$\int_{h\gamma_N - \delta}^{h\gamma_{N+M}} |\zeta(\sigma + i\tau + it)|^2 d\tau \ll_{\sigma, h} \frac{M}{\log M} (1 + |t|). \tag{14}$$

Estimate (14) and an application of the Cauchy integral formula lead to the bound

$$\int_{h\gamma_N - \delta}^{h\gamma_{N+M}} |\zeta'(\sigma + i\tau + it)|^2 d\tau \ll_{\sigma, h} \frac{M}{\log M} (1 + |t|).$$

This, estimate (14) and (12) prove the lemma. \square

Now, we are ready to state an approximation lemma.

5. Approximation in the Mean

Denote by ρ the metric in $H(D)$ which induces the topology of uniform convergence on compacta. More precisely, for $g_1, g_2 \in H(D)$,

$$\rho(g_1, g_2) = \sum_{l=1}^{\infty} 2^{-l} \frac{\sup_{s \in K_l} |g_1(s) - g_2(s)|}{1 + \sup_{s \in K_l} |g_1(s) - g_2(s)|},$$

where $\{K_l : l \in \mathbb{N}\} \subset D$ is a sequence of compact subsets such that

$$D = \bigcup_{l=1}^{\infty} K_l,$$

$K_l \subset K_{l+1}$ for all $l \in \mathbb{N}$, and every compact $K \subset D$ lies in a certain K_l .

Lemma 7. *Suppose that $N^{1/2+\varepsilon} \leq M \leq N$ and (2) is true. Then, for every $h > 0$,*

$$\lim_{n \rightarrow \infty} \limsup_{N \rightarrow \infty} \frac{1}{M+1} \sum_{k=N}^{N+M} \rho(\zeta(s + ih\gamma_k), \zeta_n(s + ih\gamma_k)) = 0.$$

Proof. In view of the definition of the metric ρ , it suffices to show that, for every compact $K \subset D$,

$$\lim_{n \rightarrow \infty} \limsup_{N \rightarrow \infty} \frac{1}{M+1} \sum_{k=N}^{N+M} \sup_{s \in K} |\zeta(s + ih\gamma_k) - \zeta_n(s + ih\gamma_k)| = 0. \tag{15}$$

Thus, let $K \subset D$ be a fixed compact set. Denote the points of K by $s = \sigma + iv$, and fix $\varepsilon > 0$ such that $1/2 + 2\varepsilon \leq \sigma \leq 1 - \varepsilon$ for $s \in K$. It is known [5] that

$$\zeta_n(s) = \frac{1}{2\pi i} \int_{\theta-i\infty}^{\theta+i\infty} \zeta(s+z) l_n(z) \frac{dz}{z},$$

where

$$l_n(s) = \frac{s}{\theta} \Gamma(s/\theta) n^s,$$

$\Gamma(s)$ is the Euler gamma-function, and θ comes from the definition of $v_n(m)$. Let $\theta_1 > 0$. From this, we have

$$\zeta(s) - \zeta_n(s) = \frac{1}{2\pi i} \int_{-\theta-i\infty}^{-\theta+i\infty} \zeta(s+z) l_n(z) \frac{dz}{z} + R_n(s),$$

with

$$R_n(s) = \frac{l_n(1-s)}{1-s}.$$

Therefore, as in the proof of Lemma 12 of [16], we find that

$$\begin{aligned} & \frac{1}{M+1} \sum_{k=N}^{N+M} \sup_{s \in K} |\zeta(s + ih\gamma_k) - \zeta_n(s + ih\gamma_k)| \\ & \ll \int_{-\infty}^{\infty} \frac{1}{M} \sum_{k=N}^{N+M} \left| \zeta \left(\frac{1}{2} + \varepsilon + i(h\gamma_k + t) \right) \right| \sup_{s \in K} \frac{|l_n(1/2 + \varepsilon - s + it)|}{|1/2 + \varepsilon - s + it|} dt \\ & \quad + \frac{1}{M} \sum_{k=N}^{N+M} \sup_{s \in K} |R_n(s + ih\gamma_k)| \stackrel{def}{=} I_1 + I_2. \end{aligned} \tag{16}$$

Denote by c_1, c_2, \dots positive constants. In view of the well-known estimate

$$\Gamma(\sigma + it) \ll \exp\{-c_1|t|\}, \tag{17}$$

we find that

$$\frac{|l_n(1/2 + \varepsilon - s + it)|}{|1/2 + \varepsilon - s + it|} \ll n^{-\varepsilon} \exp\{-c_2|t - v|\} \ll_{K,\varepsilon} n^{-\varepsilon} \exp\{-c_3|t|\}.$$

Therefore, by Lemma 5,

$$I_1 \ll_{K,\varepsilon} n^{-\varepsilon} \int_{-\infty}^{\infty} (1 + |t|) \exp\{-c_3|t|\} dt \ll_{K,\varepsilon} n^{-\varepsilon}. \tag{18}$$

Similarly, taking into account inequality (17), we find

$$\begin{aligned} I_2 &\ll \frac{n^{1/2-2\varepsilon}}{M} \sum_{k=N}^{N+M} \exp\{-c_4|h\gamma_k - v|\} \ll_K \frac{n^{1/2-2\varepsilon}}{M} \sum_{k=N}^{N+M} \exp\{-c_5h\gamma_k\} \\ &\ll_K \frac{n^{1/2-2\varepsilon}}{M} \sum_{k=N}^{N+M} \exp\{-c_6h(k/\log k)\} \ll_{K,h} \frac{n^{1/2-2\varepsilon}}{M}. \end{aligned}$$

This, Equations (18) and (16) prove (15). \square

6. A Limit Theorem for $\zeta(s)$

Using the results of Sections 3 and 4 leads to a limit theorem for $P_{N,M,h}$.

Theorem 6. *Suppose that $N^{1/2+\varepsilon} \leq M \leq N$ and estimate (2) is true. Then $P_{N,M,h}$ converges weakly to P_{ζ} as $N \rightarrow \infty$.*

Proof. In a certain probability space with measure μ define the random variable $\theta_{N,M,h}$ with the distribution

$$\mu\{\theta_{N,M,h} = h\gamma_k\} = \frac{1}{M+1}, \quad k = N, N+1, \dots, N+M,$$

and consider the $H(D)$ -valued random element

$$X_{N,M,n,h} = X_{N,M,n,h}(s) = \zeta_n(s + i\theta_{N,M,h}).$$

Moreover, let $X_n = X_n(s)$ be the $H(D)$ -valued random element with the distribution V_n . Then, by Theorem 5,

$$X_{N,M,n,h} \xrightarrow[N \rightarrow \infty]{\mathcal{D}} X_n, \tag{19}$$

where $\xrightarrow{\mathcal{D}}$ denotes the convergence in distribution. Moreover, by Proposition 1,

$$X_n \xrightarrow[n \rightarrow \infty]{\mathcal{D}} P_{\zeta}. \tag{20}$$

Define one more $H(D)$ -valued random element

$$X_{N,M,h} = X_{N,M,h}(s) = \zeta(s + i\theta_{N,M,h}).$$

Then, using Lemma 7, we find that, for every $\varepsilon > 0$,

$$\begin{aligned} &\lim_{n \rightarrow \infty} \limsup_{N \rightarrow \infty} \mu \{ \rho(X_{N,M,h}, X_{N,M,n,h}) \geq \varepsilon \} \\ &\leq \lim_{n \rightarrow \infty} \limsup_{N \rightarrow \infty} \frac{1}{\varepsilon(M+1)} \sum_{k=N}^{N+M} \rho(\zeta(s + ih\gamma_k), \zeta_n(s + ih\gamma_k)) = 0. \end{aligned}$$

Now, this, Equations (19) and (20) together with Theorem 4.2 of [20] show that

$$X_{N,M,h} \xrightarrow[n \rightarrow \infty]{\mathcal{D}} P_\zeta,$$

and theorem is proved. \square

For $A \in \mathcal{B}(H(D))$, define

$$P_{N,M,h,F}(A) = \frac{1}{M+1} \# \{N \leq k \leq N+M : F(\zeta(s+ih\gamma_k)) \in A\}.$$

Corollary 1. *Suppose that $F : H(D) \rightarrow H(D)$ is a continuous operator, and (2) is true. Then $P_{N,M,h,F}$ converges weakly to $P_\zeta F^{-1}$ as $N \rightarrow \infty$.*

Proof. The corollary follows from Theorem 5, continuity of F , equality

$$P_{N,M,h,F} = P_{N,M,h} F^{-1},$$

and Theorem 5.1 of [20]. \square

7. Proof of Universality

Theorems 2 and 3 are derived from Theorem 6 and Corollary 1, respectively, by using the Mergelyan theorem on the approximation of analytic functions by polynomials [23].

Proof of Theorem 2. We recall that

$$S = \{g \in H(D) : \text{either } g(s) \neq 0 \text{ for all } s \in D, \text{ or } g(s) \equiv 0\},$$

It is well known, see, for example, [5], that the support of the measure P_ζ is the set S . Define the set

$$G_\epsilon = \left\{ g \in H(D) : \sup_{s \in K} |g(s) - e^{p(s)}| < \frac{\epsilon}{2} \right\},$$

where $p(s)$ is a polynomial. Obviously, $e^{p(s)} \in S$. Therefore, G_ϵ is an open neighbourhood of an element of the support of the measure P_ζ . Thus, by a property of the support,

$$P_\zeta(G_\epsilon) > 0. \tag{21}$$

This, Theorem 6 and the equivalent of weak convergence in terms of open sets show that

$$\liminf_{N \rightarrow \infty} P_{N,M,h}(G_\epsilon) \geq P_\zeta(G_\epsilon) > 0.$$

Hence, by the definition of $P_{N,M,h}$ and G_ϵ ,

$$\liminf_{N \rightarrow \infty} \frac{1}{M+1} \# \left\{ N \leq k \leq N+M : \sup_{s \in K} |\zeta(s+ih\gamma_k) - e^{p(s)}| < \frac{\epsilon}{2} \right\} > 0. \tag{22}$$

Now, we apply the Mergelyan theorem and choose the polynomial $p(s)$ satisfying

$$\sup_{s \in K} |f(s) - e^{p(s)}| < \frac{\epsilon}{2}. \tag{23}$$

This and inequality (22) prove the first part of the theorem.

To prove the second part of the theorem, define the set

$$\hat{G}_\varepsilon = \left\{ g \in H(D) : \sup_{s \in K} |g(s) - f(s)| < \varepsilon \right\}.$$

Then the set \hat{G}_ε is a continuity set of the measure P_ζ for all but at most countably many $\varepsilon > 0$. This remark, Theorem 6 and the equivalent of weak convergence of probability measures in terms of open sets show that

$$\lim_{N \rightarrow \infty} P_{N,M,h}(\hat{G}_\varepsilon) = P_\zeta(\hat{G}_\varepsilon) \tag{24}$$

for all but at most countably many $\varepsilon > 0$. Inequality (23) implies the inclusion $G_\varepsilon \subset \hat{G}_\varepsilon$. Therefore, in view of inequality (21), we have $P_\zeta(\hat{G}_\varepsilon) > 0$. This, Equation (24) and the definitions of $P_{N,M,h}$ and \hat{G}_ε prove the second part of the theorem. \square

Proof of Theorem 3. Denote by S_F the support of the measure $P_\zeta F^{-1}$. We observe that S_F contains the closure of the set $H_{a_1, \dots, a_r; F}(D)$. Actually, let $g \in H_{a_1, \dots, a_r; F}(D)$ and G be any open neighborhood of g . Then the set $F^{-1}G$ is open as well, and lies in S . Hence, $P_\zeta(F^{-1}G) > 0$ because S is the support of P_ζ . Therefore,

$$P_\zeta F^{-1}(G) = P_\zeta(F^{-1}G) > 0.$$

This shows that S_F contains the set $H_{a_1, \dots, a_r; F}(D)$ and its closure.

Case $r = 1$. By the Mergelyan theorem, there exists a polynomial $p(s)$ such that

$$\sup_{s \in K} |f(s) - p(s)| < \frac{\varepsilon}{2}. \tag{25}$$

Then, $p(s) \neq a_1$ for all $s \in K$ if ε is small enough. Therefore, by the Mergelyan theorem again, we find a polynomial $q(s)$ such that

$$\sup_{s \in K} \left| (p(s) - a_1) - e^{q(s)} \right| < \frac{\varepsilon}{4}. \tag{26}$$

Since $g_1(s) \stackrel{def}{=} e^{q(s)} + a_1 \in H_{a_1; F}(D)$, the set

$$\mathcal{G}_\varepsilon = \left\{ g \in H(D) : \sup_{s \in K} |g(s) - g_1(s)| < \frac{\varepsilon}{2} \right\}$$

is an open subset of S_F . Hence,

$$P_\zeta F^{-1}(\mathcal{G}_\varepsilon) > 0. \tag{27}$$

This inequality together with Corollary 1, inequalities (25) and (26) prove the theorem in the case of the lower density.

In the case of density, consider the set \hat{G}_ε defined in the proof of Theorem 2 which is a continuity set of the measure $P_\zeta F^{-1}$ for all but at most countably many $\varepsilon > 0$. Therefore, by Corollary 1,

$$\lim_{N \rightarrow \infty} P_{N,M,h,F}(\hat{G}_\varepsilon) = P_\zeta F^{-1}(\hat{G}_\varepsilon). \tag{28}$$

Inequalities (25) and (26) show that $\mathcal{G}_\varepsilon \subset \hat{G}_\varepsilon$. Thus, by inequality (27), $P_\zeta F^{-1}(\hat{G}_\varepsilon) > 0$. This, Equation (28) and the definitions of $P_{N,M,h,F}$ and \hat{G}_ε prove the theorem in the case of density.

Case $r \geq 2$. In this case, the function $f(s)$ lies in S_F . Therefore, the Mergelyan theorem is not needed, and the theorem follows immediately from Corollary 1. \square

Author Contributions: Conceptualization, A.L. and D.Š.; methodology, A.L. and D.Š.; investigation, A.L. and D.Š.; writing—original draft preparation, A.L. and D.Š.; writing—review and editing, A.L. and D.Š. All authors have read and agreed to the published version of the manuscript.

Funding: The research of the first author is funded by the European Social Fund (project number 09.3.3-LMT-K-712-01-0037) under grant agreement with the Research Council of Lithuania (LMT LT).

Conflicts of Interest: The authors declare no conflict of interest.

References

- Bohr, H.; Courant, R. Neue Anwendungen der Theorie der Diophantischen Approximationen auf die Riemannsche Zetafunktion. *J. Reine Angew. Math.* **1914**, *144*, 249–274.
- Voronin, S.M. Theorem on the “universality” of the Riemann zeta-function. *Izv. Ross. Akad. Nauk.* **1975**, *9*, 443–453. [[CrossRef](#)]
- Bagchi, B. The Statistical Behaviour and Universality Properties of the Riemann Zeta-Function and Other Allied Dirichlet Series. Ph.D. Thesis, Indian Statistical Institute, Calcutta, India, 1981.
- Gonek, S.M. Analytic Properties of Zeta and L -Functions. Ph.D. Thesis, University of Michigan, Ann Arbor, MI, USA, 1975.
- Laurinćikas, A. *Limit Theorems for the Riemann Zeta-Function*; Kluwer Academic Publishers: Dordrecht, The Netherlands; Boston, MA, USA; London, UK, 1996.
- Steuding, J. *Value-Distribution of L -Functions*; Lecture Notes Math; Springer: Berlin/Heidelberg, Germany; New York, NY, USA, 2007; Volume 1877.
- Reich, A. Werteverteilung von Zetafunktionen. *Arch. Math.* **1980**, *34*, 440–451. [[CrossRef](#)]
- Laurinćikas, A. Discrete universality of the Riemann zeta-function and uniform distribution modulo 1. *Petersb. Math. J.* **2019**, *30*, 103–110. [[CrossRef](#)]
- Macaitienė, R. The discrete universality of the Riemann zeta-function with respect to uniformly distributed shifts. *Arch. Math.* **2017**, *108*, 271–281. [[CrossRef](#)]
- Pańkowski, Ł. Joint universality for dependent L -functions. *Ramanujan J.* **2018**, *45*, 181–195. [[CrossRef](#)]
- Garunkštis, R.; Laurinćikas, A. The Riemann hypothesis and universality of the Riemann zeta-function. *Math. Slovaca* **2018**, *68*, 741–748. [[CrossRef](#)]
- Garunkštis, R.; Laurinćikas, A.; Macaitienė, R. Zeros of the Riemann zeta-function and its universality. *Acta Arith.* **2017**, *181*, 127–142. [[CrossRef](#)]
- Montgomery, H.L. The pair correlation of zeros of the zeta-function. In *Analytic Number Theory*; Diamond, H.G., Ed.; American Mathematical Society: Providence, RI, USA, 1973; pp. 181–193.
- Balćiūnas, A.; Garbaliuskienė, V.; Karaliūnaitė, J.; Macaitienė, R.; Petuškinaitė, J.; Rimkevićienė, A. Joint discrete approximation of a pair of analytic functions by periodic zeta-functions. *Math. Modell. Anal.* **2020**, *25*, 71–87. [[CrossRef](#)]
- Laurinćikas, A.; Petuškinaitė, J. Universality of Dirichlet L -functions and non-trivial zeros of the Riemann zeta-function. *Sb. Math.* **2019**, *210*, 1753–1773. [[CrossRef](#)]
- Laurinćikas, A. Universality of the Riemann zeta-function in short intervals. *J. Number Theory* **2019**, *204*, 279–295. [[CrossRef](#)]
- Titchmarsh, E.C. *The Theory of the Riemann zeta-Function*, 2nd ed.; Heath-Brown, D.R., Ed.; Clarendon Press: Oxford, UK, 1986.
- Karatsuba, A.A.; Voronin, S.M. *The Riemann Zeta-Function*; Walter de Gruiter: New York, NY, USA; Berlin, Germany, 1992.
- Steuding, J. The roots of the equation $\zeta(s) = a$ are uniformly distributed modulo one. In *Anal. Probab. Methods Number Theory*; Laurinćikas, A., Ed.; TEV: Vilnius, Lithuania, 2012; pp. 243–249.
- Billingsley, P. *Convergence of Probability Measures*; Willey: New York, NY, USA, 1968.
- Ivić, A. *The Riemann Zeta-Function: The Theory of the Riemann Zeta-Function with Applications*; John Wiley & Sons: New York, NY, USA, 1985.

22. Montgomery, H.L. *Topics in Multiplicative Number Theory*; Lecture Notes in Mathematics; Springer: Berlin/Heidelberg, Germany; New York, NY, USA, 1971; Volume 227.
23. Mergelyan, S.N. *Uniform Approximations to Functions of a Complex Variable*; Series and Approximation; American Mathematical Society: Providence, RI, USA, 1969; pp. 294–391.

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



© 2020 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).