

Supplementary Document

Simple logistic regression of predisposing, enabling, and need factors of healthcare services utilization in relation to poor men's health behaviors (n = 257).

Variable	Crude OR (95% CI)	p-value
Age	1.014 (0.966, 1.065)	0.565
Distance to healthcare facilities	1.018 (0.879, 1.179)	0.809
Educational level		
Tertiary level	1	
Up to the secondary level	3.313 (0.429, 25.584)	0.251
Status		
Single/Divorce	1	
Married	1.045 (0.291, 3.759)	0.946
Occupation		
JBPM	1	
APM	1.515 (0.507, 4.531)	0.457
PDRM	0.784 (0.168, 3.666)	0.757
Monthly income (RM)		
2500 - 11000	1	
<2500	1.407 (0.552, 3.589)	0.474
Household income (RM)		
2500 - >11000	1	
<2500	1.196 (0.464, 3.085)	0.711
Comorbid		
No	1	
Yes	0.735 (0.232, 2.333)	0.601
Sought treatment at public healthcare facilities		
Ever	1	
Never	3.528 (0.457, 27.203)	0.226
Advice from Wife/spouse		
Not necessary	1	
Yes	0.944 (0.326, 2.730)	0.915

Advice from Family members/relatives			
	Not necessary	1	
	Yes	0.458 (0.178, 1.182)	0.106
Advice from Parents			
	Not necessary	1	
	Yes	2.029 (0.772, 5.333)	0.151
Advice from Friends/colleagues			
	Not necessary	1	
	Yes	0.388 (0.125, 1.203)	0.101
Financial capabilities			
	Not necessary	1	
	Yes	0.653 (0.224, 1.906)	0.435
Leisure time			
	Not necessary	1	
	Yes	1.865 (0.710, 4.900)	0.206
Transportation			
	Not necessary	1	
	Yes	0.510 (0.185, 1.365)	0.188