

**Table S2.** COVID-19 symptoms at the time of treatment prescription by subjects' immune status.

COVID-19 signs and symptoms	N	Immunocompromised			p-value <sup>†</sup>
		Overall, N = 143*	No, N = 37*	Yes, N = 106*	
<b>Fever</b>	143	78 (55%)	16 (43%)	62 (58%)	0.109
<b>Cough</b>	143	75 (52%)	25 (68%)	50 (47%)	0.032
<b>Ageusia-dysgeusia</b>	143	7 (4.9%)	4 (11%)	3 (2.8%)	0.074
<b>Anosmia</b>	143	7 (4.9%)	4 (11%)	3 (2.8%)	0.074
<b>Pharyngodynia</b>	143	45 (31%)	8 (22%)	37 (35%)	0.134
<b>Asthenia</b>	143	31 (22%)	9 (24%)	22 (21%)	0.650
<b>Headache</b>	143	32 (22%)	9 (24%)	23 (22%)	0.741
<b>Myalgia</b>	143	22 (15%)	2 (5.4%)	20 (19%)	0.051
<b>GI signs and symptoms</b>	143	15 (10%)	6 (16%)	9 (8.4%)	0.216
<b>Dyspnoea - tachypnoea</b>	143	5 (3.5%)	3 (8.1%)	2 (1.9%)	0.109

\*Median (IQR) or Frequency (%)

<sup>†</sup>Pearson's Chi-squared test; Fisher's exact test

Abbreviations: GI=gastrointestinal