

Supplementary Material

Table S2. Obstetrical and neonatal outcomes regarding different TSH concentrations: below and equal/over 3.0 mIU/L.

Variable	Group 1 (TSH < 3.0 mIU/L)	Group 1 (TSH ≥ 3.0 mIU/L)	Group 2	Group 3 (Controls)	<i>p</i>
Miscarriage, <i>n</i> (%)	3 (7.0)	1 (6.7)	0 (0.0)	3 (7.7)	0.359
Preterm birth, <i>n</i> (%)	4 (9.3)	1 (6.7)	1 (3.0)	5 (12.8)	0.508
Cervical insufficiency, <i>n</i> (%)	4 (9.3)	2 (13.3)	1 (3.0)	1 (2.6)	0.264
Gestational diabetes, <i>n</i> (%)	7 (16.3)	3 (20.0)	3 (9.1)	10 (25.6)	0.323
Gestational hypertension, <i>n</i> (%)	2 (4.7)	1 (6.7)	0 (0.0)	4 (10.3)	0.253
Preeclampsia, <i>n</i> (%)	1 (2.3)	1 (6.7)	0 (0.0)	1 (2.6)	0.575
Weeks of gestation at birth, Me (Q1;Q3)	39.00 (38.00;40.00)	39.00 (38.00;40.00)	39.00 (38.00;40.00)	39.00 (38.00;40.00)	0.887
Cesarean section, <i>n</i> (%)	16 (47.1)	5 (35.7)	6 (33.3)	7 (21.9)	0.200
Birth weight, g					
Males, Me (Q1;Q3)	3500.00 (3290.00;3890.00)	3340.00 (3150.00;3730.00)	3160.00 (3000.00;3632.50)	3455.00 (3242.50;3747.50)	0.528
Females, M ± SD	3366.18 ± 556.92	3243.33 ± 387.34	3495.00 ± 292.73	3056.25 ± 777.18	0.365
Apgar [0-10], Me (Q1;Q3)	10.00 (10.00;10.00)	10.00 (10.00;10.00)	10.00 (9.25;10.00)	10.00 (10.00;10.00)	0.191
Apgar below 5	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	-
NICU stay of newborns, <i>n</i> (%)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	-
SGA/FGR, <i>n</i> (%)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	-

M – mean, SD – standard deviation, Me – median, Q1 – 1st quartile, Q3 – 3rd quartile, NICU- neonatal intensive care unit; SGA- small for gestational age; FGR- fetal growth restriction.