

Supplementary materials

Article

Changes in resurgent sodium current contribute to the hyperexcitability of muscles in patients with paramyotonia congenita

Chiung-Wei Huang,^{1,2} Hsing-Jung Lai,^{3,4} Pi-Chen Lin,⁵ Ming-Jen Lee^{3, 6 and *}

1. Department of Post Baccalaureate Medicine, Kaohsiung Medical University, Kaohsiung, Taiwan. g10054b@ms51.hinet.net
2. Department of Physiology, Kaohsiung Medical University, Kaohsiung, Taiwan. g10054b@ms51.hinet.net
3. Department of Neurology, National Taiwan University Hospital, Taipei, Taiwan. i5492111@gmail.com and mjlee@ntu.edu.tw
4. Department of Neurology, National Taiwan University Hospital Jinshan Branch, New Taipei City, Taiwan. i5492111@gmail.com
5. Division of Endocrinology and Metabolism, Department of Internal Medicine, Kaohsiung Medical University Hospital, Kaohsiung, Taiwan. pichli@kmu.edu.tw
6. Department of Neurology, National Taiwan University Hospital Yunlin Branch, Yunlin, Taiwan. mjlee@ntu.edu.tw

* Correspondence: mjlee@ntu.edu.tw; Tel.: +88-62-2312-3456 (ext 65336)

Figure S1




Figure S1. Sample sweeps shows the peak currents and sustained currents in WT, p.T1313M, and p.R1448H mutant Na_v1.4 channels