

Lipophilicity Study of Fumaric and Maleic Acids

Daria Klimoszek and Alina Pyka-Pajak *

Department of Analytical Chemistry, Faculty of Pharmaceutical Sciences in Sosnowiec,
Medical University of Silesia in Katowice, Jagiellońska 4, 41-200 Sosnowiec, Poland;
daria.klimoszek@sum.edu.pl

* Correspondence: apyka@sum.edu.pl; Tel.: +48-32-364-15-30




Figure S1. Spectrodensitograms of maleic acid (pink line) and fumaric acid (celadon line).

Table S1. Average of spot area [AU] of maleic and fumaric acids analyzed by NP-TLC.

| Microgram/spot | Average of spot area [AU] | |
|----------------|---------------------------|--------------|
| | (n=5) | |
| | Maleic acid | Fumaric acid |
| 4 | 2265 | 19633 |
| 8 | 4856 | 27825 |
| 12 | 8325 | 34571 |
| 16 | 12456 | 44674 |
| 20 | 14589 | 49092 |
| 24 | 16584 | 54909 |
| 28 | 18554 | 59402 |
| 32 | 20525 | 69180 |
| 36 | 24521 | 72796 |

| | | |
|----|-------|-------|
| 40 | 26538 | 80267 |
|----|-------|-------|




Figure S2. Calibration curve of fumaric acid tested by NP-TLC chromatography.




Figure S3. Calibration curve of maleic acid tested by NP-TLC chromatography.

Table S2. R_M values of maleic acid analyzed by RP-TLC technique using methanol-water mobile phase.

| Volume fraction of methanol in the methanol-water mobile phase | R_M values on chromatographic plates | | |
|--|--|----------------------|---------------------|
| | RP18WF ₂₅₄ | RP8F _{254s} | CNF _{254s} |
| 1 | - | -2.020 | - |
| 0.9 | - | -1.797 | - |
| 0.8 | -2.756 | -1.684 | -2.999 |
| 0.7 | -2.479 | -1.571 | -2.714 |
| 0.6 | -2.103 | -1.458 | -2.500 |
| 0.5 | -1.925 | -1.412 | -2.199 |
| 0.4 | -1.734 | -1.232 | -2.038 |
| 0.3 | -1.371 | -1.202 | -1.852 |
| 0.2 | -0.985 | -1.006 | -1.599 |

Table S3. R_M values of maleic acid analyzed by RP-TLC technique using dioxane-water mobile phase.

| Volume fraction of dioxane in the dioxane-water mobile phase | R_M values on chromatographic plates | | |
|--|--|----------------------|---------------------|
| | RP18WF ₂₅₄ | RP8F _{254s} | CNF _{254s} |
| 1 | - | - | - |
| 0.9 | - | - | -2.905 |
| 0.8 | -2.988 | -2.789 | -2.625 |
| 0.7 | -2.699 | -2.714 | -2.385 |
| 0.6 | -2.532 | -2.598 | -2.155 |
| 0.5 | -2.386 | -2.499 | -1.989 |
| 0.4 | -2.199 | -2.312 | -1.843 |
| 0.3 | -1.972 | -2.255 | -1.554 |
| 0.2 | -1.702 | -1.995 | -1.345 |

Table S4. R_M values of fumaric acid analyzed by RP-TLC technique using methanol-water mobile phase.

| Volume fraction of methanol in the methanol-water mobile phase | R_M values on chromatographic plates | | |
|--|--|----------------------|---------------------|
| | RP18WF ₂₅₄ | RP8F _{254s} | CNF _{254s} |
| 1 | -2.315 | -2.560 | -2.312 |
| 0.9 | -1.999 | -2.324 | -1.898 |
| 0.8 | -1.625 | -1.995 | -1.526 |
| 0.7 | -1.345 | -1.458 | -1.289 |
| 0.6 | -1.152 | -1.288 | -0.785 |
| 0.5 | -0.789 | -0.995 | -0.565 |
| 0.4 | -0.525 | -0.702 | -0.278 |
| 0.3 | 0.119 | -0.549 | 0.089 |
| 0.2 | 0.352 | -0.158 | 0.312 |

Table S5. R_M values of fumaric acid analyzed by RP-TLC technique using dioxane-water mobile phase.

| Volume fraction of dioxane in the dioxane-water mobile phase | R_M values on chromatographic plates | | |
|--|--|----------------------|---------------------|
| | RP18WF ₂₅₄ | RP8F _{254s} | CNF _{254s} |
| 1 | -2.899 | -2.890 | - |
| 0.9 | -2.481 | -2.184 | - |
| 0.8 | -2.209 | -1.738 | -2.712 |
| 0.7 | -1.858 | -1.531 | -2.108 |
| 0.6 | -1.614 | -1.287 | -1.802 |
| 0.5 | -1.425 | -0.857 | -1.489 |
| 0.4 | -1.135 | -0.425 | -1.089 |
| 0.3 | -0.785 | -0.175 | -0.712 |
| 0.2 | -0.395 | 0.135 | -0.315 |