



## Article

# Bioactive Compounds Produced by the Accompanying Microflora in Bulgarian Yoghurt

Ivan Ivanov<sup>1</sup>, Kaloyan Petrov<sup>2</sup> , Valentin Lozanov<sup>3</sup>, Iassen Hristov<sup>3</sup>, Zhengjun Wu<sup>4</sup>, Zhenmin Liu<sup>4</sup> and Penka Petrova<sup>1,\*</sup> 

<sup>1</sup> Institute of Microbiology, Bulgarian Academy of Sciences, Acad. G. Bontchev Str., bl. 26, 1113 Sofia, Bulgaria; ivanov.ivnt@gmail.com

<sup>2</sup> Institute of Chemical Engineering, Bulgarian Academy of Sciences, Acad. G. Bontchev Str., bl. 103, 1113 Sofia, Bulgaria; kaloian04@yahoo.com

<sup>3</sup> Department of Medical Chemistry and Biochemistry, Faculty of Medicine, Medical University of Sofia, 1000 Sofia, Bulgaria; boval@abv.bg (V.L.); iass\_mu@abv.bg (I.H.)

<sup>4</sup> State Key Laboratory of Dairy Biotechnology, Dairy Research Institute, Bright Dairy & Food Co., Ltd., Shanghai 201103, China; wuzhengjun@brightdairy.com (Z.W.); liuzhenmin@brightdairy.com (Z.L.)

\* Correspondence: pepipetrova@yahoo.com; Tel.: +359-878207286

**Abstract:** Bulgarian yoghurt is associated with health benefits and longevity of consumers. The specific microflora producing bioactive metabolites is responsible for this effect. The present study examines the biodiversity in four homemade yoghurts from regions containing endemic microflora. Metagenome sequencing indicated *Lactobacillus delbrueckii* subsp. *bulgaricus* and *Streptococcus thermophilus* were predominant in all samples. In addition, yoghurts contained accompanying lactic acid bacteria (LAB) including *Lacticaseibacillus paracasei*, *Lb. helveticus*, *Limosilactobacillus fermentum*, *Lb. rhamnosus*, *Lactococcus lactis*, *Pediococcus acidilactici*, *Leuconostoc mesenteroides*, and *Leuc. pseudomesenteroides*. A negligible amount of pollutant strains was found. Twenty-four LAB strains were isolated from the yoghurts and identified. *Lb. delbrueckii* subsp. *bulgaricus* strains were genotyped by randomly amplified polymorphic DNA–PCR (RAPD), multi-locus sequence typing (MLST), and pulse field gel electrophoresis (PFGE), which demonstrated their uniqueness and non-commercial origin. To estimate the bioactive metabolites produced by the accompanying microflora, yoghurts fermented by single LAB strains were analyzed using liquid chromatography and mass spectrometry (LC-MS). The fermented samples contained large amounts of free essential amino acids (arginine, citrulline, tryptophan, lysine, and histidine), the neuroprotector indole-3-propionic acid (IPA), and significant quantities of the cyclic antimicrobial peptides cyclo(phenylalanyl-prolyl) and cyclo(leucyloprolyl). The disclosure of these special qualities draws attention to the accompanying microflora as a source of potential probiotic strains that can fortify the yoghurts' content with bioactive compounds.

**Keywords:** Bulgarian yoghurt; *Lactobacillus bulgaricus*; *Streptococcus thermophilus*; metagenomics; RAPD; MLST; PFGE; IPA; cyclic peptides



**Citation:** Ivanov, I.; Petrov, K.; Lozanov, V.; Hristov, I.; Wu, Z.; Liu, Z.; Petrova, P. Bioactive Compounds Produced by the Accompanying Microflora in Bulgarian Yoghurt. *Processes* **2021**, *9*, 114. <https://doi.org/10.3390/pr9010114>

Received: 14 December 2020

Accepted: 5 January 2021

Published: 7 January 2021

**Publisher's Note:** MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



**Copyright:** © 2021 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

## 1. Introduction

For centuries, conventional yoghurt consumption has been associated with beneficial effects on human health. Recent studies show cardio-metabolic benefits and ameliorative effects on obesity, diabetes, and biomarkers of chronic diseases, including inflammation [1]. The regular intake of yoghurt lactic acid bacteria (LAB) and their metabolites also yields anti-carcinogenic, anti-oxidative, and immunomodulatory effects and reduces blood cholesterol [2]. LAB in yoghurt can regulate the microbiome supporting the gut barrier [3] and improve the immune system response [4,5].

The remarkable properties of Bulgarian yoghurt are due exclusively to its microbial content. The microbial flora of yoghurt is not a random set of species, but a unique microecological niche [6]. The most studied yoghurt starters rely on the protocollaboration between

*Lactobacillus delbrueckii* subsp. *bulgaricus* and *Streptococcus thermophilus* [7,8]. According to the FAO/WHO definition, yoghurt must contain no less than  $10^7$  viable colony-forming units (CFU) per gram of product. Starter cultures produce volatiles, which are responsible for the typical yoghurt aroma (acetaldehyde, diacetyl, acetoin, acetone, ethanol, and 2-butanone) [9], organic acids, prebiotics such as galactooligosaccharides (GOS) [10], and bioactive compounds. Bioactive peptides, such as bacteriocins, are connected with the antimicrobial effects against pathogens in the gastrointestinal tract, such as clostridia, *Enterobacteriaceae*, pseudomonads, and *Helicobacter pylori* [11–13]. Other peptides produced by *Lb. delbrueckii* subsp. *bulgaricus* are known to harmonize the blood pressure by angiotensin-converting enzyme (ACE) inhibitory activity [14].

Contemporary culture-independent techniques, such as metagenomics, allow the comprehensive investigation of the yoghurt consortia and reveal that they contain a wide range of accompanying microflora with no less useful properties [15]. These microbial communities are an object of increasing scientific interest, at least in three different directions: (i) investigation of the symbiosis and other internal interactions between LAB to select starters with better quality; (ii) elucidation of the specific dietetic properties of yoghurt; and (iii) determination of the specific compounds produced by lactic acid bacteria (LAB), which can resolve health problems without drug use. A study engaging healthy volunteers revealed that the daily intake of yoghurt fortified with *Lb. paracasei* (*L. casei* 431<sup>®</sup>) improves immune response by enhancing NK cell activity and increasing interleukin (IL)-12 and immunoglobulin (Ig) G1 levels, and IFN- $\gamma$  concentration [16]. Another strain of this species, *Lb. paracasei* CNCM I-1518, decreases bacterial translocation, gut dysbiosis, and ileal oxidative damage, and increases ileal  $\beta$ -defensin-1 expression in rats, suggesting an improvement in the intestinal barrier integrity [17]. *Lb. helveticus* H9 yoghurt possesses excellent ACE-inhibitory (ACE I) activity with high levels of antihypertensive peptides, such as Val-Pro-Pro and Ile-Pro-Pro [18,19]. *Lb. helveticus* LH-B02 is known to favor the formation of bioactive peptides, such as  $\alpha$ S1-CN f(24–32) and  $\beta$ -CN f(193–209) [20]. The patent-protected *Lb. rhamnosus* GG (ATCC 53103) is the world's most studied probiotic bacterium with more than 800 scientific studies. The authors claim that it is bile-stable, has a great avidity for human intestinal mucosal cells, and possesses various health benefits, such as the alleviation of gastroenteritis, diarrhea, eczema, and many other disorders [21–23].

Bulgarian yoghurt (kiselo mlyako) is the hallmark of the country, being one of the most popular types of yoghurt worldwide. The present study adds novel comprehensive information about Bulgarian yoghurt microflora to complement the discovery of *Lb. delbrueckii* subsp. *bulgaricus* discovery one hundred years ago [24]. A collection of homemade Bulgarian yoghurts from remote locations revealed unique microflora content. Considering the significant influence of both starter cultures and the accompanying LAB on yoghurt's beneficial properties, the present work aims to reveal the biodiversity of Bulgarian yoghurts using metagenomics, to isolate and identify new starters and accompanying LAB strains by engaging modern genetic approaches, and to study the bioactive metabolites produced from concomitant species using precise analytical methods.

## 2. Materials and Methods

### 2.1. Yogurt Sample Collection

Four samples were collected from remote territories known with endemic yoghurt microflora: Rila, Pirin, and Rhodope Mountains, and the Thracian valley. The samples derived from cow (Samples 1 and 3), goat (Sample 2), and buffalo milk (Sample 4). They were prepared according to Bulgarian national recipes passed on by generations of local people from the respective villages. By this technology, the milk for yoghurt production is boiled and then fermented from an artisanal starter culture. The owners of the samples declared that they did not use commercial starters. The samples were collected in the period October–December 2015 in sterile containers and were stored frozen at  $-20$  °C until their analysis in 2020.

## 2.2. Bacterial Strains, Media and Cultivation Conditions

The strains were isolated by the following procedure: 10–20 µL of the yoghurt was inoculated in 5 mL MRS broth selective medium (Roth, Germany) and cultivated 24 h either at 37 °C or 44 °C. Decimal dilutions (up to 10<sup>-9</sup>) of these cultures were mixed with MRS and M17 agar selective for lactobacilli, and lactic acid cocci, respectively, and then loaded in Petri dishes. The agar was stored with a temperature below 50 °C, the colonies appeared in the agar after 48 h of cultivation. The selection was performed at two different temperatures: at 37 °C and 44 °C. At 37 °C *Pediococcus* and *Lactococcus* strains appeared, at 44 °C—*Lb. delbrueckii* subsp. *bulgaricus* and *Str. thermophilus*. Lactic acid rod-shaped bacteria were cultivated in MRS medium; lactic acid cocci in M17 medium, at 42 °C, or 37 °C, respectively; at anaerobic conditions, using an Anaerocult® A mini (Merck KGaA, Darmstadt, Germany). The cultivation of strains was done in semi-skimmed milk, containing 1.2% fat and 4.1% lactose. For cell propagation aimed at DNA isolation, the strains were cultured in MRS for 24 h, at optimal temperature for each strain. Single colonies were picked and inoculated again in MRS or M17 broth (3 mL).

## 2.3. Total DNA Isolation from Yoghurt

Total DNA from yoghurt samples was extracted using a new method that combines the first steps of the method of Lick et al. (1996) [25], followed by biomass lysis and DNA purification by the use of PureLink™ Genomic DNA Mini Kit (Invitrogen™, Thermo Fisher Scientific, Waltham, MA, USA). Briefly, a yoghurt sample with volume 8 mL was mixed with 2.4 mL of 0.4 M NaOH and 1.2 mL of 0.25 M Trisodium citrate dihydrate. The tubes were vigorously shaken (at 2500 rpm/min) for 10 s and incubated at ambient temperature for 5 min. The following centrifugation (5000× g, 10 min) separated three fractions: upper white layer (fat), yellowish supernatant, and grainy pellet (microbial cells). The biomass was re-suspended in 500 µL of the remaining supernatant, and after addition of 150 µL 0.4 M NaOH and 75 µL 40% Trisodium citrate dihydrate, the mixture was vigorously shaken again, incubated at room temperature for 5 min, and centrifuged at 12,000× g for 3 min. The obtained pellet was resuspended in 200 µL of Lysozyme Digestion Buffer of the PureLink™ Genomic DNA Mini Kit. Then, the procedure as described by the manufacturer Invitrogen™ was followed. DNA concentration and quality were measured using a QB 3000 spectrophotometer (Quawell Ltd., Hong Kong, China).

## 2.4. Molecular Approaches in Strains' Identification

### 2.4.1. Isolation of Chromosomal DNA, PCR Amplification of the 16S rRNA Gene and Sequencing

Chromosomal DNA from LAB isolates was isolated from 24 h old cultures, grown in MRS-starch medium, using a GeneJET Genomic DNA Purification Kit (Thermo Fisher Scientific Inc., Waltham, MA, USA), and following the manufacturer's recommendations. PCR amplifications were prepared with Prime Taq premix (Genet Bio, Daejeon, Korea), in a total volume of 50 µL and final concentrations of primers 0.5 pmol/µL (Macrogen Inc., Amsterdam, The Netherlands) in a QB-96 Satellite Gradient Thermal Cycler (LKB Vertriebs GmbH, Vienna, Austria). The amplification of the 16S rRNA gene was performed with universal eubacterial primer pair: 27F: 5' AGAGTTTGATCCTGGCTCAG 3' and 1492R: 5' AAGGAGGTGATCCAGCC 3'. The final concentration of the template DNA was 2 ng/µL, the temperature profile was: 95 °C for 5 min, 35 cycles consisting of 94 °C for 1 min, 55 °C for 1 min, 72 °C for 2 min, followed by final elongation at 72 °C for 5 min. All obtained PCR amplification products were purified using the GFX PCR DNA and gel band purification kit (Amersham Biosciences) and then sequenced by Macrogen Inc. (Amsterdam, The Netherlands). Sequence comparison with the GenBank data was performed using BLAST and ClustalW programs.

#### 2.4.2. Randomly Amplified Polymorphic DNA–PCR (RAPD) Analysis

RAPD analysis was performed using the modified procedure of Cebeci et al. (2011) [26]. The primers used were 1254: 5'CCGCAGCCAA3' (10-mer), rpd1: 5'CCCGACTGA3' (9-mer), rpd2: 5'CCG AGACAC3' (9-mer), rpd3: 5'GCCTGTTCTT3' (10-mer), and rpd4: 5'-GCCAGAACAA-3' (10-mer). The PCR mix contained: DNA—1 µL (100 ng), 100 pmol primers—3 µL, 2× Phusion Master Mix (Thermo Scientific Inc., Waltham, MA, USA)—10 µL, 25 mmol MgCl<sub>2</sub>—2 µL, H<sub>2</sub>O—1 µL. The program followed the temperature profile: 98 °C—30 s, 4 cycles: denaturation 98 °C 10 s, annealing 36 °C—20 s, elongation 72 °C—15 s, 30 cycles 98 °C—10 s, 20 °C—20 s, 68 °C—15 s, 72 °C—5 min. The obtained products were separated in 2% agarose, for 2.5 h, at 50 V (5 v/cm). A molecular size standard of 100 bp was used (Thermo Scientific Inc., Waltham, MA, USA).

#### 2.4.3. Multi-Locus Sequence Typing (MLST)

This analysis was conducted as described by Naser et al. (2005) [27] using the primers listed in Table 1. The primer pairs targeted the genes *rpoB* encoding the β-subunit of bacterial RNA polymerase, *groEL* encoding the protein of the chaperonin family, *pheS* for the α-subunit of phenylalanine tRNA synthetase, and *pyrG* encoding CTP synthase. The sequencing was performed with the reverse primers. Divergence data were obtained by combined similarities in 4 housekeeping genes analyzed by MEGA6 software.

**Table 1.** Primers used for multi-locus sequence typing (MLST) amplification and sequencing.

Gene	Primer Name	Sequence (5'→3')	Annealing T °C	Product Position in Gene	Position in Genome <sup>1</sup>	Template Size (bp)
<i>rpoB</i>	rpoB_F	GGCGGAAAGAGTTATCGT	58	399–1170	334519–338184	772
	rpoB_R	GATGTCGGCTGGAGTGAT				
<i>groEL</i>	groEL_F	TCGGCAAGGACGGTGTT	57	500–1134	1392354–1393967	635
	groEL_R	GTGGATTACGGCTACGC				
<i>pheS</i>	pheS_F	CATCGGCATGAGCTACCA	55	369–1042	1283032–1284081	674
	pheS_R	CCTCCTGACGGAATTGTTG				
<i>pyrG</i>	pyrG_F	AAGCCGACCCAGCAATC	57	568–1309	301679–303298	742
	pyrG_R	AGCCCAGACGCAAGGTG				

<sup>1</sup> Note: Position in the genome of *Lb. delbrueckii* subsp. *bulgaricus* ATCC11842<sup>T</sup> (GenBank Accession No: NC\_008054).

#### 2.4.4. Pulse Field Gel Electrophoresis (PFGE)

The cells for PFGE were prepared using a partially modified procedure of Michaylova et al. (2007) [28]. The cells were grown overnight, to OD<sub>600</sub> 0.6–0.8, harvested, washed in 1 mL ET buffer (1 M TRIS/HCl, 100 mM EDTA, pH 8), and resuspended in 50 µL ET buffer. After incubation for 1 min at 45 °C, the suspension was mixed with 60 µL 1.5% low melting agarose (Sigma-Aldrich, St. Louis, MO, USA) in LMP buffer (10 mM TRIS/HCl, 100 mM EDTA, 20 mM NaCl, pH 8), and was pipetted in molds (Bio-Rad Laboratories, Hercules, CA, USA). The molds were treated with 500 µL Buffer A: 1 mM TRIS/HCl, 100 mM EDTA, 1 M NaCl, 0.5% (v/v) TRITON X 100, pH 8 + 10<sup>6</sup> U/mL Lysozyme (Merck KGaA, Darmstadt, Germany) + 75 U/mL mutanolysin (Sigma-Aldrich, St. Louis, MO, USA). Incubation for 18 h at 37 °C was followed by washing for 5 h at 55 °C with 500 mL Lysis solution, which contained 1mM TRIS/HCl, 250 mM EDTA, 1% SDS, pH 8, and 30 mU/mL (1 mg/mL) Proteinase K (Merck KGaA, Darmstadt, Germany). Blocks were washed with TE Buffer (10 mM TRIS/HCl, 1 mM EDTA, pH 7.5) and 1 mM phenyl methyl sulfonyl fluoride (PMSF) for protease inactivation followed by washing 3 times with TE. Aliquots of inserts were subjected to digestion for 24 h at 37 °C in 80 µL buffer containing 10× restriction buffer and 20 U XhoI.—2 µL enzyme. After washing with TE, the blocks were inserted in the slots of 1% PFGE agarose (Bio-Rad Laboratories, Hercules, CA, USA).

The apparatus used was a CHEF—DRII (Bio-Rad Laboratories, Hercules, CA, USA), the electrophoretic buffer contained 0.5× TBB (45 mM TRIS, 45 mM Boric acid, 1 mM EDTA, pH 8.3). The electrophoresis parameters were: initial pulse—3 s, final pulse—25 s,

5 v/cm, and running time—24 h. The gels were stained with 200 µg/mL SYBR Green (Thermo Fisher Scientific, Waltham, MA, USA) and examined by UV transillumination.

### 2.5. Metagenome Library Construction, Sequencing and Bioinformatics Analysis

Metagenome library construction and sequencing was done by Macrogen Inc., Republic of Korea. Preparation of the 16S metagenomic sequencing library was performed using a Herculase II Fusion DNA Polymerase Nextera XT Index Kit V2. The sequencing (Illumina platform) was conducted with reading length of 301 bp and FastQC quality control. The assembly results showed that the quality-filtered data contained between 73,209,822 and 82,104,372 total bases, and around 243,222 read counts for each sample. The percentage of Q20 quality reads was 94.52%.

### 2.6. Data Availability

Raw sequencing data were deposited in GenBank of NCBI with the following accession numbers: BioProject ID PRJNA681359, BioSample: SAMN16961356, Sequence Read Archive (SRA) SUB8652830. The partial 16S rRNA gene sequences of strains used in this study are deposited in GenBank with the following accession numbers: MG437344, MG437345, MG437346, MG437357, MG437359, MG437363, MG437367, MG437372, MG437380, MG437386, MG437389, MG437390, MG437352, MG437355, MG438473, MG438475, MG438468, MG438466, MG437360, MG437368, MG437358, MG437375, MG437361, and MG438467.

### 2.7. Liquid Chromatography and Mass Spectrometry (LC-MS) Analysis


Analyses were carried out on a Q Exactive<sup>®</sup> mass analyzer equipped with TurboFlow<sup>®</sup>LC system and APCI atmospheric pressure electrospray ionization module (Thermo Scientific Inc., Waltham, MA, USA) using an iHILIC<sup>®</sup>-Fusion 1.8 µm, 100 Å, 100 × 2.1 mm analytical column (HILICON AB, Sweden) in gradient elution mode. The following mobile phases were used: A—20 mM ammonium acetate in water; B—buffer A/ acetonitrile (1/9 v/v) at a flow rate of 300 µL/min and gradient elution. Full-scan mass spectra over the *m/z* range 67–1000 were acquired in negative ion mode at resolution settings of 140,000. Qualification of the compounds was achieved using top 5N mode of operation of the mass spectrometer at resolution settings of 17,500 and isolation window of 0.8 over the *m/z* range 67–1000. The used value of the collision energy was HCD 30%. All MS parameters were optimized for sensitivity to the target analytes using the instrument control software program. Data acquisition and processing were carried out with the Xcalibur 2.4<sup>®</sup> software package (Thermo Scientific Inc., Waltham, MA, USA). The raw data files were processed by Compound Discoverer<sup>®</sup> 2.0 software (Thermo Scientific Co., USA).

## 3. Results and Discussion

### 3.1. Metagenomes Analysis

Several regions in Bulgaria are known to contain endemic microflora: the high mountains of Rila, Pirin, and Rhodope, and the Thracian valley. The species biodiversity of yoghurt samples collected from these locations was observed after metagenome sequencing (Figure 1, Table S1). As expected, between 68.4% and 99.6% of the total microflora in the examined samples consisted of two species: *Lb. delbrueckii* subsp. *bulgaricus* and *Str. thermophilus*, and the ratio between them varied widely. In Sample 2 (Rila), *Str. thermophilus* was the most abundant species, whereas in Sample 4 (Rhodope), *Lb. bulgaricus* strongly prevailed. Considering the distribution by region, the proto-cooperation between *Lb. bulgaricus* and *Str. thermophilus* existed in all samples, however, the recommended ratio of ~1:1 between strains was observed only in Sample 3 (from the Thracian valley). This ratio in the starter strongly affects yoghurt qualities, such as texture, color, taste, and post-acidification [29,30]. Hence, the artisanal yoghurts are usually characterized by a more pronounced sour taste [10]. Rhodope's yoghurt was the richest source of *Lb. bulgaricus*, and this species is typical for Bulgarian yoghurt and is not as widespread in the neighboring countries. For instance, in a Greek study, the isolation of only one *Lb. delbrueckii* subsp.


*bulgaricus* strain from nine analyzed yoghurts was reported [31]. Comparatively low counts of *Lb. bulgaricus* in Sample 2 could be explained by the repeated transfers of the same starter culture in an unsterile home environment. Under these conditions, bacteriophages could destroy part of the starter [32,33], thus giving rise to the microbial counts of the accompanying microflora.



**Figure 1.** Metagenome study of species abundance in four samples of Bulgarian home-made yoghurt prepared using an artisanal starter. Yoghurt origin: Sample 1 and 2, from villages in Rila and Pirin mountains; Sample 3, from the Thracian Plain; Sample 4, from Rhodope mountain.

According to Siewewerts et al. (2008), the traditional yoghurt culture is composed of *Lb. delbrueckii* subsp. *bulgaricus* and *Str. thermophilus*, and the name “yoghurt” is only allowed for those products that contain starters of these species [34]. Recent metagenome studies devoted to the juxtaposition of yoghurt microbiota and gut microbiome revealed the presence of *Lacticaseibacillus paracasei*, *Lacticaseibacillus rhamnosus*, *Lb. parabuchneri*, *Lb. parafarraginis*, *Lactiplantibacillus plantarum*, *Lactococcus lactis*, *Leuconostoc mesenteroides*, and *Pediococcus acidilactici* in both niches [35,36]. Our results partially confirmed these observations, because among autochthonic lactobacilli in Bulgarian artisanal yoghurts *Lb. rhamnosus* and *Lb. paracasei* were found. In addition, *Lb. helveticus*, *Limosilactobacillus fermentum*, and *Lb. equicursoris* were detected (Table S1). However, whereas *Lb. helveticus* was frequently observed in Bulgarian yoghurts and identified using genetic techniques [37,38], the presence of *Lb. fermentum*, *Lb. paracasei*, and *Lb. rhamnosus* is highly unusual. *Lb. plantarum*, typical of Greek yoghurt, was not found in Bulgarian yoghurt. Although *Lb. equicursoris* belongs to the *Lb. delbrueckii* phylogenetic branch, our study is the first that reports its presence in yoghurts. This species was initially isolated from the feces of a thoroughbred racehorse by Morita et al. (2010) [39] and recently found in kefir [40]. Other accompanying LAB species were *Lc. lactis*, *Lc. garvieae*, *Lc. taiwanensis*, *P. acidilactici*, *Leuc. lactis*, *Leuc. mesenteroides*, and *Leuc. paramesenteroides*. A portion of these species is known to contribute to the sweet, fruity, or vanilla aroma via the production of ethanol, 2-butanone, 2-pentanone, and 2,3-pentanedione [41], and is currently preferred by consumers. Other desirable additives to the starters are also the strains producing extracellular exopolysaccharides and, importantly, those with high proteolytic activity.

Figure 2 and Table S2 present the spoilage species found in the metagenomes. *Acinetobacter* spp. and *Chryseobacterium* spp. were identified as the main spoilage species. These Gram-negative, post-process contaminant bacteria limit the shelf life of milk produced by microfiltration and subsequent pasteurization, and have also been found in yoghurts in Germany, Austria, and Switzerland [42]. The presence of the Gram-negative, obligate anaerobe *Bacteroides gramminisolvans* in two cow samples from different regions (Samples 1 and 3) is indicative of soil contamination during milking and/or the transportation of the raw milk. This species is ubiquitous in barns and rural areas.



**Figure 2.** Metagenome study of species abundance of the contaminant microflora in four samples of Bulgarian home-made yoghurt.

Other microbial species, such as *Moraxella osloensis*, *Alcaligenes aquatilis*, *Str. ruminantium*, and *Str. salivioxodontae*, were found in counts between 0.1% and 0.01%, or less. The low microbial counts of contaminating species (Table S2) indicate that single cells from the environment contaminated the milk, but these species did not multiply due to the antagonistic activity of LAB [43]. Our observations are consistent with the research of other authors reporting the presence of contaminant Gram-negative psychrotrophs in yoghurts, such as *Pseudomonas* sp. and *Klebsiella* sp. [44]. Similarly, Bulgarian yoghurts were contaminated, but with insignificant amounts of *Ps. japonica* and *Citrobacter europaeus*. However, dangerous zoonotic pathogens, such as coliforms, *Salmonella* sp., *Listeria monocytogenes*, *Campylobacter jejuni*, *Yersinia enterocolitica*, *Brucella* sp., *Mycobacterium bovis*, or *Enterobacter sakazakii*, were not found in any of the samples, although their presence in milk has been previously reported [45–47].

### 3.2. LAB Strains Isolation and Identification


Twenty-four LAB strains belonging either to the starter or to the accompanying microflora were isolated and identified by 16S rRNA gene sequencing: *Lb. delbrueckii* subsp. *bulgaricus* (12 strains), *Str. thermophilus* (two strains), *Lb. helveticus* (two strains), *Lb. paracasei* (two strains), *Lb. fermentum* (two strains), *Lb. rhamnosus*, *P. acidilactici*, *Leuc. mesenteroides*, and *Leuc. pseudomesenteroides*. Although present in the sample, some species were not isolated as pure cultures.

To examine the diversity and specificity of *Lb. delbrueckii* subsp. *bulgaricus* strains, three different genotyping approaches were applied: RAPD, MLST, and PFGE.

#### 3.2.1. RAPD Analysis

The analysis of the profile of PCR amplified DNA fragments is shown in Figure 3. A set of five different primers was checked, and the most discriminative profiles were obtained using the combination of primers 1254: 5'CCGCAGCCAA3' (10-mer) and rpd1: 5'CCCGACTGA3' (9-mer).

The results showed that the newly isolated *Lb. delbrueckii* subsp. *bulgaricus* strains are genetically different and also that they differ from the starter cultures of the most popular Bulgarian market yoghurts, i.e., those of the companies Domlyan and Group Danone. However, RAPD was not the method that was able to distinguish all genetically close strains. Two pairs of the isolates did not provide sufficiently good profiles (9, 25, 55, and 63). To prove their genetic uniqueness, the use of other molecular techniques was needed.



**Figure 3.** Randomly amplified polymorphic DNA–PCR (RAPD) analysis of 12 *Lb. delbrueckii* subsp. *bulgaricus* isolates and two commercial *Lb. delbrueckii* subsp. *bulgaricus* strains. The strain 212 derives from the yoghurt “Domlyan”, and 222 from the yoghurt “Na baba” of Danone.

### 3.2.2. MLST Analysis


MLST is generally regarded as a bacterial identification method that depends on the genomic properties or experimental design (such as the kinds of genes analyzed), and provides a higher resolution than RAPD. MLST uses automated DNA sequencing to characterize the alleles present at different housekeeping gene loci. Because it is based on nucleotide sequences, it is highly discriminatory and provides unambiguous results. MLST analyses were performed with four housekeeping genes, shown at Figure 4: *rpoB*, *groEL*, *pheS*, and *pyrG*, encoding the  $\beta$ -subunit of bacterial RNA polymerase, the chaperonin family of molecular chaperones, the  $\alpha$ -subunit of phenylalanine tRNA synthetase, and CTP synthase, respectively. For the strains examined, parts of these genes were amplified and sequenced.

By sequencing of the genes and sequence comparison using ClustalW, *Lb. delbrueckii* subsp. *bulgaricus* strains were clustered into different branches according to the used housekeeping gene. There were ten different alleles for *rpoB*, eleven for *groEL*, eleven for *pheS*, and twelve for *pyrG*. Table 2 estimates the evolutionary divergence of combined sequences, and the number of base substitutions per site from between sequences is shown. It could be deduced that all strains are diverse and each strain is genetically different from the others. Figure 5 shows the strains’ evolutionary relationships.


**Table 2.** Estimates of evolutionary divergence between combined sequences. The number of base substitutions per site between sequences is shown. Analyses were conducted using the maximum composite likelihood model. The analysis involved 14 combined nucleotide sequences. There were a total of 2474 positions in the final dataset.

	1	1	2	9	25	27	34	38	44	55	63	69	71	212
2		0.004												
9		0.003	0.004											
25		0.003	0.004	0.001										
27		0.008	0.009	0.010	0.010									
34		0.007	0.008	0.009	0.009	0.002								
38		0.003	0.004	0.001	0.001	0.009	0.008							
44		0.002	0.004	0.002	0.002	0.009	0.007	0.002						
55		0.008	0.008	0.009	0.009	0.001	0.001	0.009	0.009					
63		0.008	0.009	0.010	0.010	0.002	0.001	0.009	0.008	0.001				
69		0.009	0.009	0.010	0.010	0.002	0.001	0.009	0.009	0.002	0.001			
71		0.008	0.009	0.010	0.010	0.002	0.001	0.009	0.008	0.001	0.001	0.001		
212		0.003	0.004	0.001	0.001	0.010	0.008	0.001	0.001	0.009	0.009	0.009	0.009	
222		0.003	0.004	0.001	0.001	0.009	0.009	0.001	0.001	0.009	0.009	0.010	0.009	0.001






**Figure 4.** Phylogenetic trees of *Lb. delbrueckii* subsp. *bulgaricus* isolates based on the sequences of four housekeeping genes: *rpoB* (A), *groEL* (B), *pheS* (C), and *pyrG* (D). Evolutionary analyses were conducted using MEGA6 software. The evolutionary history was inferred using the neighbor-joining method. The evolutionary distances were computed using the maximum composite likelihood method and are in the units of the number of base substitutions per site. The analysis involved 14 nucleotide sequences. Codon positions included were 1st + 2nd + 3rd + Noncoding. All positions containing gaps and missing data were eliminated.



**Figure 5.** Evolutionary relationships between newly isolated *Lb. delbrueckii* subsp. *bulgaricus* strains. The evolutionary history was inferred using the minimum evolution method. The optimal tree with the sum of branch length = 0.01438984 is shown. The evolutionary distances were computed using the maximum composite likelihood method and are in the units of the number of base substitutions per site. The ME tree was searched using the Close-Neighbor-Interchange (CNI) algorithm at a search level of 1. All positions containing gaps and missing data were eliminated by MEGA6.

### 3.2.3. PFGE

The PFGE method is strain-specific and provides precise insight into the isolated strains (Figure 6).



**Figure 6.** Macro-restriction profiles of *Xho* I-digested DNA of *Lb. delbrueckii* subsp. *bulgaricus* strains.


Although several enzymes were engaged (*Not*I, *Apa*I), the most highly discriminative profiles were obtained by the digest of chromosomal DNA using the restriction enzyme *Xho*I. The results presented in Figure 6 display the affiliation of the strains to 14 different groups, i.e., they are clustered separately because they are genetically different, indicating the large genetic diversity among *Lb. delbrueckii* subsp. *bulgaricus* strains isolated from home-made yoghurt.

### 3.3. Bioactive Metabolites Produced by the Accompanying Microflora of Bulgarian Yoghurts

#### 3.3.1. Amino Acids and Their Derivatives


Aiming to elucidate the ability of the accompanying LAB strains to produce valuable metabolites, single-strains fermented yoghurts were subjected to LC-MS analysis (Table S3). Interestingly, *Lb. delbrueckii* subsp. *bulgaricus* is an auxotroph for several amino acids (for instance, branched-chained amino acids) and the means of obtaining them is to use those formed by the accompanying microflora in the course of biosynthetic processes [48]. Considering the available data, reports of free amino acids accumulation by LAB are scarce. However, a *Lb. helveticus* strain isolated from kefir reached 53.38 mg free amino acids per 100 g product [49]. In our study, the amino acid that was accumulated in the largest amount was arginine, and *Lb. helveticus* 16 and 22 produced it the most compared to the other LAB strains (Figure 7). Arginine is an essential amino acid and has been suggested as a possible prebiotic [50]. L-arginine and its precursor L-citrulline are widely used in the treatment of cardiovascular diseases associated with endothelial dysfunction, such as hypertension, heart failure, atherosclerosis, diabetic vascular disease, and ischemia-reperfusion injury [51]. The ability of the studied LAB to accumulate L-citrulline is highly significant. Citrulline is known as a laxative and as a blood-plasma marker for human irritable bowel syndrome. Supplementation of citrulline with *Lb. helveticus* can improve

intestinal barrier functions [52]. L-arginine and L-citrulline oral intake improve erection hardness in men; both amino acids are used as a treatment of erectile dysfunction [53,54].



**Figure 7.** Production of amino acids arginine, lysine, and histidine by analyzed LAB. The LC-MS analyses were carried out using a Q Exactive<sup>®</sup> mass analyzer equipped with TurboFlow<sup>®</sup> LC system and APCI atmospheric pressure electrospray ionization module. Data acquisition and processing were carried out with the Xcalibur 2.4<sup>®</sup> software package.

*Lb. fermentum* 28, *Lb. paracasei* 25 and 47, and *P. acidilactici* 27 accumulated lysine and tryptophan (Figure 8), and LAB enzymes involved in these pathways may have future industrial applications [55,56]. Almost all strains produced histidine, but the potentially toxic biogenic amines deriving from its degradation were not detected.




**Figure 8.** Production of indole-3-propionic acid, L-tryptophan, and L-citrulline by analyzed LAB. The LC-MS analyses were carried out using a Q Exactive<sup>®</sup> mass analyzer equipped with a TurboFlow<sup>®</sup> LC system and APCI atmospheric pressure electrospray ionization module. Data acquisition and processing were carried out with the Xcalibur 2.4<sup>®</sup> software package.


### 3.3.2. Antimicrobial Peptides

One of the most important bioactive metabolites produced by the accompanying microflora is indole-3-propionic acid (IPA) because it is being studied as a therapeutic in the treatment of many neurodegenerative disorders. According to Mimori et al. (2019), IPA acts as “chemical chaperone”, suppressing stress-induced cell death of neurons and ceasing the aggregation of the denatured proteins in vitro [57]. IPA, and its esters or salts, prevent the cytotoxic effects of amyloid beta-protein on cells and can be used as a

treatment for fibrillogenic diseases. This inhibition of beta-amyloid fibril formation implies novel therapy for Alzheimer's disease [58,59]. IPA is an even more potent scavenger of hydroxyl radicals than melatonin, moreover, it does so without generating subsequent reactive and pro-oxidant intermediate compounds. In 2017, elevated concentrations of IPA in human blood plasma were found to correlate with a lower risk of type 2 diabetes [60]. Because the phenomenon was also connected with the higher consumption of fiber-rich foods (prebiotics), this effect was assigned to IPA, which is endogenously produced by the human microbiota. Other beneficial metabolites produced by LAB of the accompanying microflora were two cyclic peptides: cyclo(phenylalanyl-prolyl) and cyclo(leucylprolyl) (Figures 9 and 10).



**Figure 9.** Two- (2D), and three-dimensional (3D) structural formulas of antimicrobial compounds produced by accompanying LAB in yoghurt. (A,B) Cyclo(phenylalanyl-prolyl), Mw 244 Da; (C,D) Cyclo(leucylprolyl), 210 Da. The models were obtained from the free chemical structure database of the Royal Society of Chemistry *ChemSpider* [61].



**Figure 10.** Production of antimicrobial peptides by LAB strains. The LC-MS analyses were carried out using a Q Exactive<sup>®</sup> mass analyzer equipped with a TurboFlow<sup>®</sup> LC system and APCI atmospheric pressure electrospray ionization module. Data acquisition and processing were carried out with the Xcalibur 2.4<sup>®</sup> software package.

Cyclo(phenylalanyl-prolyl) has broad-spectrum antimicrobial activity, including antibacterial, antifungal, antiviral, anti-settlement, antiprotozoal, antiparasitic, and antitumor properties, and has a radioprotective effect [62]. It is produced in significant amounts by *Leuc. pseudomesenteroides* 24, *Str. thermophilus* 21, *Lb. fermentum* 28, and *P. acidilactici* 27.

Cyclo(leucylprolyl), also known as the antibiotic Gancidin W, is another antimicrobial compound produced by the concomitant LAB. It is synthesized in higher amounts by *Lb. fermentum*, *Leuc. mesenteroides*, and *P. acidilactici*. It is a bioactive natural product, and an alkaloid that provides the bitter component of sake and contributes to the flavor of the beer.

Recently, the anticancer effect of Gandicin W against breast cancer was demonstrated [63]. Until now, both antimicrobial compounds have been found only in marine bacteria (named drugs from the ocean) [64]. Our investigation reports for the first time their presence in yoghurt.

#### 4. Conclusions

Metagenomic study of authentic Bulgarian yoghurts revealed the presence of “classical” starter strains *Lb. delbrueckii* subsp. *bulgaricus* and *Str. thermophilus*, accompanied by many LAB and spoilage genera. RAPD, MLST, and PFGE analyses showed that the *Lb. delbrueckii* subsp. *bulgaricus* strains in homemade yoghurts are different, authentic, and do not participate in commercial starters. The formation of unique metabolites belonging to the class of amino acids is a significant novelty of the present work. The accumulation of arginine, lysine, and histidine by the biosynthetic properties of the accompanying microflora is beneficial for the propagation of the starter cultures *Lb. bulgaricus*, but also for the consumer of the yoghurt. Metabolites that are valuable for human health, such as indole-3-propionic acid and L-citrulline, were detected. This identification could address the issue of the “magic” effects of Bulgarian yoghurt on the elderly, who retain clear minds and sexual activity to an advanced age. The strains producing the antimicrobial peptides cyclo(phenylalanyl-prolyl) and cyclo(leucyloprolyl) may be included in future starter formulations. In summary, the specific natural and climatic conditions in Bulgaria contributed to the evolution of yoghurt starter cultures with unique features. The comprehensive examination of the current status of the authentic Bulgarian yoghurt microbial diversity showed that homemade yoghurts, produced in small farms in Bulgarian mountainous areas, are the source of LAB with health-promoting effects.

**Supplementary Materials:** The following are available online at <https://www.mdpi.com/2227-9717/9/1/114/s1>, Table S1: Metagenome study of Lactic acid bacteria—species abundance in four samples of Bulgarian home-made yoghurt prepared using an artisanal starter. Yoghurt origin: Sample 1 and 2, from villages in Rila and Pirin mountains, Sample 3, from the Thracian Plain, Sample 4, from Rhodope mountain; Table S2: Metagenome study of contaminant microflora species abundance in four samples of Bulgarian home-made yoghurt prepared using an artisanal starter. Yoghurt origin: Sample 1 and 2, from villages in Rila and Pirin mountains, Sample 3, from the Thracian Plain, Sample 4, from Rhodope mountain; Table S3: LC-MS (Liquid chromatography/Mass spectrometry) analysis of the valuable metabolites produced by the accompanying LAB strains in single-strain fermented yoghurts. LC-MS analysis was carried out on Q Exactive® mass analyzer equipped with TurboFlow® LC system and APCI atmospheric pressure electrospray ionization module. Data acquisition and processing were carried out with Xcalibur 2.4® software package.

**Author Contributions:** Conceptualization, P.P. and K.P.; methodology, I.I., V.L., I.H.; writing—original draft preparation, I.I.; writing—review and editing, P.P., Z.W., Z.L. All authors have read and agreed to the published version of the manuscript.

**Funding:** Metagenomics experiments were funded by the Bulgarian Ministry of Education and Science with the financial support by the “Healthy Foods for a Strong Bio-Economy and Quality of Life” National Research Program approved by DCM # 577/17.08.2018. The study was financially supported also by the company Bright Dairy & Food Co. Ltd, China, grant number 1/18.06.2015.

**Institutional Review Board Statement:** Not applicable.

**Informed Consent Statement:** Not applicable.

**Data Availability Statement:** Publicly available datasets were analyzed in this study. This data can be found here: <https://www.ncbi.nlm.nih.gov/bioproject/?term=PRJNA681359>; [https://www.ncbi.nlm.nih.gov/biosample?Db=biosample&DbFrom=bioproject&Cmd=Link&LinkName=bioproject\\_biosample&LinkReadableName=BioSample&ordinalpos=1&IdsFromResult=681359](https://www.ncbi.nlm.nih.gov/biosample?Db=biosample&DbFrom=bioproject&Cmd=Link&LinkName=bioproject_biosample&LinkReadableName=BioSample&ordinalpos=1&IdsFromResult=681359).

**Conflicts of Interest:** The authors declare no conflict of interest. The funders had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript, or in the decision to publish the results.

## References

1. Fernandez, M.A.; Panahi, S.; Daniel, N.; Tremblay, A.; Marette, A. Yogurt and cardiometabolic diseases: A critical review of potential mechanisms. *Adv. Nutr.* **2017**, *8*, 812–829. [[CrossRef](#)] [[PubMed](#)]
2. Berthold-Pluta, A.M.; Pluta, A.S.; Garbowska, M.; Stasiak-Róžańska, L. Exopolysaccharide-producing Lactic Acid Bacteria—Health-promoting properties and application in the dairy industry. *Adv. Microbiol.* **2019**, *58*, 191–204. [[CrossRef](#)]
3. Popović, N.; Brdarić, E.; Đokić, J.; Dinić, M.; Veljović, K.; Golić, N.; Terzić-Vidojević, A. Yogurt produced by novel natural starter cultures improves gut epithelial barrier in vitro. *Microorganisms* **2020**, *8*, 1586. [[CrossRef](#)] [[PubMed](#)]
4. Julio, P.D.; Carolina, G.L.; Luis, F.; Angel, G. Modulation of immunity and inflammatory gene expression in the gut, in inflammatory diseases of the gut and in the liver by probiotics. *World J. Gastroenterol.* **2014**, *20*, 15632–15649.
5. Mousavi, S.N.; Somayeh, S.; Asbaghi, O. Effect of daily probiotic yogurt consumption on inflammation: A systematic review and meta-analysis of randomized Controlled Clinical trials. *Obes. Med.* **2020**, *18*, 100221. [[CrossRef](#)]
6. De Filippis, F.; Parente, E.; Ercolini, D. Metagenomics insights into food fermentations. *Microb. Biotechnol.* **2016**, *10*, 91–102. [[CrossRef](#)] [[PubMed](#)]
7. Hao, P.; Zheng, H.; Yu, Y.; Ding, G.; Gu, W.; Chen, S.; Yu, Z.; Ren, S.; Oda, M.; Konno, T.; et al. Complete sequencing and pan-genomic analysis of *Lactobacillus delbrueckii* subsp. *bulgaricus* reveal its genetic basis for industrial yogurt production. *PLoS ONE* **2011**, *6*, e15964. [[CrossRef](#)]
8. Liu, M.; Siezen, R.J.; Nauta, A. In Silico prediction of horizontal gene transfer events in *Lactobacillus bulgaricus* and *Streptococcus thermophilus* reveals protocoeoperation in yogurt manufacturing. *Appl. Environ. Microbiol.* **2009**, *75*, 4120–4129. [[CrossRef](#)] [[PubMed](#)]
9. Beshkova, D.; Simova, E.; Frengova, G.; Simov, Z. Production of flavour compounds by yogurt starter cultures. *J. Ind. Microbiol. Biotechnol.* **1998**, *20*, 180–186. [[CrossRef](#)]
10. Velikova, P.; Petrov, K.; Lozanov, V.; Tsvetanova, F.; Stoyanov, A.; Wu, Z.; Liu, Z.; Petrova, P. Microbial diversity and health-promoting properties of the traditional Bulgarian yogurt. *Biotechnol. Biotechnol. Equip.* **2018**, *32*, 1205–1217. [[CrossRef](#)]
11. Suzuki, Y.; Ikeda, K.; Sakuma, K.; Kawai, S.; Sawaki, K.; Asahara, T.; Takahashi, T.; Tsuji, H.; Nomoto, K.; Nagpal, R.; et al. Association between yogurt consumption and intestinal microbiota in healthy young adults differs by host gender. *Front. Microbiol.* **2017**, *8*, 847. [[CrossRef](#)] [[PubMed](#)]
12. Boyanova, L.; Stephanova-Kondratenko, M.; Mitov, I. Anti-*Helicobacter pylori* activity of *Lactobacillus delbrueckii* subsp. *bulgaricus* strains: Preliminary report. *Lett. Appl. Microbiol.* **2009**, *48*, 579–584. [[CrossRef](#)] [[PubMed](#)]
13. Boyanova, L.; Gergova, G.; Markovska, R.; Yordanov, D.; Mitov, I. Bacteriocin-like inhibitory activities of seven *Lactobacillus delbrueckii* subsp. *bulgaricus* strains against antibiotic susceptible and resistant *Helicobacter pylori* strains. *Lett. Appl. Microbiol.* **2017**, *65*, 469–474. [[CrossRef](#)] [[PubMed](#)]
14. Stefanova, T.; Urshev, Z.; Dimitrov, Z.; Fatchikova, N.; Minkova, S. Species affiliation of dairy lactobacilli with angiotensin converting enzyme inhibitory activity. *Biotechnol. Biotechnol. Equip.* **2009**, *23*, 1250–1254. [[CrossRef](#)]
15. Cocolin, L.; Ercolini, D. Zooming into food-associated microbial consortia: A “cultural” evolution. *Curr. Opin. Food Sci.* **2015**, *2*, 43–50. [[CrossRef](#)]
16. Lee, A.; Lee, Y.J.; Yoo, H.J.; Kim, M.; Chang, Y.; Lee, D.S.; Lee, J.H. Consumption of dairy yogurt containing *Lactobacillus paracasei* ssp. *paracasei*, *Bifidobacterium animalis* ssp. *lactis* and heat-treated *Lactobacillus plantarum* improves immune function including natural killer cell activity. *Nutrients* **2017**, *9*, 558. [[CrossRef](#)]
17. Sánchez, E.; Nieto, J.; Vidal, S.; Santiago, A.; Martínez, X.; Sancho, F.J.; Sancho-Bru, P.; Mirelis, B.; Corominola, H.; Juárez, C.; et al. Fermented milk containing *Lactobacillus paracasei* subsp. *paracasei* CNCM I-1518 reduces bacterial translocation in rats treated with carbon tetrachloride. *Sci. Rep.* **2017**, *7*, 45712.
18. Wang, J.; Li, C.; Xue, J.; Yang, J.; Zhang, Q.; Zhang, H.; Chen, Y. Fermentation characteristics and angiotensin I-converting enzyme-inhibitory activity of *Lactobacillus helveticus* isolate H9 in cow milk, soy milk, and mare milk. *J. Dairy Sci.* **2015**, *98*, 3655–3664. [[CrossRef](#)]
19. Zhou, T.; Huo, R.; Kwok, L.Y.; Li, C.; Ma, Y.; Mi, Z.; Chen, Y. Effects of applying *Lactobacillus helveticus* H9 as adjunct starter culture in yogurt fermentation and storage. *J. Dairy Sci.* **2019**, *102*, 223–235. [[CrossRef](#)]
20. Cavalheiro, G.F.; Baptista, D.P.; Galli, B.D.; Negrão, F.; Eberlin, M.N.; Gigante, M.L. High protein yogurt with addition of *Lactobacillus helveticus*: Peptide profile and angiotensin-converting enzyme ACE-inhibitory activity. *Food Chem.* **2020**, *333*, 127482. [[CrossRef](#)]
21. Cameron, D.; Hock, Q.S.; Kadim, M.; Mohan, N.; Ryoo, E.; Sandhu, B.; Yamashiro, Y.; Jie, C.; Hoekstra, H.; Guarino, A. Probiotics for gastrointestinal disorders: Proposed recommendations for children of the Asia-Pacific region. *World J. Gastroenterol.* **2017**, *23*, 7952–7964. [[CrossRef](#)] [[PubMed](#)]
22. Szajewska, H.; Guarino, A.; Hojsak, I.; Indrio, F.; Kolacek, S.; Shamir, R.; Vandenplas, Y.; Weizman, Z. Use of probiotics for management of acute gastroenteritis: A position paper by the ESPGHAN Working Group for Probiotics and Prebiotics. *J. Ped. Gastroenterol. Nutr.* **2014**, *58*, 531–539. [[CrossRef](#)] [[PubMed](#)]
23. Szajewska, H.; Horvath, A. *Lactobacillus rhamnosus* GG in the primary prevention of eczema in children: A systematic review and meta-analysis. *Nutrients* **2018**, *10*, 1319. [[CrossRef](#)] [[PubMed](#)]
24. Grigoroff, S. Étude sur une lait fermentée comestible. Le “Kissélo mléko” de Bulgarie. *Rev. Méd. Suisse Romande* **1905**, *25*, 714–720.

25. Lick, S.; Keller, M.; Bockelmann, M.; Heller, K.J. Optimized DNA extraction method for starter cultures from yoghurt. *Milchwissenschaft* **1996**, *51*, 183–186.
26. Cebeci, A.; Gurakan, C. Comparative typing of *L. delbrueckii* subsp. *bulgaricus* strains using multilocus sequence typing and RAPD-PCR. *Eur. Food. Res. Technol.* **2011**, *233*, 377–385. [[CrossRef](#)]
27. Naser, S.M.; Thompson, F.L.; Hoste, B.; Gevers, D.; Dawyndt, P.; Vancanneyt, M.; Swings, J. Application of multilocus sequence analysis (MLSA) for rapid identification of *Enterococcus* species based on *rpoA* and *pheS* genes. *Microbiology* **2005**, *151*, 2141–2150. [[CrossRef](#)]
28. Michaylova, M.; Minkova, S.; Kimura, K.; Sasaki, T.; Isawa, K. Isolation and characterization of *Lactobacillus delbrueckii* ssp. *bulgaricus* and *Streptococcus thermophilus* from plants in Bulgaria. *FEMS Microbiol. Lett.* **2007**, *269*, 160–169. [[CrossRef](#)]
29. Undugoda, L.J.S.; Nilmini, A.H.L. Effect of lactic acid microbial ratio of yoghurt starter culture in yoghurt fermentation and reduction of post acidification. *J. Food Ind. Microbiol.* **2019**, *5*, 1.
30. Surono, I.S.; Hosono, A. Fermented milks. Types and Standards of Identity. In *Encyclopedia of Dairy Sciences*, 2nd ed.; Fuquay, J.W., Fox, P.F., McSweeney, P.L.H., Eds.; Academic Press: New York, NY, USA, 2011; pp. 470–476.
31. Georgalaki, M.; Zoumpopoulou, G.; Mavrogonatou, E.; Van Driessche, G.; Alexandraki, V.; Anastasiou, R.; Papadelli, M.; Kazou, M.; Manolopoulou, E.; Kletsas, D.; et al. Evaluation of the antihypertensive angiotensin-converting enzyme inhibitory (ACE-I) activity and other probiotic properties of lactic acid bacteria isolated from traditional Greek dairy products. *Int. Dairy J.* **2017**, *75*, 10–21. [[CrossRef](#)]
32. Urshev, Z.L.; Pashova-Baltova, K.N.; Dimitrov, Z.P. Tracing *Streptococcus thermophilus* strains in three-component yogurt starters. *World J. Microbiol. Biotechnol.* **2006**, *22*, 1223–1228. [[CrossRef](#)]
33. Ma, C.; Chen, Z.; Gong, G.; Huang, L.; Li, S.; Ma, A. Starter culture design to overcome phage infection during yogurt fermentation. *Food Sci. Biotechnol.* **2015**, *24*, 521–527. [[CrossRef](#)]
34. Sieuwerts, S.; De Bok, F.A.; Hugenholtz, J.; Van Hylckama Vlieg, J.E. Unraveling microbial interactions in food fermentations: From classical to genomics approaches. *Appl. Environ. Microbiol.* **2008**, *74*, 4997–5007. [[CrossRef](#)]
35. Alvarez, A.S.; Tap, J.; Chambaud, I.; Cools-Portier, S.; Quinquis, L.; Bourlioux, P.; Marteau, P.; Guillemard, E.; Schrezenmeir, J.; Derrien, M. Safety and functional enrichment of gut microbiome in healthy subjects consuming a multi-strain fermented milk product: A randomised controlled trial. *Sci. Rep.* **2020**, *10*, 15974. [[CrossRef](#)] [[PubMed](#)]
36. Roder, T.; Wüthrich, D.; Bär, C.; Sattari, Z.; von Ah, U.; Ronchi, F.; Macpherson, A.J.; Ganai-Vonarburg, S.C.; Bruggmann, R.; Vergères, G. In Silico Comparison Shows that the Pan-Genome of a Dairy-Related Bacterial Culture Collection Covers Most Reactions Annotated to Human Microbiomes. *Microorganisms* **2020**, *8*, 966. [[CrossRef](#)]
37. Miteva, V.I.; Abadjieva, A.N.; Stefanova, T.T. M13 DNA fingerprinting, a new tool for classification and identification of *Lactobacillus* spp. *J. Appl. Bacteriol.* **1992**, *73*, 349–354. [[CrossRef](#)]
38. Dimitrov, Z.; Michaylova, M.; Mincova, S. Characterization of *Lactobacillus helveticus* strains isolated from Bulgarian yogurt, cheese, plants and human faecal samples by sodium dodecyl sulfate polyacrylamide gel electrophoresis of cell-wall proteins, ribotyping, and pulsed-field gel fingerprinting. *Int. Dairy J.* **2005**, *15*, 998–1005. [[CrossRef](#)]
39. Morita, H.; Shimazu, M.; Shiono, H.; Toh, H.; Nakajima, F.; Akita, H.; Takagi, M.; Takami, H.; Murakami, M.; Masaoka, T.; et al. *Lactobacillus equicursoris* sp. nov., isolated from the faeces of a thoroughbred racehorse. *Int. J. Syst. Evol. Microbiol.* **2010**, *60*, 109–112. [[CrossRef](#)]
40. Beleza do Campo, K. Available online: <https://www.belezadocampo.com.br> (accessed on 10 December 2020).
41. Chen, C.; Zhao, S.; Hao, G.; Yu, H.; Tian, H.; Zhao, G. Role of lactic acid bacteria on the yogurt flavour: A review. *Int. J. Food Prop.* **2017**, *20*, S316–S330. [[CrossRef](#)]
42. Verena, S.J.; Schmidt, V.K.; Kulozik, U.; Scherer, S.; Wenning, M. Microbial biodiversity, quality and shelf life of microfiltered and pasteurized extended shelf life (ESL) milk from Germany, Austria and Switzerland. *Int. J. Food Microbiol.* **2012**, *154*, 1–9.
43. Kagkli, D.M.; Vancanneyt, M.; Hill, C.; Vandamme, P.; Cogan, T.M. *Enterococcus* and *Lactobacillus* contamination of raw milk in a farm dairy environment. *Int. J. Food Microbiol.* **2007**, *114*, 243–251. [[CrossRef](#)] [[PubMed](#)]
44. Agu, K.; Archibong, E.; Anekwe, D.; Ago, C.; Okafor, A.; Awah, N. Assessment of bacteria present in yoghurt sold on Awka Metropolis. *Sch. J. Appl. Med. Sci.* **2014**, *2*, 3071–3075.
45. Pal, M.; Awel, H. Public health significance of *Listeria monocytogenes* in milk and milk products. *J. Vet. Public Health* **2014**, *12*, 1–5.
46. Addis, M.F.; Tanca, A.; Uzzau, S.; Oikonomou, G.; Bicalho, R.C.; Moroni, P. The bovine milk microbiota: Insights and perspectives from -omics studies. *Mol. Biosyst.* **2016**, *12*, 2359–2372. [[CrossRef](#)] [[PubMed](#)]
47. Lal, M.; Kaur, H.; Gupta, L.K.; Sood, N.K. Isolation of *Yersinia enterocolitica* from raw milk and pork in Ludhiana. *Ind. J. Pathol. Microbiol.* **2005**, *48*, 286–287.
48. Lee, K.; Lee, J.; Kim, Y.H.; Moon, S.H.; Park, Y.H. Unique properties of four *Lactobacilli* in amino acid production and symbiotic mixed culture for lactic acid biosynthesis. *Curr. Microbiol.* **2001**, *43*, 383–390. [[CrossRef](#)] [[PubMed](#)]
49. Simova, E.; Simov, Z.; Beshkova, D.; Frengova, G.; Dimitrov, Z.; Spasov, Z. Amino acid profiles of lactic acid bacteria, isolated from kefir grains and kefir starter made from them. *Int. J. Food Microbiol.* **2006**, *107*, 112–123. [[CrossRef](#)]
50. Nascimento, M.M. Potential uses of arginine in dentistry. *Adv. Dent. Res.* **2018**, *29*, 98–103. [[CrossRef](#)]
51. Romero, M.J.; Platt, D.H.; Caldwell, R.B.; Caldwell, R.W. Therapeutic use of citrulline in cardiovascular disease. *Cardiovasc. Drug Rev.* **2006**, *24*, 275–290. [[CrossRef](#)]

52. Ho, S.W.; El-Nezami, H.; Shah, N.P. Effects of supplementation of citrulline and *Lactobacillus helveticus* ASCC 511 on intestinal epithelial cell integrity. *J. Funct. Foods* **2020**, *64*, 103571. [[CrossRef](#)]
53. Stanislavov, R.; Nikolova, V. Treatment of erectile dysfunction with pycnogenol and L-arginine. *J. Sex Marital Ther.* **2003**, *29*, 207–213. [[CrossRef](#)] [[PubMed](#)]
54. Cormio, L.; De Siati, M.; Lorusso, F.; Selvaggio, O.; Mirabella, L.; Sanguedolce, F.; Carrieri, G. Oral L-citrulline supplementation improves erection hardness in men with mild erectile dysfunction. *Urology* **2011**, *77*, 119–122. [[CrossRef](#)] [[PubMed](#)]
55. Wang, X.; Yang, L.; Cao, W.; Ying, H.; Chen, K.; Ouyang, P. Efficient production of enantiopure d-lysine from l-lysine by a two-enzyme cascade system. *Catalysts* **2016**, *6*, 168. [[CrossRef](#)]
56. Lim, Y.H.; Foo, H.L.; Loh, T.C.; Mohamad, R.; Abdul-Rahim, R. Rapid evaluation and optimization of medium components governing tryptophan production by *Pediococcus acidilactici* tp-6 isolated from Malaysian food via statistical approaches. *Molecules* **2020**, *25*, 779. [[CrossRef](#)]
57. Mimori, S.; Kawada, K.; Saito, R.; Takahashi, M.; Mizoi, K.; Okuma, Y.; Hosokawa, M.; Kanzaki, T. Indole-3-propionic acid has chemical chaperone activity and suppresses endoplasmic reticulum stress-induced neuronal cell death. *Biochem. Biophys. Res. Commun.* **2019**, *517*, 623–628. [[CrossRef](#)]
58. Pappolla, M. Indole-3-propionic Acids, Salts and Esters Thereof Used as Medicaments. U.S. Patent WO1999042102A1, 26 August 1999.
59. Zhang, L.S.; Davies, S.S. Microbial metabolism of dietary components to bioactive metabolites: Opportunities for new therapeutic interventions. *Genome Med.* **2016**, *8*, 46. [[CrossRef](#)]
60. De Mello, V.D.; Paananen, J.; Lindström, J.; Lankinen, M.A.; Shi, L.; Kuusisto, J.; Bergdahl, I.A. Indolepropionic acid and novel lipid metabolites are associated with a lower risk of type 2 diabetes in the Finnish diabetes prevention study. *Sci. Rep.* **2017**, *7*, 46337. [[CrossRef](#)]
61. Chemspider: Search and Share Chemistry, a Site of Royal Society of Chemistry. Available online: <http://www.chemspider.com> (accessed on 6 December 2020).
62. Lee, K.H.; Rhee, K.H. Radioprotective effect of cyclo(l-phenylalanyl-l-prolyl) on irradiated rat lung. *J. Microbiol. Biotechnol.* **2008**, *18*, 369–376.
63. Ravi, L.; Ragunathan, A.; Krishnan, K. Marine *Streptomyces paradoxus* VITALK03 derived gancidin W mediated cytotoxicity through Ras-Raf-MEK-ERK signalling pathway. *Ind. J. Biotechnol.* **2017**, *16*, 164–175.
64. Nweze, J.A.; Mbaoji, F.N.; Huang, G.; Li, Y.; Yang, L.; Zhang, Y.; Huang, S.; Pan, L.; Yang, D. Antibiotics development and the potentials of marine-derived compounds to stem the tide of multidrug-resistant pathogenic bacteria, fungi, and protozoa. *Mar. Drugs* **2020**, *18*, 145. [[CrossRef](#)]