

Article

Supplementary materials: Hydrodeoxygenation of Guaiacol over Pd–Co and Pd–Fe Catalysts: Deactivation and Regeneration

Nga Tran ^{1,2,*}, Yoshimitsu Uemura ^{1,*}, Thanh Trinh ³ and Anita Ramli ^{1,4}

¹ Centre for Biofuel and Biochemical Research, Department of Chemical Engineering, Universiti Teknologi PETRONAS, Seri Iskandar, Perak, 32610, Malaysia

² Key Laboratory of Chemical Engineering and Petroleum Processing, Ho Chi Minh City University of Technology (HCMUT), 268 Ly Thuong Kiet Street, 72506 District 10, Ho Chi Minh City, Vietnam

³ Faculty of Food Science and Technology, HCMC University of Food Industry (HUFI), 140 Le Trong Tan, 72009 Tan Phu District, Ho Chi Minh City, Vietnam; onalone2000@gmail.com

⁴ Department of Fundamental & Applied Science, Universiti Teknologi PETRONAS, 32610 Seri Iskandar, Perak, Malaysia; anita_ramli@utp.edu.my

* Correspondence: Tonga.tnt@gmail.com (N.T.); YUemura.my@gmail.com (Y.U.); Tel.: (+84)985516524 (N.T.); (+81)9025099726 (Y.U.)




Figure S1. De-convoluted Si 2p, Al 2p and Pd 3d XPS spectra of fresh, used and regenerated Pd-Co/Al-MCM-41 catalysts.




Figure S2. De-convoluted Si 2p, Al 2p and Pd 3d XPS spectra of fresh, used and regenerated Pd-Fe/Al-MCM-41 catalysts.