

Supplementary Material

Antimicrobial Steroids from Poisonous Mushroom *Gymnopilus orientispectabilis* and Their Molecular Docking Studies

Bowon Jung ^{1,†}, Eun Jin Heo ^{1,†}, Dieu Linh Nguyen ^{2,3,4}, Ui Joung Youn ^{4,5}, Ki Hyun Kim ⁶, Boram Son ^{7,*} and Seulah Lee ^{1,*}

¹ Department of Oriental Medicine Biotechnology, Graduate School of Biotechnology, Kyung Hee University, Yongin 17104, Republic of Korea

² Center for Molecular and Translational Human Infectious Diseases Research, Houston Methodist Research Institute, Houston, TX 77030, USA

³ Department of Pathology and Genomic Medicine, Houston Methodist Hospital, Houston, TX 77030, USA

⁴ Division of Life Sciences, Korea Polar Research Institute, Incheon 21990, Republic of Korea

⁵ Department of Polar Sciences, University of Science and Technology, Incheon 21990, Republic of Korea

⁶ School of Pharmacy, Sungkyunkwan University, Suwon 16419, Republic of Korea



⁷ Department of Bio & Fermentation Convergence Technology, Kookmin University, Seoul 02707, Republic of Korea

* Correspondence: boramson@kookmin.ac.kr (B.S.) lee.seulah@khu.ac.kr (S.L.)

† These authors contributed equally to this work.

Table of Contents

Figure S1. ^1H NMR spectrum (600 MHz, CDCl_3) of 1	3
Figure S2. ^{13}C NMR spectrum (150 MHz, CDCl_3) of 1	3
Figure S3. ^1H NMR spectrum (600 MHz, CDCl_3) of 2	4
Figure S4. ^{13}C NMR spectrum (150 MHz, CDCl_3) of 2	4
Figure S5. ^1H NMR spectrum (600 MHz, CDCl_3) of 3	5






Figure S3. ^1H NMR spectrum (600 MHz, CDCl_3) of **2**.




Figure S4. ^{13}C NMR spectrum (150 MHz, CDCl_3) of **2**.




Figure S5. ^1H NMR spectrum (600 MHz, CDCl_3) of **3**.