


Portable Raman spectrometer as a screening tool for characterization of Iberian dry-cured ham

A. Martín-Gómez¹, N. Arroyo-Manzanares^{2*}, M. García-Nicolás², A.I. López-Lorente¹, S. Cárdenas¹, I. López-García², P. Viñas², M. Hernández-Córdoba² and L. Arce¹


¹ Department of Analytical Chemistry, Institute of Fine Chemistry and Nanochemistry, Marie Curie Annex Building, Campus de Rabanales, 14071 Córdoba, Spain; q02magoa@uco.es (A.M.-G); q32loloa@uco.es (A.I.L.-L); qa1caarm@um.es (S.C.); qa1arjil@um.es (L.A.)

² Department of Analytical Chemistry, Faculty of Chemistry, Regional Campus of International Excellence “Campus Mare Nostrum”, University of Murcia, E-30100 Murcia, Spain; maria.garcia66@um.es (M.G.-N.); ilgarcia@um.es (I.L.-G.); pilarvi@um.es (P.V.); hcordoba@um.es (M.H.C.)

* Correspondence: natalia.arroyo@um.es (N.A.-M.)



Supplemental Figure S1. Spectra from the different samples categories: A) 50% Iberian acorn-fed, B) 100% Iberian acorn-fed, C) 75% Iberian feed-fed.



Supplemental Figure S2. Laser power optimization.