

Supplementary Materials

Hypoglycemic Effects of Novel *Panax notoginseng* Polysaccharide in Mice with Diet-Induced Obesity

Table S1. Food intakes and the contents of LDL-C, HDL-C, TG, TC, insulin, and HbA1c in the HFD-induced obese mice.

Groups	Food in- take (g/kg/d)	LDL-C (μ M)	HDL-C (mM)	TG (mM)	TC (mM)	Insulin (mM)	HbA1c (mM)
ND	182 \pm 27	0.26 \pm 0.06	3.82 \pm 0.23	1.28 \pm 0.26	3.14 \pm 0.32	17.8 \pm 1.57	172.34 \pm 15.38
HF	112 \pm 18	0.31 \pm 0.04	4.92 \pm 0.42	1.58 \pm 0.06	4.32 \pm 0.22	19.53 \pm 2.18	218.22 \pm 30.21
50 μ g/mL	108 \pm 19	0.30 \pm 0.05	4.52 \pm 0.23	1.72 \pm 0.09	3.41 \pm 0.32	18.26 \pm 3.12	163.52 \pm 12.59
100 μ g/mL	118 \pm 21	0.32 \pm 0.06	4.61 \pm 0.27	1.91 \pm 0.46	3.27 \pm 0.26	18.64 \pm 1.94	181.47 \pm 19.63
200 μ g/mL	107 \pm 22	0.30 \pm 0.12	4.91 \pm 0.32	1.68 \pm 0.16	3.51 \pm 0.18	17.52 \pm 1.37	171.43 \pm 15.29