


Correction


## Correction: Xia et al. Resveratrol Alleviates Zearalenone-Induced Intestinal Dysfunction in Mice through the NF-κB/Nrf2/HO-1 Signalling Pathway. *Foods* 2024, *13*, 1217

Sugan Xia <sup>1,2,3</sup>, Chaoyue Yan <sup>1,2</sup>, Jianhong Gu <sup>1,2</sup>, Yan Yuan <sup>1,2</sup>, Hui Zou <sup>1,2,3</sup>, Zongping Liu <sup>1,2,3</sup> and Jianchun Bian <sup>1,2,3,\*</sup>

- <sup>1</sup> College of Veterinary Medicine, Yangzhou University, Yangzhou 225009, China; liuzongping@yzu.edu.cn (Z.L.)
- <sup>2</sup> Jiangsu Co-Innovation Center for Prevention and Control of Important Animal Infectious Diseases and Zoonoses, Yangzhou 225009, China
- <sup>3</sup> Joint International Research Laboratory of Agriculture and Agri-Product Safety, Ministry of Education of China, Yangzhou University, Yangzhou 225009, China
- \* Correspondence: jcbian@yzu.edu.cn; Tel.: +86-13338850772

In the original publication [1], there was a mistake in "Figure 7. Impact of RSV on the Nrf2 signalling pathway in zearalenone-induced gut oxidative stress in mice". The research images were duplicated due to carelessness in drawing the combined images, resulting in ZEA + 50 mg/kg RSV and ZEA + 200 mg/kg RSV being the same (Figure 7d). We have corrected Figure 7d on the basis that the scientific conclusions are unaffected. The corrected Figure 7 is as follows:







Citation: Xia, S.; Yan, C.; Gu, J.; Yuan, Y.; Zou, H.; Liu, Z.; Bian, J. Correction: Xia et al. Resveratrol Alleviates Zearalenone-Induced Intestinal Dysfunction in Mice through the NF-κB/Nrf2/HO-1 Signalling Pathway. *Foods* 2024, *13*, 1217. *Foods* **2024**, *13*, 1686. https://doi.org/ 10.3390/foods13111686

Received: 14 May 2024 Accepted: 20 May 2024 Published: 28 May 2024



**Copyright:** © 2024 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (https:// creativecommons.org/licenses/by/ 4.0/). analyses of Nrf2 (magnification:  $100 \times$  and scale bar:  $50 \mu$ m). Representative Western blot images (e) and quantification of (f) HO-1 and (g) Nrf2 (n = 3). Relative expression of mRNA in the jejunum (h) NQO1, (i) GSH-PX, and (j)  $\gamma$ -GCS (n = 6). Values are presented as the mean  $\pm$  SD of each treatment. "##" p < 0.01 compared to control group; "\*" p < 0.05 and "\*\*" p < 0.01 compared ZEA group.

This correction has been approved by the Academic Editor. The original publication has also been updated.

## Reference

1. Xia, S.; Yan, C.; Gu, J.; Yuan, Y.; Zou, H.; Liu, Z.; Bian, J. Resveratrol Alleviates Zearalenone-Induced Intestinal Dysfunction in Mice through the NF-κB/Nrf2/HO-1 Signalling Pathway. *Foods* **2024**, *13*, 1217. [CrossRef] [PubMed]

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.