

Brief Report

FD-2, an Anticervical Stenosis Device for Patients Undergoing Radical Trachelectomy or Cervical Conization

Seiji Mabuchi ^{1,*}, Shoji Kamiura ¹, Takuya Saito ², Hayato Furukawa ², Azusa Abe ³ and Takashi Sasagawa ⁴

¹ Department of Gynecology, Osaka International Cancer Institute, 3-1-69, Otemae, Chuo-ku, Osaka-shi 541-8567, Japan; kamiura-sh@oici.jp

² Medical Technical Sec., Fuji Latex Co., Ltd., 1705 Chizuka-machi, Tochigi 328-0135, Japan; t-saitou@fujilandex.jp (T.S.); ha-furukawa@fujilandex.jp (H.F.)

³ Quality Control Sec., Fuji Latex Co., Ltd., 1705 Chizuka-machi, Tochigi 328-0135, Japan; a-tanifuji@fujilandex.jp

⁴ Fuji Latex Co., Ltd., 3-19-1 Kanda Nishiki-cho, Chiyoda-ku, Tokyo 101-0054, Japan; sasagawa@fujilandex.jp

* Correspondence: seiji.mabuchi@oici.jp; Tel.: +81-6-6945-1181

Supplementary

1. Designing FD-2



2. Manufacturing FD-2 (Figure 1)



3. Bench testings

- Assessment of the durability of a FD-2 (Tensile test: Figure 2)
- Assessment of leachable impurities (Extractables and Leachables test: Table 1)



- Sterilization
- Informed consent

4. Preliminary use of FD-2 (Case series: Figure 3)



5. Future clinical trials

Figure S1. Flow diagram of the FD-2 development