

SUPPORTING INFORMATION

Article

An Algerian Soil-Living *Streptomyces alboflavus* Strain as Source of Antifungal Compounds for the Management of the Pea Pathogen *Fusarium oxysporum* f. sp. *pisi*

Marco Masi ^{1,*}, Dorsaf Nedjar ^{2,†}, Moustafa Bani ^{2,*}, Ivana Staiano ³, Maria Michela Salvatore ¹, Karima Khenaka ², Stefany Castaldi ³, Jesus Garcia Zorrilla ^{1,4}, Anna Andolfi ¹, Rachele Istatico ³ and Alessio Cimmino ¹

¹ Department of Chemical Sciences, University of Naples Federico II, Complesso Universitario Monte S. Angelo, Via Cinthia 4, 80126 Naples, Italy; mariamichela.salvatore@unina.it (M.M.S.); jesus.zorrilla@uca.es (J.G.Z.); andolfi@unina.it (A.A.); alessio.cimmino@unina.it (A.C.)

² Laboratory of Biotechnology, Higher National School of Biotechnology Taoufik Khaznadar, Nouveau Pôle Universitaire Ali Mendjeli, BP. E66, Constantine 25100, Algeria; dorsafnedjar0@gmail.com (D.N.); khenakak@yahoo.fr (K.K.)

³ Department of Biology, University of Naples Federico II, Complesso Universitario Monte S. Angelo, Via Cinthia 4, 80126 Naples, Italy; ivana.staiano@unina.it (I.S.); stefany.castaldi@unina.it (S.C.); isticato@unina.it (R.I.)

⁴ Allelopathy Group, Department of Organic Chemistry, Facultad de Ciencias, Institute of Biomolecules (INBIO), University of Cadiz, C/Avenida República Saharaui, s/n, 11510 Puerto Real, Spain

* Correspondence: marco.masi@unina.it (M.M.); m.bani@ensbiotech.edu.dz (M.B.)

† Authors contributed the same way to the work.

Supporting information list

Figure S1. ¹H NMR spectrum of metacycloprodigiosin (**1**) (CHCl₃, 400 MHz).

Figure S2. ¹³C NMR spectrum of metacycloprodigiosin (**1**) (CHCl₃, 100 MHz).

Figure S3. ESI-MS spectrum of metacycloprodigiosin (**1**), recorded in positive mode.

Figure S4. EI mass spectrum at 70 eV of 4-nitrobenzamide, TMS (RI = 1864). TMS = trimethylsilyl group.

Figure S5. EI mass spectrum at 70 eV of furandimethanol, 2TMS (RI = 1427). TMS = trimethylsilyl group.

Figure S6. ¹H NMR spectrum of FC (CHCl₃, 400 MHz).

Figure S7. ¹H NMR spectrum of FE (CHCl₃, 400 MHz).




Figure S1. ¹H NMR spectrum of metacycloprodigiosin (1) (CHCl₃, 400 MHz).




Figure S2. ¹³C NMR spectrum of metacycloprodigiosin (1) (CHCl₃, 100 MHz).




Figure S3. ESI-MS spectrum of metacycloprodigiosin (1), recorded in positive mode.




Figure S4. EI mass spectrum at 70 eV of 4-nitrobenzamide, TMS (RI = 1864). TMS = trimethylsilyl group.




Figure S5. EI mass spectrum at 70 eV of furandimethanol, 2TMS (RI = 1427). TMS = trimethylsilyl group.




Figure S6. ^1H NMR spectrum of FC (CHCl_3 , 400 MHz).




Figure S7. ^1H NMR spectrum of FE (CHCl_3 , 400 MHz).