

Supplementary Material

Potent and Selective Inhibitors of Human Monoamine Oxidase A from an Endogenous Lichen Fungus *Diaporthe mahothocarpus*

Geum Seok Jeong ¹, Prima F. Hillman ², Myung-Gyun Kang ³, Sungbo Hwang ³, Jong-Eun Park ¹, Sang-Jip Nam ², Daeui Park ³ and Hoon Kim ^{1,*}

¹ Department of Pharmacy, and Research Institute of Life Pharmaceutical Sciences, Sunchon National University, Suncheon 57922, Republic of Korea; fever41@naver.com (G.S.J.); nji8009@naver.com (J.I.N.); park140201@naver.com (J.E.P.); sungtae@scnu.ac.kr (S.T.Y)

² Department of Chemistry and Nanoscience, Ewha Womans University, Seoul 03760, Republic of Korea; pri-mafitriah@gmail.com (P.F.H); sjnam@ewha.ac.kr (S.J.N)

³ Department of Predictive Toxicology, Korea Institute of Toxicology, Daejeon 34114, Republic of Korea; myung-gyun.kang@kitox.re.kr (M.-G.K.); sungbo.hwang@kitox.re.kr (S.H.); daeui.park@kitox.re.kr (D.P.)

* Correspondence: Correspondence: hoon@sunchon.ac.kr; Tel.: +82-61-750-3751

Table of Contents

Figure S1. Percentage purity of AT (1)
Figure S2. Percentage purity of HAT (2)
Figure S3. Percentage purity of MED (3)
Figure S4. ¹H NMR spectrum of alternariol (AT) (1) in CD₃OD
Figure S5. ¹³C NMR Spectrum of alternariol (AT) (1) in CD₃OD
Figure S6. COSY Spectrum of alternariol (AT) (1) in CD₃OD
Figure S7. HSQC Spectrum of alternariol (AT) (1) in CD₃OD
Figure S8. HMBC Spectrum of alternariol (AT) (1) in CD₃OD
Figure S9. ¹H NMR spectrum of 5'-hydroxy-alternariol (HAT) (2) in DMSO-<i>d</i>₆
Figure S10. ¹³C NMR Spectrum of 5'-hydroxy-alternariol (HAT) (2) in DMSO-<i>d</i>₆
Figure S11. COSY Spectrum of 5'-hydroxy-alternariol (HAT) (2) in DMSO-<i>d</i>₆
Figure S12. HSQC Spectrum of 5'-hydroxy-alternariol (HAT) (2) in DMSO-<i>d</i>₆
Figure S13. HMBC Spectrum of 5'-hydroxy-alternariol (HAT) (2) in DMSO-<i>d</i>₆
Figure S14. ¹H NMR Spectrum of mycoepoxydiene (MED) (3) in DMSO-<i>d</i>₆
Figure S15. ¹³C NMR Spectrum of mycoepoxydiene (MED) (3) in DMSO-<i>d</i>₆
Figure S16. COSY Spectrum of mycoepoxydiene (MED) (3) in DMSO-<i>d</i>₆
Figure S17. HSQC Spectrum of mycoepoxydiene (MED) (3) in DMSO-<i>d</i>₆
Figure S18. HMBC Spectrum of mycoepoxydiene (MED) (3) in DMSO-<i>d</i>₆
Figure S19. Plots of root mean square deviation (RMSD) during 1 ns MD simulation of the complex with (a) hMAO-A and AT, and (b) hMAO-B and AT




Figure S1. Percentage purity of AT (1).




Figure S2. Percentage purity of HAT (2).






Figure S3. Percentage purity of MED (3).






Figure S6. COSY spectrum of Alternariol (AT) (1) in CD₃OD.




Figure S7. HSQC spectrum of Alternariol (AT) (1) in CD₃OD.




Figure S8. HMBC spectrum of Alternariol (AT) (1) in CD₃OD.




Figure S9. ¹H NMR Spectrum of 5'-hydroxy-alternariol (HAT) (2) in DMSO-*d*₆.




Figure S10. ^{13}C NMR Spectrum of 5'-hydroxy-alternariol (HAT) (2) in $\text{DMSO-}d_6$.




Figure S11. COSY Spectrum of 5'-hydroxy-alternariol (HAT) (2) in $\text{DMSO-}d_6$.




Figure S12. HSQC Spectrum of 5'-hydroxy-alternariol (HAT) (2) in DMSO- d_6 .




Figure S13. HMBC Spectrum of 5'-hydroxy-alternariol (HAT) (2) in DMSO- d_6 .




Figure S14. ¹H NMR spectrum of Mycoepoxydiene (MED) (3) in DMSO-*d*₆.



Figure S15. ¹³C NMR Spectrum of Mycoepoxydiene (MED) (3) in DMSO-*d*₆.




Figure S16. COSY Spectrum of Mycoepoxydiene (MED) (3) in DMSO-*d*₆.




Figure S17. HSQC Spectrum of Mycoepoxydiene (MED) (3) in DMSO-*d*₆.




Figure S18. HMBC Spectrum of Mycoepoxydiene (MED) (3) in DMSO- d_6 .




Figure S19. Plots of root mean square deviation (RMSD) during 1 ns MD simulation of (a) hMAO-A and (b) hMAO-B complexes with AT.