

Fluconazole

	P1	P1- <i>mfs7</i> Δ	P1- <i>cdr1</i> Δ	P1- <i>mrr1</i> Δ	P1- <i>mfs7</i> Δ <i>cdr1</i> Δ	P1- <i>mrr1</i> Δ <i>cdr1</i> Δ
P1	0	-1	-1	1	-2	1
P1- <i>mfs7</i> Δ	1	0	0	2	-1	2
P1- <i>cdr1</i> Δ	1	0	0	2	-1	2
P1- <i>mrr1</i> Δ	-1	-2	-2	0	-3	0
P1- <i>mfs7</i> Δ <i>cdr1</i> Δ	2	1	1	3	0	3
P1- <i>mrr1</i> Δ <i>cdr1</i> Δ	-1	-2	-2	0	-3	0

Posaconazole

	P1	P1- <i>mfs7</i> Δ	P1- <i>cdr1</i> Δ	P1- <i>mrr1</i> Δ	P1- <i>mfs7</i> Δ <i>cdr1</i> Δ	P1- <i>mrr1</i> Δ <i>cdr1</i> Δ
P1	0	-1	0	-1	-1	-1
P1- <i>mfs7</i> Δ	1	0	1	0	0	0
P1- <i>cdr1</i> Δ	0	-1	0	-1	-1	-1
P1- <i>mrr1</i> Δ	1	0	1	0	0	0
P1- <i>mfs7</i> Δ <i>cdr1</i> Δ	1	0	1	0	0	0
P1- <i>mrr1</i> Δ <i>cdr1</i> Δ	1	0	1	0	0	0

Itraconazole

	P1	P1- <i>mfs7</i> Δ	P1- <i>cdr1</i> Δ	P1- <i>mrr1</i> Δ	P1- <i>mfs7</i> Δ <i>cdr1</i> Δ	P1- <i>mrr1</i> Δ <i>cdr1</i> Δ
P1	0	0	0	0	-1	-3
P1- <i>mfs7</i> Δ	0	0	0	0	-1	-3
P1- <i>cdr1</i> Δ	0	0	0	0	-1	-3
P1- <i>mrr1</i> Δ	0	0	0	0	-1	-3
P1- <i>mfs7</i> Δ <i>cdr1</i> Δ	1	1	1	1	0	-2
P1- <i>mrr1</i> Δ <i>cdr1</i> Δ	3	3	3	3	2	0




Figure S1: Pairwise comparisons of MICs between isolate P1 and derived mutants. Fold-changes are given in log2 scale. Comparisons were made always starting from top row. Values > 0 indicate MIC decreases, while values < 0 indicate MIC increases. Greyed colored boxes indicate comparisons between same isolates.