

## Supplementary material


# Preparation and optimization of bovine serum albumin nano-particles as a promising gelling system for enhanced nasal drug administration

Sandra Aulia Mardikasari 1,2, Gábor Katona 1,\* , Bence Sipos 1, Rita Ambrus 1 and Ildikó Csóka 1

<sup>1</sup> Institute of Pharmaceutical Technology and Regulatory Affairs, Faculty of Pharmacy, University of Szeged, Eötvös St. 6, H-6720 Szeged, Hungary; [sandraaulia@unhas.ac.id](mailto:sandraaulia@unhas.ac.id) (S.A.M.); [katona.gabor@szte.hu](mailto:katona.gabor@szte.hu) (G.K.); [sipos.bence@szte.hu](mailto:sipos.bence@szte.hu) (B.S.); [ambrus.rita@szte.hu](mailto:ambrus.rita@szte.hu) (R.A.); [csoka.ildiko@szte.hu](mailto:csoka.ildiko@szte.hu) (I.C.)

<sup>2</sup> Faculty of Pharmacy, Hasanuddin University, Makassar 90245, Indonesia; [sandraaulia@unhas.ac.id](mailto:sandraaulia@unhas.ac.id) (S.A.M.)

\* Correspondence: [katona.gabor@szte.hu](mailto:katona.gabor@szte.hu) (G.K.)



**Figure S1.** Response surface plots showing the effect of independent variables PW – Ethanol (a), BSA – Ethanol (b) and PW – BSA (c) on Z-average as well as PW – Ethanol (d), BSA – Ethanol (e) and PW – BSA (f) on PdI.