

**Table S1.** Univariable (A) and Multivariable Cox Regression (B) analyses for TTFT with identification of significant Independent Predictors via backward elimination strategy.

Univariable analysis (A)		95% CI		
Variables	HR	Lower limit	Upper limit	P-value
<i>IGHV</i> , unmutated	4.708	3.085	7.185	<0.001
ZAP-70, positive	2.914	1.914	4.434	<0.001
del11q	6.341	3.554	11.315	<0.001
del17p	4.076	1.63	10.191	0.003
NOTCH1, mutated	2.112	1.272	3.507	0.004
B2M, abnormal	2.23	1.42	3.503	<0.001
Rai stage, I-II	1.782	1.157	2.747	0.009
Lymphocytosis, >5x10 <sup>9</sup> /L	1.806	1.005	3.245	0.048
CD38, positive	2.692	1.754	4.134	<0.001
Multivariable Cox regression analyses (B)				
<i>IGHV</i> , unmutated	3.452	2.273	5.244	<0.001
del11q	2.871	1.556	5.299	<0.001
del17p	9.616	2.212	41.813	0.003
NOTCH1, mutated	1.866	1.129	3.086	0.015
B2M, abnormal	1.940	1.351	2.785	<0.001
Rai stage, I-II	2.117	1.443	3.107	<0.001