

Heterogenous organo- and metal catalysis using phosphine oxide derivatives anchored on multiwalled carbon nanotubes

Maria Cristina Ligi ¹, Anna Flis ², Giacomo Biagiotti ^{1,3}, Giulia Serrano ^{3,4}
K. Michał. Pietrusiewicz ^{2,*} and Stefano Cicchi ^{1,3,5,*}

¹ Department of Chemistry, Università di Firenze, via della Lastruccia 3-13, 50019 Sesto Fiorentino, Firenze, Italy; mariacristina.ligi@unifi.it (M.C.L.); giacomo.biagiotti@unifi.it (G.B.)

² Department of Organic Chemistry, Maria Curie-Skłodowska University, ul. Gliniana 33, 20-614 Lublin, Poland; a.flis@poczta.umcs.lublin.pl

³ INSTM (National Interuniversity Consortium of Materials Science and Technology), Via G. Giusti 9, 50121 Firenze, Italy

⁴ Department of Industrial Engineering, Università di Firenze, via di S. Marta, 3-50139 Firenze, Italy; giulia.serrano@unifi.it

⁵ Institute of Chemistry of OrganoMetallic Compounds ICCOM-CNR, Via Madonna del Piano 10, 50019 Sesto Fiorentino, Italy

* Correspondence: kazimierz.pietrusiewicz@poczta.umcs.lublin.pl (K.M.P.); stefano.cicchi@unifi.it (S.C.)

* Correspondence: kazimierz.pietrusiewicz@poczta.umcs.lublin.pl; stefano.cicchi@unifi.it.

Supporting material




Figure S1. ^1H NMR of 1-phenyl-3-benzylaminophospholane 1-oxide **A**




Figure S2. ^{13}C NMR of 1-phenyl-3-benzylaminophospholane 1-oxide **A**




Figure S3. ^{13}P NMR of 1-phenyl-3-benzylaminophospholane 1-oxide **A**




Figure S4. ^1H NMR of 3-azido-1-phenylphospholane 1-oxide (**5**).




Figure S5. ^{13}C NMR Spectrum of 3-azido-1-phenylphospholane 1-oxide (**5**).




Figure S6. gCosy NMR Spectrum of 3-azido-1-phenylphospholane 1-oxide (**5**).




Figure S7. FT-IR Spectrum of 3-azido-1-phenylphospholane 1-oxide (5).




Figure S8. ESI-MS of 3-azido-1-phenylphospholane 1-oxide (5).




Figure S9. ox-CNTs 2: TEM.




Figure S10 FT-IR Spectrum of ox-CNTs 2.




Figure S11. FT-IR spectrum of 6




Figure S12. FT-IR spectrum of 7.




Figure S13. FT-IR spectrum of **8**.




Figure S14. FT-IR spectrum of **9**




Figure S15. FT-IR spectrum of **10**.




Figure S16. GC-MS Spectrum of N-Benzylbenzamide.




Figure S17. ¹H NMR Spectrum of N-Benzylbenzamide.




Figure S18. GC-MS Spectrum of methyl cinnamate.




Figure S19. ¹H NMR of methyl cinnamate.




Figure S20. GC-MS Spectrum of benzyl 4-nitrobenzoate.




Figure S21. ¹H NMR of benzyl 4-nitrobenzoate.




Figure S22. GC-MS Spectrum of trans-Stilbene.




Figure S23. ¹H NMR of trans-stilbene.




Figure S24. GC-MS Spectrum of trans-cinnamic acid.




Figure S25. ¹H NMR of trans-cinnamic acid.