

Supplementary Figure

Drought and Salinity Stresses Response in Three Korean Native Herbaceous Plants and Their Suitability as Garden Plants

Saeng Geul Baek ^{1,2}, Jung Won Shin ¹, Jae Ik Nam ¹, Jeong Min Seo ¹, Jung Min Kim ¹ and Su Young Woo ^{2,*}

¹ Gardening material Division, Korea Arboreta and Gardens Institute, Sejong-si 30106, Republic of Korea; bsg1175@koagi.or.kr (S.G.B.); jwony1208@naver.com (J.W.S.); silhouette@koagi.or.kr (J.I.N.); jmseo3@koagi.or.kr (J.M.S.); kimini@koagi.or.kr (J.M.K.)

² Department of Environmental Horticulture, University of Seoul, Seoul 02504, Republic of Korea;

* Correspondence: wsy@uos.ac.kr ; Tel.: +82-10-3802-5242 (S.Y.W.)




Figure S1. Change of air temperature and relative humidity in the greenhouse during a month(June~July) of study period. Data were represented as the daily average.




Figure S2. Mean relative growth rate (RGR) of (a) height, (b) diameter and (c) no.leaf in three native species under drought stress. Significant differences, based on Tukey's HSD test, are denoted by lowercase letters. The data are presented as mean \pm standard deviation ($n = 5$). Bars marked with the same letters are not significantly different ($p < 0.05$). ns: not significant.




Figure S3. Mean relative growth rate (RGR) of (a) height, (b) diameter and (c) no.leaf in three native species under salinity stress. Significant differences, based on Tukey's HSD test, are denoted by lowercase letters. The data are presented as mean \pm standard deviation ($n = 5$). Bars marked with the same letters are not significantly different ($p < 0.05$). ns: not significant.