





Review

Microorganisms, the Ultimate Tool for Clean Label Foods?

Giorgia Perpetuini ¹, Pumnat Chuenchomrat ^{2,†}, Valentin Pereyron ^{3,†}, Maxime Haure ^{3,†}, Da Lorn ^{3,4,†}, Le-Ha Quan ^{5,†}, Phu-Ha Ho ^{5,6,†}, Tien-Thanh Nguyen ^{5,6,†}, Thi-Yen Do ^{6,†}, Quyet-Tien Phi ^{7,†} , Thi Kim Chi Nguyen ^{3,†}, Hélène Licandro ^{3,†}, Son Chu-Ky ^{5,6,†}, Rosanna Tofalo ¹ , Warissara Kasikonsunthonchai ², Saowalak Adunphatcharaphon ², Awanwee Petchkongkaew ^{2,8,*} , and Yves Waché ^{2,3,*} 

- ¹ Faculty of Bioscience and Technology for Food, Agriculture and Environment, University of Teramo, 64100 Teramo, Italy
- ² School of Food Science and Technology, Faculty of Science and Technology, Thammasat University, Pathumthani 12121, Thailand; cpumnat@tu.ac.th (P.C.); warissara.kasi@dome.tu.ac.th (W.K.); s.adunphatcharaphon@outlook.co.th (S.A.)
- ³ International Joint Laboratory, Tropical Bioresources & Biotechnology, School of Biotechnology and Food Technology, Hanoi University of Science and Technology, PAM UMR A 02.102, AgroSup Dijon, University Bourgogne Franche-Comté, Dijon 21000, France
- ⁴ Chemical and Food Engineering Department, Institute of Technology of Cambodia, Phnom Penh, Cambodia
- ⁵ School of Biotechnology and Food Technology, Hanoi University of Science and Technology, Hanoi 100000, Vietnam
- ⁶ International Joint Laboratory, Tropical Bioresources & Biotechnology, School of Biotechnology and Food Technology, Hanoi University of Science and Technology, PAM UMR A 02.102, AgroSup Dijon, University Bourgogne Franche-Comté, Hanoi 100000, Vietnam
- ⁷ Institute of Biotechnology, Vietnam Academy of Science and Technology, Hanoi 100000, Vietnam
- ⁸ Institute for Global Food Security, School of Biological Science, Queen's University Belfast, Belfast BT7 1NN, Northern Ireland, UK
- * Correspondence: awanwee@tu.ac.th (A.P.); y.wache@agrosupdijon.fr (Y.W.)
- † Research Network: Tropical Fermentation Network.



Citation: Perpetuini, G.; Chuenchomrat, P.; Pereyron, V.; Haure, M.; Lorn, D.; Quan, L.-H.; Ho, P.-H.; Nguyen, T.-T.; Do, T.-Y.; Phi, Q.-T.; et al. Microorganisms, the Ultimate Tool for Clean Label Foods? *Inventions* **2021**, *6*, 31. <https://doi.org/10.3390/inventions6020031>

Academic Editor: Monique Lacroix

Received: 26 March 2021

Accepted: 27 April 2021

Published: 30 April 2021

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2021 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

Abstract: Clean label is an important trend in the food industry. It aims at washing foods of chemicals perceived as unhealthy by consumers. Microorganisms are present in many foods (usually fermented), they exhibit a diversity of metabolism and some can bring probiotic properties. They are usually well considered by consumers and, with progresses in the knowledge of their physiology and behavior, they can become very precise tools to produce or degrade specific compounds. They are thus an interesting means to obtain clean label foods. In this review, we propose to discuss some current research to use microorganisms to produce clean label foods with examples improving sensorial, textural, health and nutritional properties.

Keywords: clean-label; technology additive; sensorial additive; anti-staling; bioremediation; biosurfactants; bio-preservation; antibiofilm; antinutrition; beneficial microorganism

1. Introduction

Clean label is a marketing concept aiming at giving confidence to consumers. Indeed, in the last few decades, consumers may have perceived the food industry as at risk of poisons in which all possibilities are used to do business at the expense of consumers, society and the environment. Applying the clean-label concept to food consists in washing the label from additives, especially those perceived as chemical and artificial, to go back to traditional foods reminding us of “Grandma’s cooking”.

Whereas in some fields, biotechnology is only limited by technical possibilities, in the food domain in which consumers are pushing the debate on ethical concepts, naturality and sustainability, biotechnology grows between many constraints that have arose to preserve people and the environment. As a result, the food biotechnologist is used to trying to bring

about new technologies responding to technical issues as well as ethical and environmental ones. The use of microorganisms to wash the food label is one of these typical questions addressed by food biotechnologists. The food industry can produce high technical quality food, but this food is highly processed, using thorough cracking additives, many pesticides and crop preservation chemicals resulting in a high carbon cost, environmental pollution and food inducing metabolic syndrome and cancers in consumers. The concept of using microorganisms to achieve clean-label food is thus quite simple: microorganisms should, by their activity, produce active molecules from precursors naturally present in the food matrix. However, this concept should be usable also for non-fermented foods, meaning that, in this case, the microbial activity should not modify the sensorial properties of the product towards fermented notes.

From a regulatory point of view, this concept brings much discussion with business trying to occupy the field to obtain benefits while food and health agencies try to protect the consumers against this. In this review, we propose to present several examples of microbial applications in a clean-label strategy. These examples will deal with the use of microorganisms to replace technology, sensorial, biopreservation, bioremediation and health additives. We shall focus on recent work or current strategies and only refer briefly to already existing applications.

2. Technology Additives

Foods are usually very complex structures including all nutritional components, whatever their hydrophobicity, solubility, physicochemical status. Their textural organisation is thus prone to modification during shelf life and many chemical agents can be added to stabilise them. However, this domain of technology additives is very controversial as good quality products in terms of texture/structure and physico-chemical stability are often in the category of over-processed food, which results in bad marks in food score applications. In this context, microorganisms can bring a lot of functionalities without addition of chemicals. In this part, we will present some examples concerning how we can use microorganisms to avoid starch retrogradation in bread products and how microbial biosurfactants can bring interesting textural properties to food.

2.1. Staling

Starch retrogradation occurs in bread and starch products [1]. It is an issue in this field as it is responsible for stale bread, but it brings also desirable properties to other products like breakfast cereals or rice vermicelli. It is the result of a rearrangement of amylose and amylopectin molecules from gelatinised starch upon cooling [2]. During cooling, amylose forms a network around amylopectin granules. This network is reinforced by the rearrangement of amylose into double helices crystalline structures. Later during storage, amylopectin rearranges to form also crystalline structures, contributing to the hardness of the system. Several additives can interact with amylose, mobilising the molecules out of the network. For instance, monoglycerides, coded as E471 additives in the European system, can decrease amylose crystallisation. However, these E471 additives are typically a target of the clean-label strategy.

In the microorganism-induced clean-label strategy, microbial catalysts hydrolyse triglycerides present in natural plant oil into diglycerides, monoglycerides and free fatty acids. Contrasting with the use of enzymes, they can be labelled in the well-accepted “starter” category. One microorganism we have tested is the yeast *Yarrowia lipolytica*. This species is well-known and studied for its capacity to degrade hydrophobic compounds [3]. It possesses a wide family of lipases, including extracellular ones that are produced depending on the fatty substrate present in the medium [4]. From a technological point of view, mutants altered for the regulation of lipase synthesis or lipase production would be more attractive as they can be more efficient in the precision catalysis required. However, one of the constraints of microorganisms for foods is that, in almost all world markets, microorganisms for food usage cannot be genetically modified and only natural mutants

are usable. This constraint is often not insurmountable even if no examples are available to produce specific lipases in *Yarrowia lipolytica*. Indeed, the difficulty is to find the right and easy-to-use screening procedure. Natural improvement of the tolerance of *Y. lipolytica* to toxic alcohols has already been made [5]. Another constraint is that *Y. lipolytica* must not exhibit any sensorial impact other than decreasing staling. This yeast species is well known for its ability to degrade lipids and proteins, producing thereby aroma compounds [3,6]. In the case of this aerobic yeast, this point can also be relatively easily overcome through a sequential utilisation of the yeast in the production process and inactivation after use. Eventually, the yeast must not pose any risks to consumers' health and this yeast, which is Generally Recognized As Safe (GRAS), has been studied for its applications as a starter showing high benefits [7].

Another family of additive popular to limit staling is composed of glucidic hydrocolloids. These compounds can have an impact on the plasticity of the amorphous regions of crumbs, where they can increase water retention or inhibit gluten-starch interactions [8]. Lactic acid bacteria can produce several products of this family under the form of exopolysaccharides [9]. Dextrane is one such bacterial compound which effect has been studied on starch retrogradation [10,11].

2.2. Microbial Biosurfactants

Emulsifiers are amphiphilic compounds i.e., compounds possessing both hydrophobic and hydrophilic parts, exhibiting surface activity properties. They tend to accumulate at interfaces making them suitable to stabilise emulsions. These molecules can come from diverse origins, including petroleum industry and they can also exhibit many bioactivity properties. They could thus have a role to play in many modern food-related diseases [12]. Research has thus been oriented towards the development of new natural emulsifiers [13]. Biosurfactants are produced by living cells, especially microorganisms like bacteria, molds and yeasts. As emulsifiers, they are like chemical synthetic surfactants, amphiphilic compounds [14] consisting of hydrophilic and hydrophobic moieties and they can reduce surface and interfacial tensions [15]. In biosurfactants, hydrophilic moieties can be carbohydrates, carboxylic acids, phosphates, amino acids, cyclic peptides, and alcohols. However, the hydrophobic moieties of the biosurfactants are usually long-chain fatty acids, hydroxyl fatty acids and α -alkyl- β -hydroxyl fatty acids [16]. Based on their chemical structures, the microbial biosurfactants are classified into four groups: glycolipids, phospholipids, and fatty acids, lipopeptides and polymeric biosurfactants [17,18] as shown in Table 1.

Table 1. Microbial biosurfactants.

Biosurfactants	Producing Microbes	References
Glycolipids		
Rhamnolipids	<i>Pseudomonas aeruginosa</i>	[19]
Sophorolipids	<i>Candida bombicola</i>	[20]
Trehalolipids	<i>Rhodococcus erythropolis</i>	[21]
	<i>Mycobacterium</i> sp.	[22]
Lipopeptides		
Putisolvins I and II	<i>Pseudomonas putida</i>	[23]
Surfactin	<i>Bacillus subtilis</i>	[24]
Pseudofactin II	<i>Pseudomonas fluorescens</i>	[25]
Serrawettin	<i>Serratia marcescens</i>	[26]
Iturin A	<i>Bacillus amyloliquefaciens</i>	[27]
Fengycin	<i>Bacillus licheniformis</i>	[28]

Biosurfactant agents also show potential properties such as emulsification, functional additives, detergency, lubrication, phase dispersion, foaming, and solubilisation in many industries [29,30]. They show unique advantages including lower toxicity, better environmental compatibility, higher biodegradability, and specific activity when compared with chemical agents [31]. Mouafo et al. (2018) [32] reported that a glycolipid biosurfactant pro-

duced by *Lactobacillus* spp. could be used as an emulsifier in the food industry. Varvaresou and Iakovou (2015) [33] reviewed that sophorolipid ester was interesting as an ingredient in cosmetic products such as rouge, lip cream, and eye shadow. Furthermore, trehalose lipid produced by *Rhodococcus erythropolis* 3C-9 exhibited oil spill cleanup application [34]. In food, it can be noted that the bacteria themselves can exhibit surface active properties as shown on the use of *Lactococcus* strains to stabilise or destabilise emulsions [35–37].

Several studies are currently being carried out to develop the use of microbial biosurfactants instead of chemical ones in food. However, biosurfactants not only show the aforementioned properties, but they can also exhibit biological activities such as anti-microbial, anti-adhesion, and anti-biofilm formation activities. These properties can be of interest, but they require also a complete check before using a biosurfactant-producing microorganism.

3. Sensorial Additives

A major quality of food is to be attractive for consumers. This is true when a company wants consumers to buy back its products as well as to maintain a good nutritious state for patients losing their appetite. In the food transition towards a more sustainable system, sensorial properties are particularly important when new products are formulated with plants bringing off flavour or off colours. The traditional strategy in this case consists in using flavours or flavour-masking compounds that will lengthen the list of ingredients while the microorganism-based clean-label strategy proposes to select microorganisms able to produce flavour or colour and degrade off-flavours. Some examples concerning the bitterness of naringin and legumes off-colours are given in this section.

3.1. Naringin

Naringin (4',5,7-trihydroxy flavanone 7-rhamnoglucoside) is a flavanone glycoside that is abundant in citrus fruits, mostly in the albedo and the peel [38]. With the limonin glycoside, naringin is considered as the molecule responsible of their bitterness, major off flavour when processing juice from citrus [39]. The naringin content is closely linked to the maturity of the fruit, its content being reduced with the maturity of the fruit [40]. Because of its high rate, the industrial processing of citrus generally uses immature fruits containing high contents of naringin. Thus, researchers have put efforts into finding ways to decrease the content of naringin in citrus. To do so, some physico-chemical methods have been developed, generally implying the use of resins, affinity polymers, cyclodextrin [41–43]. But these techniques involve the inclusion of additives and tend to impact the organoleptic characteristics of the processed juice [43,44]. Naringin can also be converted into naringenin by naringinase, an enzyme containing both α -L-rhamnosidase (E.C 3.2.1.40) and β -D-glucosidase (E.C 3.2.1.21) activities [43,45]. First, the enzyme breaks the bond between the rhamnose and glucose moieties of the naringin, producing pruning. Pruning is then hydrolysed, producing both D-glucose and naringenin, bitterless compound. This enzyme can directly be added to the juice—freely or immobilised [42,43] and can easily be produced by microorganisms, mostly filamentous fungi [43,46–48]. The enzyme production is generally induced by the addition of naringin, from 0.1 to 0.5% of the total medium nutrients [49]. The purified enzymes have a maximum activity temperature around 50 °C but are more thermically stable at 40 °C [50,51]. The range of pH stability is generally from 4 to 8 [45,50,51]. In 2016, Srikantha et al. [52] reached an activity as high as 449.58 U/g of dry matter in solid state fermentation for *Aspergillus flavus*. Some studies focused on the capacity of bacteria to produce naringinases, such as *Bacillus* spp. [53–55], *Lactiplantibacillus* (*L.*) *plantarum* [56], *Clostridium stercorarium* [57] or *Pseudomonas paucimobilis* [58]. Under optimum conditions for submerged culture, the production of naringinase reached 12.05 U/L for *Bacillus methylotrophicus* [54]. Similarly, Zhu et al. [55] characterized an enzyme produced by *Bacillus amyloliquefaciens*, which could reduce 97% of initial naringin in a pomelo juice. These results clearly indicate that both filamentous fungi and bacteria have the capacity of debittering citrus in juice processing industry. The goal now is to find a microorganism able to degrade multiple phenolic glycosides, which could be used for

different applications. Indeed, most of enzymes have an activity highly specific for the nature of the bond between the glycosidic and aglyconic moiety (rutinoside-7-O-heperetin versus rutinoside-3-O-quercetin for example) and for the nature of the bond between the two sugars moieties (2 versus 6-O- α -L-rhamnosyl-D-glucose for example). Information about enzymes showing activities independent of the nature of the bond are scarce but are highly interesting for futures screening of glycosidases-producing microorganisms, which can possibly be used for a wide variety of applications.

3.2. Green-Notes in Legume Products

Legume-based products represent an interesting source of non-animal proteins due to their rich amount and diversity of essential and non-essential amino-acids [59]. In Europe, the main issue for the development of such products is the sensory acceptance by consumers. Indeed legume-based products are linked to “green”, “grassy” or “leafy” descriptors [60,61]. Removing or masking undesirable tastes by means of biotechnology is a way of developing new alternative food products without using additives or heavy processes. The development of green-notes flavours is linked to the oxidative degradation of fatty-acid by enzymatic and non-enzymatic pathways during process and storage [62,63]. Green notes are related to many volatile compounds such as aldehydes, alcohols, esters, or ketones [64]. Hexanal and its derivatives have been widely associated with green characteristics such as cut grass and leafy descriptors [65,66]. Nevertheless, green characteristics appear to depend not on the presence of isolate molecules but on the association of multiple compounds leading to various green description. Moreover, each modification on the aromatic mix leads to changes on the green perception balancing between green fruity and green grass/leafy [67]. Reducing the green characteristic of legume-based products might be complex according to multiple origins of it and its evolution during the making process. Fermentation appears to be a safe, cheap, and natural way to try to improve aromatics properties of legume-based products. This process has been widely used since thousands of years in order to preserve and improve food quality. Fermentation by lactic acid bacteria (LAB) on legumes derivatives products such as protein extract, legume-based milk or raw legumes have been investigated among the literature. Fermentation of pea and lupin protein extracts by *L. plantarum* and *Pediococcus pentosaceus* separately, leads to a modification of green markers quantity, such as a diminution of hexanal content [68,69]. Fermentation of soy milk and peanut milk by *L. acidophilus*, *L. (Lactocaseibacillus) casei*, *L. delbrückii*, *Streptococcus thermophilus* also demonstrates the ability to decrease and eliminates hexanal from milk [70,71]. The elimination of hexanal is a good start for improving organoleptic quality of legume-based products, but not enough to completely eliminate green notes due to other compounds. Fermentation by co-cultures of *L. delbrückii* ssp. *bulgaricus* and *S. salivarius* ssp. *thermophilus* leads to a modification of the aromatic profile of peanut milk, by decreasing green flavour and enhancing creamy flavour and sourness [71]. The transformation by LAB allows us to modify the aromatic profile by decreasing green-related compounds and enhancing other flavour. Moreover, the anti-green note-effect provided by some microbial cultures can be sufficient in one food matrix but not in another. Investigations are still needed to apply this clean label mean of inactivation of off flavours in all conditions but reaching this goal might be possible by selecting strains exhibiting precise metabolic activities. Our recent results have shown that when screening LAB activities towards aldehydes, it was possible to discriminate between strains reducing all aldehydes and strains reducing preferably a class of aldehydes depending on carbon chain saturation or length [72].

4. Bio-Preservation and Bioremediation Agents

The use of microorganisms for bio-preservation purposes has already been the subject of several reviews papers and will not be developed in this section. Bacteria able to produce antifungal weak acids are already used in bread applications to avoid the use of chemical preservatives [73] and bacteria able to produce antimicrobial peptides such as bacteriocins are used as starter in several products [74]. In this section, we will review the use of biosurfactants-producing microorganisms in bio-preservation strategies.

4.1. Antibacterial Activity of Biosurfactants

Biosurfactants exhibiting antimicrobial activity to control the growth of food pathogens are the subject of many studies [75–83]. Biosurfactant often exhibit detergent properties causing cell membrane destructure and permeabilisation [84]. These interactions are according to the theory of “like dissolves like”. Therefore, their combinations cause the leakage of variety of substances [85].

Several microbial biosurfactants have shown antimicrobial activity against bacteria and fungi [86–88]. Rhamnolipid biosurfactant from *Pseudomonas aeruginosa* AT10 showed inhibitory activity against several microorganisms including, *Escherichia coli*, *Micrococcus luteus*, *Alcaligenes faecalis*, *Staphylococcus epidermidis*, *Penicillium crysogenum* and *Rhizoctonia solani* [89]. Rufino et al. (2011) [88] reported that a biosurfactant named after the author Rufisan, produced by *Candida lipolytica* UCP 0988 showed antibacterial activity against *Streptococcus* spp. with concentration of 12 mg/mL. Padmapriya et al. (2013) [90] reported that biosurfactant from *Candida tropicalis* also showed antimicrobial activities against *Bacillus* spp., *C. albicans*, *Citrobacter* spp., *E. coli*, *Klebsiella pneumoniae*, *Proteus mirabilis*, *Pseudomonas aeruginosa*, *Salmonella* spp. and *Staphylococcus aureus*. A potent anti-bacterial activity of *Brevibacterium casei* affected the growth reduction of *P. aeruginosa* and *E. coli* [91]. Lipopeptide from *B. licheniformis* strain M104 demonstrated high activity against *S. aureus* [92]. Falardeau et al. (2013) [93] also reported that cyclic lipopeptides produced by *B. subtilis* showed anti-microbial activity against plant pathogenic fungi. Biosurfactant from *B. pumilus* DSVP18 showed anti-microbial activity against *B. cereus*, *S. aureus*, *S. enteritidis*, *E. coli*, and *Paenibacillus larvae* with concentration of 30–35 µg/mL [94]. The lipopeptide derived from *B. cereus* NK1 showed anti-microbial properties against Gram-positive, Gram-negative bacteria, and fungi [95]. In addition, Das et al. (2008) [96] also reported that lipopeptide biosurfactant from marine *B. circulans* showed growth inhibition of *E. coli*, *S. typhimurium*, and *S. aureus*.

However, all these strains are hardly usable as clean label starters in food because of potential hazards or sensorial impact. Fortunately, lactic acid bacteria which are often Qualified Presumption of Safety species used in foods, are also microbes reputed to produce biosurfactants [97]. Biosurfactants derived from *Lactococcus lactis* showed microbial inhibition against multi-drug resistant pathogens including *E. coli* and methicillin resistant *S. aureus* [98]. *Lactocaseibacillus paracasei* biosurfactant presented an antibacterial activity against *E. coli*, *Streptococcus agalactiae* and *S. pyogene* with concentration of 25 mg/mL [87]. Sharma and Saharan (2014) [99] also reported that biosurfactants from *L. casei* MRTL3 showed antimicrobial activity against several pathogens, including *S. aureus* ATCC 6538P, *S. epidermidis* ATCC 12228, *B. cereus* ATCC 11770, *Listeria monocytogenes* MTCC 657, *L. innocua* ATCC 33090, *Shigella flexneri* ATCC 9199, *S. typhi* MTCC 733 and *P. aeruginosa* ATCC 15442. Biosurfactant produced by *L. plantarum* CFR 2194 also showed antimicrobial activity against *E. coli* ATCC31705, *E. coli* MTCC 108, *S. typhi*, *Yersinia enterocolitica* MTCC 859 and *S. aureus* F 722 by using well diffusion method [100]. Gudina et al. (2015) [101] reported that 5 mg/mL of biosurfactant from *L. agalis* CCUG31450 exhibited the growth inhibition of *S. aureus*, *P. aeruginosa* and *S. agalactiae*.

4.2. Antifungal Activity of Biosurfactants

Biosurfactants also represent antifungal activity against fungal mycelium and spore [102]. *Botrytis cinerea*, a fruit spoilage mold was inhibited by the lipopeptide biosurfactant from *B. amyliuefaciens* [103]. Kilani-Feki et al. (2016) [104] also reported that microbial biosurfactants showed a 79% decay inhibition of tomato colonisation by *B. cinerea*. The lipopeptide biosurfactant derived from *Bacillus marinus* B-9987 was published as a safe antifungal substance against grey mold of *B. cinerea* [105]. In addition, Torres et al. (2016) [106] reported that the biosurfactants from *Bacillus* spp. showed antifungal activity against soybean pathogenic fungus of *Macrophomina phaseolina*. An interesting antifungal activity was reported by Abalos et al. (2001) [89]. They showed that *Aspergillus niger*, *B. cinerea*, *Chaetoniium globosum*, *P. chrysogenum*, and *Rhizoctonia solani* were inhibited by rhamnolipid produced by *P. aeruginosa* AT10.

This activity can also concern pathogenic molds. This is of course less related with food processing but can contribute to decrease the number of pesticides in food. For instance, *Phytophthora cryptogea*, causing rotting of fruits and flowers, was inhibited by lipopeptide produced by strains of *P. fluorescent* [107]. Mnif et al. (2015) [102] revealed that *Fusarium solani*, a potato pathogenic fungus was undergoing a 78% inhibition by *B. subtilis* SPB1 lipopeptide biosurfactant after 20 days of incubation. Moreover, the 0.02 and 3.3 mg/mL SPB1 lipopeptide biosurfactant also inhibited the seed-borne pathogenic fungus of *R. bataticola* and *R. solani*, respectively [83]. Furthermore, Joshi et al. (2008a) [81] studied the antifungal activity of *B. subtilis* 20B lipopeptide biosurfactant by using the disc diffusion method. The results of this study showed that *B. subtilis* 20B lipopeptide biosurfactant has antifungal activity against several natural contaminating fungi such as *Fusarium oxysporum*, *Alternaria burnsii*, *Cryosporium indicum* and *R. bataticola*. The antifungal activity of biosurfactant was explained by González-Jaramillo et al. (2017) [108]. They studied the effect of fengycin C, a lipopeptide biosurfactant from *B. subtilis* EA-CB0015 on *Mycosphaerella fijiensis* mycelium and spore morphology changes by using dipalmitoylphosphatidylcholine (DPPC), a fungal membrane model. The results revealed that fengycin C, the lipopeptide biosurfactant was able to change the fungal membrane model by dehydrating the polar head groups of cell membranes bilayer, causing the loss of its permeable properties. Moreover, repulsion of charges of amino acid and polar bilayer might also be involved in the destabilisation of cell structure [108].

Interestingly, Jadhav et al. (2011) [109] studied the biosurfactant produced by *Enterobacter* sp. MS16 on *A. niger* and *P. chrysogenum* spore germination. These fungal spore germinations were also inhibited by 12.5 mg/mL biosurfactant. Yoo et al. (2005) [110] also demonstrated that the rhamnolipid and sophorolipid biosurfactants showed the zoospore lysis activity against *Phytophthora* spp. and *Pythium* spp. Gond et al. (2015) [111] investigated for antifungal activity by iturin A from *B. amyloliquifaciens* against maize phytopathogenic fungus, *F. moniliforme*. They revealed that this lipopeptide biosurfactant with 500 µg/disk was strongly inhibited the mycelium elongation of *F. moniliforme* by interacting with fungal hyphae.

As a conclusion, many microbial biosurfactants are efficient against food spoilage or pathogenic strains. LAB biosurfactants can be used against food bacteria whereas bacilli bacteria produce often antifungal compounds. However, it is important to check whether these surface-active compounds can exhibit other properties that could limit their use in food.

4.3. Bioremediation

Apart from bio-preservation, numerous microorganism can also exhibit some ability to degrade toxic substances. This is referred to as a “bioremediation process” which is a bioprocess that can convert toxic substances (e.g., pesticides) or toxic contaminants (i.e., mycotoxins) or anti-nutrients such as phytates (which cause a decrease in iron availability) or biogenic amines. Nowadays, a worldwide serious agricultural threat is mycotoxin. It is recognized as an unavoidable risk. Many factors that influence the contamination level are environmental (such as weather and insect infestation) which are difficult or impossible to control. Therefore, this section attempts to review and discuss mainly on mycotoxin bioremediation.

Mycotoxins, a large group of toxic secondary metabolites, are produced primarily by a group of filamentous fungi mainly in the genera *Fusarium*, *Penicillium*, *Aspergillus* and *Alternaria*. They can contaminate food and feedstuffs at pre- and post-harvest stages. Currently, approximately of 60–80% all global agricultural commodities are contaminated with mycotoxins [112]. The most frequently found are aflatoxins, ochratoxins, zearalenone, deoxynivalenol, fumonisin B₁, T₂ and HT-2. There are numerous strategies, either based on physical or chemical treatments, that can be applied to mitigate against this problem. However, the application of biological means of mycotoxin reduction using microorganisms is received increasing interest from scientists due to its low cost, the broad spectrum of

mycotoxins that can be targeted, the minimal side effects regarding nutrient status of the food, minimal training requirements for those applying the microorganisms, and its suitability for a wide range of liquid and solid food types [113]. Mechanism of action will involve either adsorption by cell wall or degradation by enzyme depending on species and strains of microorganisms. Watanakij et al., 2020 [114] demonstrated the application of an extracellular fraction from *Bacillus subtilis* BCC42005 with water as a soaking agent for maize. The result revealed that aflatoxin B₁ was reduced after 120 min contact time without any changed appearance of the corn kernel. Table 2 summarises some microorganisms which exhibit the potential to reduce mycotoxin loads.

Table 2. Potential microorganism for mycotoxins bioremediation.

Mycotoxin	Microorganism	Reduction Capacity (%)	References
Adsorption			
Aflatoxins	<i>L. casei</i>		
	<i>L. plantarum</i>	25–61	[115]
	<i>L. fermentum</i>		
	<i>L. casei</i>	14–49	[116]
	<i>L. rhamnosus</i> GG	80	[117]
	<i>L. rhamnosus</i> LC-705		
	<i>Lactobacillus</i> spp.		
	<i>Bifidobacterium</i>	5.6–59.7	[118]
	<i>Lactococcus</i> strains		
	<i>Enterococcus faecium</i> M74 and EF031	29.0–33.7	[119]
	<i>L. plantarum</i>	45–100	[120]
	<i>B. bifidum</i> 1900		
	<i>B. pseudolongum</i> 20,099		
	<i>B. infantis</i> 1912	20–50	[121]
	<i>L. casei</i>		
	<i>Lactobacillus delbrueckii</i> subsp. <i>bulgaricus</i> CH-2	18.7	[122]
	<i>L. plantarum</i>	81	[123]
	<i>Lactococcus lactis</i>		
	<i>S. thermophilus</i>		
	<i>L. bulgaricus</i>	11–34	[124]
<i>L. plantarum</i>			
<i>L. paracasei</i> LOCK 0920			
<i>L. brevis</i> LOCK 0944	39–55	[125]	
<i>L. plantarum</i> LOCK 0945			
<i>L. plantarum</i> C88	60	[126]	
Fumonisin	<i>L. paraplantarum</i> CNRZ1885	2–27	[127]
	LAB strains	32–100	[128]
Zearalenone	<i>L. rhamnosus</i> GG		
	<i>L. rhamnosus</i> LC-705	47–52	[129]
	<i>Lactobacillus</i> spp.	26–69	[130]
Deoxynivalenol	<i>L. paracasei</i>	55	[131]
	<i>L. lactis</i>		
Patulin	LAB strains	13–54	[128]
	<i>L. plantarum</i> GT III	56–66	[132]
	<i>Enterococcus faecium</i> M74 and EF031	41.6–45.3	[119]
Ochratoxins	LAB strains	3–78	[133]
	<i>L. brevis</i> 20023	ND	[134]
	LAB strains	2–96	[133]
	LAB strains	31–57	[135]
	<i>Oenococcus oeni</i>	26–33	[136]
	<i>L. casei</i> LOCK 0920		
	<i>L. brevis</i> LOCK 0944	50	[127]
	<i>L. plantarum</i> LOCK 0945		
	<i>L. acidophilus</i> VM20	95	[133]
	<i>B. animalis</i> VM12		
	<i>Pediococcus parvulus</i>	90	[137]
	<i>L. rhamnosus</i> CECT 278 ^T	97	[138]

Table 2. Cont.

Mycotoxin	Microorganism	Reduction Capacity (%)	References
Degradation			
Aflatoxins	<i>B. subtilis</i>	74	[139]
	<i>B. subtilis</i> BCC42005	45	[114]
Ochratoxins	<i>B. subtilis</i>	92.5	[139]
Zearalenone	<i>B. licheniformis</i>	100	[140]
	<i>B. natto</i>	75	[141]

5. Nutritional Additives and Properties

With the population becoming older, consumers are getting more interested in health issues and big industrial food groups transform their strategy and communication around health [142]. However, putting away compounds that are undesired by some consumers may be difficult and adding some healthy additives is still based on additives. In this section, some examples of use of microorganisms to selectively destroy antinutritional factors or to produce vitamins will be given.

5.1. Cleaning Food of Their Antinutritional Factors (ANF)

Antinutritional factors (ANF) are present in cultivated legumes, seeds and cereals [143]. ANF regroups multiple compounds which are lowering nutritional value of foods by inhibiting protein digestion and nutrient intakes, have deleterious effect on the digestive tract and health or cause gut disorders like flatulence [144,145]. Based on the previous literature, protease inhibitors, tannins, phytic acid are the main molecules responsible for the decreasing of proteolytic activity due to the inactivation of gut protease and denaturation of protein (protease inhibitors and tannins respectively) and the capture of positive-charged mineral ions (phytic acid). Lectins are glycoproteins characterised by their ability to interfering with intestinal epithelium leading to inflammatory state and a lack of nutrient absorption. Flatulence is linked to the digestion of α -galactosides like raffinose, stachyose and verbascose by the microbiota. The development of legume-based diet as protein source and the demands for healthy product poses the challenge for developing processes that keep nutritional benefits and clear products from ANF. First approach consisting in thermal processes as boiling, microwaving or pressurised cooking, such processes have shown great efficiency for decreasing trypsin inhibitors, phytic acid, hemagglutinins activity (lectins), saponins and some oligosaccharides of chickpeas [146]. The second approach is based on the supplementation of the cooking by germination or fermentation. The germination of seeds has shown significant results by eliminating flatulence-linked oligosaccharides [147] and decreasing the level of phytic acid, tannins and trypsin inhibitors [148]. The combination between germination and cooking allows us to significantly decrease or eliminate ANF in seeds and cereals. Nevertheless, few legume-based foods are produced following the germination process. Fermentation could appear as a safe way to tackles ANF from ungerminated legumes. Lactic acid fermentation by *L. plantarum* on bean flour shown multiple effects on ANF, such as the elimination of oligosaccharides and a significant diminution of lectins level [149]. The fermentation by *L. brevis* also shown great improvement on soybean digestibility due to the reduction of protease inhibitors and oligosaccharides [150]. Significant decrease of raffinose, stachyose, trypsin inhibitors and tannins have been reported for lactic acid fermentation of black bean by *L. casei* and *L. plantarum* [151]. Similar results have been reported for lactic acid fermentation of pearl millet [152]. Fungi fermentation can also eliminate ANF, and *Rhizopus oligosporu* has shown significant activity against oligosaccharides and protease inhibitors [147]. But the fungi fermentation must be well characterised to avoid the production of any toxic compounds. As reported by the literature, fermentation could help to reduce or eliminate some ANF without using heavy processes or chemical treatments. It can be used on raw products or at further stage of transformation. More

investigations are needed due to the variability of fermentation effects caused by strains and legumes' specificity. Indeed, lactic acid fermentation of plant-based product could lead to the production of biogenic amines [153], and this production is hugely dependent on the strains and the variety of legumes. The combination of thermic processes, germination and fermentation seems to be a great way for improving nutritional quality of plant-based product, but studies must be carried out to avoid any deleterious effects. Characterisation of plant cultivars composition and the activity of microorganism on it is the only way to develop clean and healthy plant-based products.

5.2. Vitamins Like Folate

Vitamins are organic compounds involved in several metabolic functions including energy production, red blood cell synthesis, etc. They are grouped into 2 main groups: lipid-soluble (vitamins A, D, E, K) and water-soluble (vitamin C and eight kinds of B vitamins) vitamins [154].

Vitamins of group A comprise retinoids, retinol, retinal, retinoic acid and retinyl esters. Pro-vitamin A is composed of various carotenoids (β -carotene, α -carotene, and β -cryptoxanthin), which are then converted in their active forms in the body [154].

Vitamin D derives from cholesterol and ergosterol. Cholesterol is converted into 7-dehydrocholesterol, which can be cleaved by ultraviolet (UV)-radiation to form cholecalciferol (vitamin D3), while ergosterol results in ergocalciferol (vitamin D2). Vitamins D2 and D3, used by humans, require further hydroxylations [154].

The vitamin E group is formed by different chemical forms: four tocopherol and four tocotrienol forms. Tocopherols are often used as dietary supplements for humans, food preservatives, and in manufacture of cosmetics and sunscreens. However, α -tocopherol is the most predominant and active form in most human and animal tissues [155].

Vitamin K can be divided into phyloquinone (vitamin K1) with a phytyl group obtained from plants and menaquinones (vitamin K2) [154]. Vitamin C or ascorbic acid is an essential dietary component that humans are unable to synthesize.

B vitamins contain thiamine (vitamin B1), riboflavin (vitamin B2), niacin (vitamin B3), pantothenic acid (vitamin B5), vitamin B6, biotin (vitamin B7 or vitamin H), folic acid (vitamin B9), and cobalamin (vitamin B12) [154].

The absence of adequate amounts of these compounds in the diet can cause several health problems not only to humans but also to animals. Therefore, they are produced industrially and used widely not only as food and feed additives, but also as cosmetics, therapeutic agents and health and technical aids [154]. However, these processes require the use of solvents, which are undesirable pollutants harmful to the environment. To overcome this drawback several studies are focused on the selection of microorganisms able to produce vitamins (Table 3).

Table 3. Vitamins, their functions and microorganisms used for their production.

Vitamin	Physiological Functions	Microbial Producer	References
Vitamin A	Immune system regulation, vision, reproduction, cellular communication, cell growth and differentiation.	<i>Cyberlindnera jadinii</i> (teleomorph <i>Candida utilis</i>), <i>Saccharomyces cerevisiae</i> , <i>Pichia pastoris</i> , <i>Y. lipolytica</i>	[156–160]
Vitamin D	Calcium absorption and mineralization of bones, modulation of cell growth, neuromuscular, immune and inflammation functions	<i>S. cerevisiae</i> , <i>Saccharomyces uvarum</i> and <i>Cyberlindnera jadinii</i> (teleomorph <i>C. utilis</i>)	[161]
Vitamin E	Antioxidant activity, cellular membrane stabilizer	Microalgae: <i>Spirulina platensis</i> , <i>Dunaliella tertiolecta</i> , <i>Synechocystis</i> spp., <i>Nannochloropsis oculata</i> , <i>Tetraselmis suecica</i> , <i>Chlorella</i> spp., <i>Clamydomonas</i> spp., and <i>Ochromonas</i> spp., <i>Euglena gracilis</i> , <i>Dunaliella salina</i> , <i>Isochrysis galbana</i> , and <i>Diacronema vlkianum</i>	[155]

Table 3. Cont.

Vitamin	Physiological Functions	Microbial Producer	References
Vitamin K	Blood coagulation	<i>Flavobacterium</i> sp., <i>B subtilis</i> , and <i>Propionibacterium freudenreichii</i>	[162–164]
B vitamins	Energy production, red blood cell synthesis	<i>B. subtilis</i> , <i>Corynebacterium ammoniagenes</i> , <i>L. plantarum</i> , <i>Leuconostoc mesenteroides</i> , <i>Lactococcus lactis</i> , <i>Rhodococcus rhodochrous</i> , <i>Agrobacterium</i> sp., <i>Corynebacterium glutamicum</i> , <i>Flavobacterium</i> sp., <i>Sinorhizobium meliloti</i> (ex <i>Rhizobium meliloti</i>), <i>B. sphaericus</i> , <i>Serratia marcescens</i> , <i>Propionibacterium shermanii</i> , <i>Pseudomonas denitrificans</i> , <i>Bacillus megaterium</i> , <i>Methanobacterium ivanovii</i> , <i>Rhodobacter capsulatus</i> , <i>Ashbya gossypii</i> , <i>Candida parapsilosis</i> , <i>Candida flaueri</i> and <i>Candida famata</i> (teleomorph <i>Debaryomyces hansenii</i>).	[165]
Vitamin C	Antioxidant activity, biosynthesis of collagen, l-carnitine and certain neurotransmitters, protein metabolism	<i>Gluconobacter</i> spp., <i>Acetobacter</i> spp., <i>Ketogulonicigenium</i> spp., <i>Pseudomonas</i> spp., <i>Erwinia</i> spp., and <i>Corynebacterium</i> spp.	[166,167]

Presently, several studies are focusing on Vitamin B9 or folate since it plays very important functions in human health including amino acid metabolism and DNA replication and repair and is thus essential for cell division. In pregnant women daily intake of folic acid is recommended since it reduces the risk of low birth weight, maternal anemia and neural tube defects (NTD): spina bifida and anencephaly [168]. There are many forms of vitamin B9, called vitamers, which are more resistant to technological processes. Folic acid, the synthetic form of B9 vitamin, presents only a glutamate molecule, while naturally occurring forms are characterized by a polyglutamate chain. In addition, folic acid exhibits a fully oxidized pteridine ring, while the other vitamers are generally either partially reduced (at the 7,8-position) in the case of dihydrofolate forms, or fully reduced (at the 5,6,7,8-position) in the case of tetrahydrofolate compounds [169].

Humans do not synthesize folate de novo and folate deficiency represents a problem worldwide. In fact, several countries adopted mandatory fortification programs in foods of mass consumption such as flours and rice [169]. The main strategies used to address the problem of vitamin deficiencies are (i) supplementation, (ii) food fortification, and (iii) dietary diversification [170]. Unfortunately, folate-rich foods are not always available, depending on the season, and on the geographic, agro-ecological and socio-economic context, and the intake of folic acid could exert some adverse secondary effects, such as masking symptoms of vitamin B12 deficiency and possibly promoting colorectal cancer. These side effects are not observed when natural folates, such as those found in foods or produced by certain microorganisms, are consumed [169].

The main producers of folate are LAB and bifidobacteria (Table 4). Folate production is strain-dependent and is influenced by growth kinetics and medium composition. Several studies reviewed in [169] highlighted that folate bacterial production occurs during the exponential growth phase or at the beginning of the stationary phase and is then consumed.

Table 4. Main bacterial species producing folate.

Microorganism	Outcome Range (ng/mL)	References
Bifidobacteria		
<i>Bifidobacterium (B.) adolescentis</i>	50–150	[171]
<i>B. dentium</i>	0–25	
<i>B. animalis</i>	1–65	[172]
<i>B. bifidum</i>	26	
<i>B. breve</i>	1	
<i>B. catenulatum</i>	1–3	
<i>B. longum</i>	3	
<i>B. pseudocatenulatum</i>	29	
<i>B. adolescentis</i>	10–30	[173]
Lactic acid bacteria		
<i>Lactobacillus acidophilus</i>	0–38	[174]
<i>Lb. amylovorus</i>	75–87	
<i>L. casei</i>	0–2	
<i>L. paracasei</i>	0–40	
<i>Levilactobacillus brevis</i> (ex <i>Lb. brevis</i>)	0–150	[175]
<i>Latilactobacillus curvatus</i> (ex <i>Lb. curvatus</i>)	0–20	
<i>Fructilactobacillus fructivorans</i> (ex <i>Lb. fructivorans</i>)	0–20	
<i>Lb. helveticus</i>	2–89	
<i>Limosilactobacillus reuteri</i> (ex <i>Lb. reuteri</i>)	0–125	
<i>Loigolactobacillus coryniformis</i> (ex <i>Lb. coryniformis</i>)	80–100	[176]
<i>L. pentosus</i>	0–4	
<i>Lb. sakei</i>	101–107	
<i>Pediococcus. parvulus</i>	40–60	
<i>Pediococcus pentosaceus</i>	0–40	
<i>Weissella confusa</i>	0–20	
<i>Lb. delbrueckii</i>	50–200	[171]
<i>L. plantarum</i>	36–60	
<i>L. fermentum</i>	0–148	[177]
<i>Lb. johnsonii</i>	28	[178]
<i>Lactococcus lactis</i>	57–291	[179]
<i>Leuconostoc lactis</i>	45	
<i>Leuconostoc paramesenteroides</i>	44	
<i>S. thermophilus</i>	0–170	[180]

The majority of studies concerning folate production by eukaryotic microorganisms were carried out on *S. cerevisiae* and *A. gossypii* [173]. However, also other yeast genera are reported as folate producers such as *Candida*, *Debaryomyces*, *Kodamea*, *Metchnikowia*, *Wickerhamiella* [174]. *A. gossypii* can naturally synthesize 40 µg/L of folates and after metabolic engineering is able to reach 6595 µg/L. This result was obtained overexpressing 3 genes involved in folate production (*FOL1*, *FOL2*, *FOL3*) and deleting the gene *MET7* which encodes for a FPGS (folypolyglutamate synthetase) which catalyses the polyglutamylation of folates in their gamma-carboxyl residue [173]. The elimination of competing pathways, such as riboflavin and adenine favours folate production [173].

Despite the efforts undertaken so far, microbial folate production is still low and not competitive in terms of cost and final concentration with industrial processes. A possibility to increase folate production could be the development of co-cultures of folate producing strains or folate vitamers that are resistant to oxidation, acid pH, and heat treatments. Finally, the possibility to use probiotic strains could be an advantage since folate could be produced in the gut. Future research should also focus on the understanding the complex regulatory mechanisms governing the enzymatic activities involved in the folate pathway; the optimization of the fermentation conditions and further development of downstream processes for the recovery and purification of the product.

6. Use of Taste-Active Microbial Amino Acids, and Peptides in Food Fermentation

Eventually, we will see some examples concerning inactive microorganisms that can be used for some compounds active for food properties.

Salt is an irreplaceable additive, flavouring foods. Culinary salt is a chemical compound consisting of the elements sodium and chlorine. Salty taste is given mainly by Na^+ . The ions of the alkaline metal group exhibit also a salty taste but causing less feeling than Na^+ . The size of the ions Li^+ and K^+ is also close to that of Na^+ , creating a salty taste that is almost similar. The salinity of substances is assessed in comparison to the sodium chloride standard [181,182]. KCl is the main ingredient used to replace salt with an index of 0.6 (when the salinity of NaCl is 1).

Monosodium glutamate (MSG) gives the taste of meat and umami, which is one of the five basic tastes with sourness, sweetness, saltiness, bitterness. In 1909, Kikunae Ikeda discovered MSG from seaweed. The taste strength of glutamate is quite strong. The sensory threshold of MSG is 1/3000 (one gram over three liters of water). This intensity is much stronger than salt and sugar. However, in addition, glutamate enhances also the perception of salty taste, and helps therefore to reduce the amount of salt added to food. Reducing salt is a goal in daily meals for humans to avoid certain diseases such as high blood pressure, kidney failure. But reducing salt will lead to food with poor taste. Using KCl as a substitute for culinary salt will create a bitter and metallic taste. Research results have shown that MSG combines culinary salt, significantly improving the sensory properties of foods. Yamaguchi [183,184] reported that the addition of MSG to broth could help to decrease the rate of sodium chloride for a similar sensorial result. Thus, MSG can replace culinary salt while ensuring the deliciousness of food.

MSG is present in different amounts in most natural food sources such as tomatoes, fish meat or oysters. It can be present as a free form or bind with other amino acids to create certain peptides and proteins. The content of MSG in nature has been determined [185,186]. The highest content of free glutamate in food (100 g) are found in Pamesano cheese, 1.680 mg; seaweed, 1.608 mg, oyster, 140 mg; tomatoes, 246 mg, or Japanese fish sauce, 1.323 mg.

In the human body, approximately 70% of body weight is water, 20% is protein and of which glutamate accounts for about 2%. MSG is a natural part of metabolism and about 50 g per day is formed by the human body. The average person consumes 10–20 g of bound glutamate per day and about 1 g of free food glutamate. Daily intake of glutamate is the main source of intestinal energy.

Saccharomyces yeast is a rich-in-protein source (protein content accounts for 48–50% dry matter) and yeast hydrolysed products are considered as rich sources of amino acids and peptides. They have many applications in food such as salad dressings, ice creams, crackers or meat products. They are used as additives, enhancing the flavour of the food products. Beer production can be a source of yeast. For instance, in a country like Vietnam with beer consumption of about 4.6 billion liters in 2019 according to data from the World Bank and Euromonitor, the production can generate around 7000 tons of spent yeast that can be used for either food consumption and feed. Utilising a large source of protein from brewer's yeast to produce hydrolysed products for application in food and food additives has a high real-life benefit. The composition of some amino acids in the brewer's yeast hydrolysates (BYH) varies depending on hydrolysis techniques. Continuous circulation hydrolysis method with heat shock and processed by autolysis gives the highest total amino acid content. The glutamate content accounts for 3.14 g/ 100 g BYH (55% dry matter) when the total amino acid composition achieved 32.3 g/100 g BYH.

However, bitterness in hydrolysates is one of the major undesirable aspects for various applications in food processing. It has been reported that the bitterness of brewer's yeast hydrolysate obtained by using flavourzyme is the lowest and that this product keeps a good umami taste [187].

The second limitation in the use of yeast and hydrolysate is the high content of nucleic acid in the yeast. There are many methods for reducing or separating nucleic acids in hydrolysed products such as extracellular ribonuclease enzymes, chemical agents, thermal

shock and autolysis. Using extracellular ribonuclease enzyme for hydrolysis of nucleic acid gives good efficiency but suffers high production cost. Chemical agents negatively affect the quality of the hydrolysed products used in the food industry. It has been reported that a method using combination of heat shock treatment, autolysis and continuous circulation hydrolysis techniques gave the smallest content of nucleic acid in the brewer's yeast hydrolysate in comparison with using the batch and continuous overflow process [188].

In addition to the contribution of inactivated yeast to the taste of products, this popular microorganism can also bring health-active compounds. One of the most economically important components of yeast biomass is ergosterol, which, as already discussed in the previous paragraph, could be used as a precursor of vitamin D2 and another sterol drug [189]. Thanks to advanced technology in biotechnology, modified strains of yeast have been developed to enhance the production of ergosterol or the co-production of ergosterol with other products [190–192]. In Vietnam, the National Institute of Nutrition has conducted investigation on the production of ergosterol from *S. cerevisiae* and its application in functional food production. From 50 yeast samples of bakerhouses and 50 samples of fresh grapefruits from markets in Hanoi, two yeast strains, namely MB14.2.2 and N42.2.2, were found with the highest concentration of ergosterol in comparison with dry biomass (3.7% and 3.5%, respectively). Furthermore, optimized conditions and apparatus system for ergosterol production from these strains were established. For the applications in function foods, cookies (for children) and soya milk powder (for adults) were supplemented with vitamin D2 (1600 IU/100 g and 2261 UI/100 g in cookies and soya milk powder, respectively), that was transformed from ergosterol using radiation method. After using the products, the group of children had better transformation of the z-score index height/age and body mass index (BMI). The adult group improved bone health and improved blood biochemical indicators. Concentrations of 25- (OH) D of both groups with vitamin D2 were significantly higher than that of the control group ($p < 0.001$). The percentage of vitamin D deficiency noticeably decreased in both intervention groups.

Furthermore, brewing yeast is a great source for β -glucan. When yeasts are grown for seasoning purposes, molasses from sugar production is used as raw material for yeast fermentation. Presently, there are three products: spray-dried whole cell yeasts, yeast extract in paste form and spray dried yeast extract. The yeast cell wall separated after centrifuge goes to wastewater and causes complications and costs in wastewater treatment. Therefore, there would be a great opportunity to add value to yeast by using the cell wall as a source for production of β -glucan, a functional food.

7. Conclusions

With the growing concern of consumers towards the food that they eat, the clean label strategy has been generalised in many companies. From the first efforts which could often be assimilated to green washing, some companies have now developed a systematic struggle against additives. In this cleaning effort, microorganisms can be an efficient tool. This review illustrates what microorganisms can bring to the clean label concept through examples of recent strategies. In fact, besides the use of microorganisms producing antifungal weak acids in bread products, exopolysaccharides or of strains able to consume lipids or sugars to decrease the caloric properties of foods, or compounds with a positive effect on human effects, the efficacy of microbial strains to obtain good foods without additives is always subject to evaluation. The use of microorganisms could be useful to reduce the employment of additives since some strains are able to transform food components, degrade off-flavors, antinutritional factors, toxins, and chemical pollutants, or bring new molecules that are active for taste or health. Further studies are necessary to improve this “clean label” approach to reduce the list of ingredients used in food products.

Author Contributions: All authors have participated to the redaction of the manuscript. All authors have read and agreed to the published version of the manuscript.

Funding: This research was funded by the BUALUANG Chair Professorship of the University of Thammasat.

Acknowledgments: Acknowledgements are given to Center of Excellence in Food Science and Innovation, Faculty of Science and Technology of Thammasat University, and Institute for Global Food Security, School of Biological Science, Queen's University Belfast for the support.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Wang, S.; Li, C.; Copeland, L.; Niu, Q.; Wang, S. Starch Retrogradation: A Comprehensive Review. *Compr. Rev. Food Sci. Food Saf.* **2015**, *14*, 568–585. [[CrossRef](#)]
2. Goesaert, H.; Brijs, K.; Veraverbeke, W.S.; Courtin, C.M.; Gebruers, K.; Delcour, J.A. Wheat Flour Constituents: How They Impact Bread Quality, and How to Impact Their Functionality. *Trends Food Sci. Technol.* **2005**, *1*, 12–30. [[CrossRef](#)]
3. Fickers, P.; Benetti, P.H.; Waché, Y.; Marty, A.; Mauersberger, S.; Smit, M.S.; Nicaud, J.M. Hydrophobic Substrate Utilisation by the Yeast *Yarrowia lipolytica*, and Its Potential Applications. *FEMS Yeast Res.* **2005**, *5*, 527–543. [[CrossRef](#)] [[PubMed](#)]
4. Fickers, P.; Marty, A.; Nicaud, J.M. The Lipases from *Yarrowia lipolytica*: Genetics, Production, Regulation, Biochemical Characterization and Biotechnological Applications. *Biotechnol. Adv.* **2011**, *29*, 632–644. [[CrossRef](#)] [[PubMed](#)]
5. Smit, M.; Mokgoro, M.; Setati, M.; Nicaud, J.-M. Preparation of Dodecanol-Tolerant Strains of *Yarrowia lipolytica*. *Biotechnol. Lett.* **2004**, *26*, 849–854. [[CrossRef](#)]
6. Romero-Guido, C.; Belo, I.; Ta, T.M.; Cao-Hoang, L.; Alchihab, M.; Gomes, N.; Thonart, P.; Teixeira, J.A.; Destain, J.; Waché, Y. Biochemistry of Lactone Formation in Yeast and Fungi and Its Utilisation for the Production of Flavour and Fragrance Compounds. *Appl. Microbiol. Biotechnol.* **2011**, *89*, 535–547. [[CrossRef](#)] [[PubMed](#)]
7. Groenewald, M.; Boekhout, T.; Neuvéglise, C.; Gaillardin, C.; van Dijk, P.W.; Wyss, M. *Yarrowia lipolytica*: Safety Assessment of an Oleaginous Yeast with a Great Industrial Potential. *Crit. Rev. Microbiol.* **2014**, *40*, 187–206. [[CrossRef](#)] [[PubMed](#)]
8. Davidou, S.; Le Meste, M.; Debever, E.; Bekaert, D. A Contribution to the Study of Staling of White bread: Effect of Water and Hydrocolloid. *Food Hydrocoll.* **1996**, *10*, 375–383. [[CrossRef](#)]
9. Zhou, Y.; Cui, Y.; Qu, X. Exopolysaccharides of Lactic Acid Bacteria: Structure, Bioactivity and Associations: A Review. *Carbohydr. Polym.* **2019**, *207*, 317–332. [[CrossRef](#)]
10. Sandra, G.; Schwab, C.; Bello, F.D.; Coffey, A.; Gänzle, M.; Arendt, E. Comparison of the Impact of Dextran and Reuteran on the Quality of Wheat Sourdough Bread. *J. Cereal Sci.* **2012**, *56*, 531–537. [[CrossRef](#)]
11. Zhang, Y.; Guo, L.; Li, D.; Jin, Z.; Xu, X. Roles of Dextran, Weak Acidification and Their Combination in the Quality of Wheat Bread. *Food Chem.* **2019**, *286*, 197–203. [[CrossRef](#)]
12. Csáki, K.F.; Sebestyén, É. Who Will Carry Out the Tests That Would be Necessary for Proper Safety Evaluation of Food Emulsifiers? *Food Sci. Hum. Well.* **2019**, *8*, 126–135. [[CrossRef](#)]
13. Ozturk, B.; McClements, D.J. Progress in Natural Emulsifiers for Utilization in Food Emulsions. *Curr. Opin. Food Sci.* **2016**, *7*, 1–6. [[CrossRef](#)]
14. Ayudhya, M.S.N.; Chuenchomrat, P. Screening and Characterization of Bacterial Biosurfactant from Bangkok and Vicinities. *Int. J. Life Sci. Biotechnol. Pharma Res.* **2015**, *4*, 168–171.
15. Choopraserdchok, T.; Athinuwat, D.; Chuenchomrat, P. Effects of Glucose and Ferrous Supplements and Culture Conditions on Lipopeptide Biosurfactant from *Pseudomonas* spp. *Int. J. Environ. Sci. Dev.* **2018**, *8*, 772–775. [[CrossRef](#)]
16. Katemai, W.; Maneerat, S.; Kawai, F.; Kanzaki, H.; Nitoda, T.; H-kittikun, A. Purification and Characterization of a Biosurfactant Produced by *Issatchenkia orientalis* SR4. *J. Gen. Appl. Microbiol.* **2008**, *54*, 79–82. [[CrossRef](#)] [[PubMed](#)]
17. Nitschke, M.; Costa, S.G.V.A.O. Biosurfactants in Food Industry. *Trends Food Sci. Technol.* **2007**, *18*, 252–259. [[CrossRef](#)]
18. Hemlata, B.; Selvin, J.; Tukaram, K. Optimization of Iron Chelating Biosurfactant Production by *Stenotrophomonas maltophilia* NBS-11. *Biocatal. Agric. Biotechnol.* **2015**, *4*, 135–143. [[CrossRef](#)]
19. Zhao, F.; Shi, R.; Ma, F.; Han, S.; Zhang, Y. Oxygen Effects on Rhamnolipids Production by *Pseudomonas aeruginosa*. *Microb. Cell Fact.* **2018**, *17*, 39. [[CrossRef](#)] [[PubMed](#)]
20. Elshafie, A.E.; Joshi, S.J.; Al-Wahaibi, Y.M.; Al-Bemani, A.S.; Al-Bahry, S.N.; Al-Maqbali, D.A.; Banat, I.M. Sophorolipids Production by *Candida bombicola* ATCC 22214 and Its Potential Application in Microbial Enhanced Oil Recovery. *Front. Microbiol.* **2015**, *6*, 1–11. [[CrossRef](#)] [[PubMed](#)]
21. Zaragoza, A.; Teruel, J.A.; Aranda, F.J.; Ortiz, A. Interaction of a Trehalose Lipid Biosurfactant Produced by *Rhodococcus erythropolis* 51T7 with a Secretory Phospholipase A2. *J. Colloid Interface Sci.* **2013**, *408*, 132–137. [[CrossRef](#)] [[PubMed](#)]
22. Santos, D.K.F.; Rufino, R.D.; Luna, J.M.; Santos, V.A.; Sarubbo, L.A. Biosurfactants: Multifunctional Biomolecules of the 21st Century. *Int. J. Mol. Sci.* **2016**, *17*, 401. [[CrossRef](#)]
23. Kuiper, I.; Legendijk, E.L.; Pickford, R.; Derrick, J.P.; Lamers, G.E.; Thomas-Oates, J.E.; Lugtenberg, B.J.; Bloemberg, G.V. Characterization of Two *Pseudomonas putida* Lipopeptide Biosurfactants, Putisolvin I and II, which Inhibit Biofilm Formation and Break Down Existing Biofilms. *Mol. Microbiol.* **2004**, *51*, 97–113. [[CrossRef](#)]
24. Vedaraman, N.; Venkatesh, N. Production of Surfactin by *Bacillus subtilis* MTCC 2423 from Waste Frying Oils. *Braz. J. Chem. Eng.* **2011**, *28*, 175–180. [[CrossRef](#)]

25. Biniarz, P.; Baranowska, G.; Feder-Kubis, J.; Krasowska, A. The Lipopeptides Pseudofactin II and Surfactin Effectively Decrease *Candida albicans* adhesion and hydrophobicity. *Antonie Van Leeuwenhoek* **2015**, *108*, 343–353. [[CrossRef](#)] [[PubMed](#)]
26. Chan, X.Y.; Chang, C.Y.; Hong, K.W.; Tee, K.K.; Yin, W.F.; Chan, K.G. Insights of Biosurfactant Producing *Serratia marcescens* strain W2.3 Isolated from Diseased Tilapia fish: A Draft Genome Analysis. *Gut Pathog.* **2013**, *5*, 29. [[CrossRef](#)]
27. Shi, J.; Zhu, X.; Lu, Y.; Zhao, H.; Lu, F.; Lu, Z. Improving Iturin A Production of *Bacillus amyloliquefaciens* by Genome Shuffling and Its Inhibition Against *Saccharomyces cerevisiae* in Orange Juice. *Front. Microbiol.* **2018**, *9*, 2683. [[CrossRef](#)]
28. Hanif, A.; Zhang, F.; Li, P.; Li, C.; Xu, Y.; Zubair, M.; Zhang, M.; Jia, D.; Zhao, X.; Liang, J.; et al. Fengycin Produced by *Bacillus amyloliquefaciens* FZB42 Inhibits *Fusarium graminearum* Growth and Mycotoxins Biosynthesis. *Toxins* **2019**, *11*, 295. [[CrossRef](#)]
29. Campos, J.M.; Stamford, T.L.; Sarubbo, L.A.; de Luna, J.M.; Rufino, R.D.; Banat, I.M. Microbial Biosurfactants as Additives for Food Industries. *Biotechnol. Prog.* **2013**, *29*, 1097–1108. [[CrossRef](#)]
30. Passos da Silva, D.; Schofield, M.C.; Parsek, M.R.; Tseng, B.S. An Update on the Sociomicrobiology of Quorum Sensing in Gram-Negative Biofilm Development. *Pathogens* **2017**, *6*, 51. [[CrossRef](#)] [[PubMed](#)]
31. Pacwa-Płociniczak, M.; Płaza, G.A.; Piotrowska-Seget, Z.; Cameotra, S.S. Environmental Applications of Biosurfactants: Recent Advances. *Int. J. Mol. Sci.* **2011**, *12*, 633–654. [[CrossRef](#)] [[PubMed](#)]
32. Mouafo, T.H.; Mbawala, A.; Ndjouenkeu, R. Effect of Different Carbon Sources on Biosurfactants' Production by Three Strains of *Lactobacillus* spp. *Biomed. Res. Int.* **2018**, *2018*, 5034783. [[CrossRef](#)] [[PubMed](#)]
33. Lvaresou, A.; Iakovou, K. Biosurfactants in Cosmetics and Biopharmaceuticals. *Lett. Appl. Microbiol.* **2015**, *61*, 214–223. [[CrossRef](#)]
34. Peng, F.; Liu, Z.; Wang, L.; Shao, Z. An Oil-Degrading Bacterium: *Rhodococcus erythropolis* Strain 3C-9 and Its Biosurfactants. *J. Appl. Microbiol.* **2007**, *102*, 1603–1611. [[CrossRef](#)] [[PubMed](#)]
35. Ly, M.H.; Vo, N.H.; Le, T.M.; Belin, J.M.; Waché, Y. Diversity of the Surface Properties of *Lactococci* and Consequences on Adhesion to Food Components. *Colloids Surf. B Biointerfaces* **2006**, *52*, 149–153. [[CrossRef](#)] [[PubMed](#)]
36. Ly, M.H.; Aguedo, M.; Goudot, S.; Le, M.L.; Cayot, P.; Teixeira, J.A.; Le, T.M.; Belin, J.M.; Waché, Y. Interactions between Bacterial Surfaces and Milk Proteins, Impact on Food Emulsions Stability. *Food Hydrocoll.* **2008**, *22*, 742–751. [[CrossRef](#)]
37. Ly, M.H.; Covarrubias-Cervantes, M.; Dury-Brun, C.; Bordet, S.; Voilley, A.; Le, T.M.; Belin, J.M.; Waché, Y. Retention of Aroma Compounds by Lactic Acid Bacteria in Model Food Media. *Food Hydrocoll.* **2008**, *22*, 211–217. [[CrossRef](#)]
38. Ma, G.; Zhang, L.; Sugiura, M.; Kato, M. Chapter 24—Citrus and Health, in *The Genus Citrus*; Talon, M., Caruso, M., Gmitter, F.G., Eds.; Woodhead Publishing (Elsevier Inc.): Cambridge, UK, 2020; pp. 495–511.
39. Yusof, S.; Ghazali, H.M.; King, G.S. Naringin Content in Local Citrus Fruits. *Food Chem.* **1990**, *37*, 113–121. [[CrossRef](#)]
40. Giannuzzo, A.N.; Boggetti, H.J.; Nazareno, M.A.; Mishima, H.T. Supercritical Fluid Extraction of Naringin from the Peel of *Citrus paradisi*. *Phytochem. Anal.* **2003**, *14*, 221–223. [[CrossRef](#)]
41. Singh, S.V.; Gupta, A.K.; Jain, R.K. Adsorption of Naringin on Nonionic (Neutral) Macroporous Adsorbent Resin from Its Aqueous Solutions. *J. Food Eng.* **2008**, *86*, 259–271. [[CrossRef](#)]
42. Ribeiro, M.H. Naringinases: Occurrence, Characteristics, and Applications. *Appl. Microbiol. Biotechnol.* **2011**, *90*, 1883–1895. [[CrossRef](#)] [[PubMed](#)]
43. Busto, M.D.; Cavia-Saiz, M.; Ortega, N.; Muñoz, P. Chapter 20—Enzymatic Debittering on Antioxidant Capacity of Grapefruit Juice. In *Processing and Impact on Antioxidants in Beverages*; Preedy, V., Ed.; Academic Press: San Diego, CA, USA, 2014; pp. 195–202.
44. Puri, M.; Seth, M.; Marwaha, S.; Kothari, R. Debittering of Kinnow Mandarin Juice by Covalently Bound Naringinase on Hen Egg White. *Food Biotechnol.* **2007**, *15*, 13–23. [[CrossRef](#)]
45. Puri, M.; Banerjee, U.C. Production, Purification, and Characterization of the Debittering Enzyme Naringinase. *Biotechnol. Adv.* **2000**, *18*, 207–217. [[CrossRef](#)]
46. Mendoza-Cal, A.; Cuevas-Glory, L.; Lizama-Uc, G.; Ortiz-Vázquez, E. Naringinase Production from Filamentous Fungi Using Grapefruit Rind in Solid State Fermentation. *Afr. J. Microbiol. Res.* **2010**, *4*, 1964–1969.
47. Chen, Y.; Ni, H.; Chen, F.; Cai, H.; Li, L.; Su, W. Purification and Characterization of a Naringinase from *Aspergillus aculeatus* JMUb058. *J. Agric. Food Chem.* **2013**, *61*, 931–938. [[CrossRef](#)]
48. Weiz, G.; Breccia, J.D.; Mazzaferro, L.S. Screening and Quantification of the Enzymatic Deglycosylation of the Plant Flavonoid Rutin by UV-Visible Spectrometry. *Food Chem.* **2017**, *229*, 44–49. [[CrossRef](#)]
49. Yadav, M.; Sehrawat, N.; Sharma, A.; Kumar, V.; Kumar, A. Naringinase: Microbial Sources, Production and Applications in Food Processing Industry. *J. Microbiol. Biotechnol. Food Sci.* **2018**, *8*, 717–720. [[CrossRef](#)]
50. Koseki, T.; Mese, Y.; Nishibori, N.; Masaki, K.; Fujii, T.; Handa, T.; Yamane, Y.; Shiono, Y.; Murayama, T.; Iefuji, H. Characterization of an Alpha-L-Rhamnosidase from *Aspergillus kawachii* and Its Gene. *Appl. Microbiol. Biotechnol.* **2008**, *80*, 1007–1013. [[CrossRef](#)]
51. Chang, H.-Y.; Lee, Y.-B.; Bae, H.-A.; Huh, J.-Y.; Nam, S.-H.; Sohn, H.-S.; Lee, H.J.; Lee, S.-B. Purification and Characterisation of *Aspergillus sojae* Naringinase: The Production of Prunin Exhibiting Markedly Enhanced Solubility with in vitro Inhibition of HMG-CoA Reductase. *Food Chem.* **2011**, *124*, 234–241. [[CrossRef](#)]
52. Srikantha, K.; Kapilan, R.; Vasantharuba, S. Characterization of Best Naringinase Producing Fungus Isolated from the Citrus Fruits. *Int. J. Biol. Res.* **2016**, *4*, 83. [[CrossRef](#)]
53. Hashimoto, W.; Miyake, O.; Nankai, H.; Murata, K. Molecular Identification of an Alpha-L-Rhamnosidase from *Bacillus* sp Strain GL1 as an Enzyme Involved in Complete Metabolism of Gellan. *Arch. Biochem. Biophys.* **2003**, *415*, 235–244. [[CrossRef](#)]
54. Mukund, P.; Belur, P.D.; Saidutta, M.B. Production of Naringinase from a New Soil Isolate, *Bacillus methylotrophicus*: Isolation, Optimization and Scale-up Studies. *Prep. Biochem. Biotechnol.* **2014**, *44*, 146–163. [[CrossRef](#)]

55. Zhu, Y.; Jia, H.; Xi, M.; Xu, L.; Wu, S.; Li, X. Purification and Characterization of a Naringinase from a Newly Isolated Strain of *Bacillus amyloliquefaciens* 11568 Suitable for the Transformation of Flavonoids. *Food Chem.* **2017**, *214*, 39–46. [[CrossRef](#)] [[PubMed](#)]
56. Beekwilder, J.; Marcozzi, D.; Vecchi, S.; de Vos, R.; Janssen, P.; Francke, C.; van Hylckama Vlieg, J.; Hall, R.D. Characterization of Rhamnosidases from *Lactobacillus plantarum* and *Lactobacillus acidophilus*. *Appl. Environ. Microbiol.* **2009**, *75*, 3447–3454. [[CrossRef](#)] [[PubMed](#)]
57. Kaur, A.; Singh, S.; Singh, R.S.; Schwarz, W.H.; Puri, M. Hydrolysis of Citrus Peel Naringin by Recombinant α -L-rhamnosidase from *Clostridium stercorarium*. *J. Chem. Technol. Biotechnol.* **2010**, *85*, 1419–1422. [[CrossRef](#)]
58. Miake, F.; Satho, T.; Takesue, H.; Yanagida, F.; Kashige, N.; Watanabe, K. Purification and Characterization of Intracellular α -l-rhamnosidase from *Pseudomonas paucimobilis* FP2001. *Arch. Microbiol.* **2000**, *173*, 65–70. [[CrossRef](#)] [[PubMed](#)]
59. Iqbal, A.; Khalil, I.A.; Ateeq, N.; Sayyar Khan, M. Nutritional Quality of Important Food Legumes. *Food Chem.* **2006**, *97*, 331–335. [[CrossRef](#)]
60. Kaczmarek, K.T.; Chandra-Hioe, M.V.; Frank, D.; Arcot, J. Aroma Characteristics of Lupin and Soybean after Germination and Effect of Fermentation on Lupin Aroma. *LWT* **2018**, *87*, 225–233. [[CrossRef](#)]
61. Kaneko, S.; Kumazawa, K.; Nishimura, O. Studies on the Key Aroma Compounds in Soy Milk Made from Three Different Soybean Cultivars. *J. Agric. Food Chem.* **2011**, *59*, 12204–12209. [[CrossRef](#)]
62. Stephany, M.; Kapusi, K.; Bader-Mittermaier, S.; Schweiggert-Weisz, U.; Carle, R. Odour-Active Volatiles in Lupin Kernel Fibre Preparations (*Lupinus angustifolius* L.): Effects of Thermal Lipoxygenase Inactivation. *Eur. Food Res. Technol.* **2016**, *242*, 995–1004. [[CrossRef](#)]
63. Boatright, W.L.; Lei, Q. Compounds Contributing to the “Beany” Odor of Aqueous Solutions of Soy Protein Isolates. *J. Food Sci.* **1999**, *64*, 667–670. [[CrossRef](#)]
64. Hongsoongnern, P.; Chambers, E., IV. A Lexicon for Green Odor or Flavor and Characteristics of Chemicals Associated with Green. *J. Sens. Stud.* **2008**, *23*, 205–221. [[CrossRef](#)]
65. Lozano, P.R.; Drake, M.; Benitez, D.; Cadwallader, K.R. Instrumental and Sensory Characterization of Heat-Induced Odorants in Aseptically Packaged Soy Milk. *J. Agric. Food Chem.* **2007**, *55*, 3018–3026. [[CrossRef](#)]
66. Torres-Penaranda, A.V.; Reitmeier, C.A. Sensory Descriptive Analysis of Soymilk. *J. Food Sci.* **2001**, *66*, 352–356. [[CrossRef](#)]
67. Vara-Ubol, S.; Chambers, E.; Chambers, D.H. Sensory Characteristics of Chemical Compounds Potentially Associated with Beany Aroma in Food. *J. Sens. Stud.* **2004**, *19*, 15–26. [[CrossRef](#)]
68. Schindler, S.; Wittig, M.; Zelena, K.; Krings, U.; Bez, J.; Eisner, P.; Berger, R.G. Lactic Fermentation to Improve the Aroma of Protein Extracts of Sweet Lupin (*Lupinus angustifolius*). *Food Chem.* **2011**, *128*, 330–337. [[CrossRef](#)]
69. Schindler, S.; Zelena, K.; Krings, U.; Bez, J.; Eisner, P.; Berger, R. Improvement of the Aroma of Pea (*Pisum sativum*) Protein Extracts by Lactic Acid Fermentation. *Food Biotechnol.* **2012**, *26*, 58–74. [[CrossRef](#)]
70. Blagden, T.; Gilliland, S. Reduction of Levels of Volatile Components Associated with the “Beany” Flavor in Soymilk by *Lactobacilli* and *Streptococci*. *J. Food Sci.* **2006**, *70*, M186–M189. [[CrossRef](#)]
71. Lee, C.; Beuchat, L.R. Changes in Chemical Composition and Sensory Qualities of Peanut Milk Fermented with Lactic Acid Bacteria. *Int. J. Food Microbiol.* **1991**, *13*, 273–283. [[CrossRef](#)]
72. Lorn, D. Screening of Lactic Acid Bacteria for Their Potential Use as Aromatic Starters in Fermented Vegetables. Ph.D. Thesis, University of Burgundy, Dijon, France, 15 December 2020.
73. Axel, C.; Zannini, E.; Arendt, E.K. Mold Spoilage of Bread and Its Biopreservation: A Review of Current Strategies for Bread Shelf Life Extension. *Crit. Rev. Food Sci. Nutr.* **2017**, *57*, 3528–3542. [[CrossRef](#)] [[PubMed](#)]
74. O’Connor, P.M.; Kuniyoshi, T.M.; Oliveira, R.P.; Hill, C.; Ross, R.P.; Cotter, P.D. Antimicrobials for Food and Feed; a Bacteriocin Perspective. *Curr. Opin. Biotechnol.* **2020**, *61*, 160–167. [[CrossRef](#)] [[PubMed](#)]
75. Bansal-Mutalik, R.; Nikaido, H. Quantitative Lipid Composition of Cell Envelopes of *Corynebacterium glutamicum* Elucidated through Reverse Micelle Extraction. *Proc. Natl. Acad. Sci. USA* **2011**, *108*, 15360–15365. [[CrossRef](#)]
76. Shekhar, S.; Sundaramanickam, A.; Thangavel, B. Biosurfactant Producing Microbes and Their Potential Applications: A Review. *Crit. Rev. Environ. Sci. Technol.* **2015**, *45*, 1522–1554. [[CrossRef](#)]
77. Rahman, P.; Edward, G. Production, Characterisation and Applications of Biosurfactants-Review. *Biotechnology* **2008**, *7*, 360–370. [[CrossRef](#)]
78. Amani, H.; Kariminezhad, H. Study on Emulsification of Crude Oil in Water Using Emulsan Biosurfactant for Pipeline Transportation. *Pet. Sci. Technol.* **2016**, *34*, 216–222. [[CrossRef](#)]
79. Chaprão, M.J.; Ferreira, I.N.S.; Correa, P.F.; Rufino, R.D.; Luna, J.M.; Silva, E.J.; Sarubbo, L.A. Application of Bacterial and Yeast Biosurfactants for Enhanced Removal and Biodegradation of Motor Oil from Contaminated Sand. *Electron. J. Biotechnol.* **2015**, *18*, 471–479. [[CrossRef](#)]
80. Sajna, K.; Höfer, R.; Sukumaran, R.; Gottumukkala, L.; Pandey, A. White Biotechnology in Biosurfactants. In *Industrial Biorefineries & White Biotechnology*; Elsevier: Amsterdam, The Netherlands, 2015; pp. 499–521.
81. Joshi, S.; Bharucha, C.; Desai, A.J. Production of Biosurfactant and Antifungal Compound by Fermented Food Isolate *Bacillus subtilis* 20B. *Bioresour. Technol.* **2008**, *99*, 4603–4608. [[CrossRef](#)]
82. Magalhães, L.; Nitschke, M. Antimicrobial Activity of Rhamnolipids against *Listeria monocytogenes* and Their Synergistic Interaction with Nisin. *Food Control* **2013**, *29*, 138–142. [[CrossRef](#)]

83. Mnif, I.; Ghribi, D. Glycolipid Biosurfactants: Main Properties and Potential Applications in Agriculture and Food Industry. *J. Sci. Food Agric.* **2016**, *96*, 4310–4320. [[CrossRef](#)]
84. Zhao, Z.; Wang, Q.; Wang, K.; Brian, K.; Liu, C.; Gu, Y. Study of the Antifungal Activity of *Bacillus vallismortis* ZZ185 in vitro and Identification of Its Antifungal Components. *Bioresour. Technol.* **2010**, *101*, 292–297. [[CrossRef](#)]
85. Otzen, D.E. Biosurfactants and Surfactants Interacting with Membranes and Proteins: Same but Different? *Biochim. Biophys. Acta Biomembr.* **2017**, *1859*, 639–649. [[CrossRef](#)]
86. Rodrigues, L.; van der Mei, H.C.; Teixeira, J.; Oliveira, R. Influence of Biosurfactants from Probiotic Bacteria on Formation of Biofilms on Voice Prostheses. *Appl. Environ. Microbiol.* **2004**, *70*, 4408–4410. [[CrossRef](#)] [[PubMed](#)]
87. Gudiña, E.J.; Rocha, V.; Teixeira, J.A.; Rodrigues, L.R. Antimicrobial and Antiadhesive Properties of a Biosurfactant Isolated from *Lactobacillus paracasei* ssp. *paracasei* A20. *Lett. Appl. Microbiol.* **2010**, *50*, 419–424. [[CrossRef](#)]
88. Rufino, R.D.; Luna, J.M.; Sarubbo, L.A.; Rodrigues, L.R.M.; Teixeira, J.A.C.; Campos-Takaki, G.M. Antimicrobial and Anti-Adhesive Potential of a Biosurfactant Rufisan Produced by *Candida lipolytica* UCP 0988. *Colloids Surf. B Biointerfaces* **2011**, *84*, 1–5. [[CrossRef](#)]
89. Abalos, A.; Pinazo, A.; Infante, M.R.; Casals, M.; García, F.; Manresa, A. Physicochemical and Antimicrobial Properties of New Rhamnolipids Produced by *Pseudomonas aeruginosa* AT10 from Soybean Oil Refinery Wastes. *Langmuir* **2001**, *17*, 1367–1371. [[CrossRef](#)]
90. Padmapriya, B.; Suganthi, S.; Anishya, R.S. Screening, Optimization and Production of Biosurfactants by *Candida* Species Isolated from Oil Polluted Soils. *Am. Eurasian J. Agric. Environ. Sci.* **2013**, *13*, 227–233.
91. Kiran, G.S.; Sabarathnam, B.; Selvin, J. Biofilm Disruption Potential of a Glycolipid Biosurfactant from Marine *Brevibacterium casei*. *FEMS Immunol. Med. Microbiol.* **2010**, *59*, 432–438. [[CrossRef](#)] [[PubMed](#)]
92. Goma, E. Antimicrobial Activity of a Biosurfactant Produced by *Bacillus licheniformis* Strain M104 Grown on Whey. *Braz. Arch. Biol. Technol.* **2013**, *56*, 259–268. [[CrossRef](#)]
93. Falardeau, J.; Wise, C.; Novitsky, L.; Avis, T.J. Ecological and Mechanistic Insights into the Direct and Indirect Antimicrobial Properties of *Bacillus subtilis* Lipopeptides on Plant Pathogens. *J. Chem. Ecol.* **2013**, *39*, 869–878. [[CrossRef](#)] [[PubMed](#)]
94. Sharma, D.; Ansari, M.J.; Gupta, S.; Al Ghamdi, A.; Pruthi, P.; Pruthi, V. Structural Characterization and Antimicrobial Activity of a Biosurfactant Obtained from *Bacillus pumilus* DSVP18 Grown on Potato Peels. *Jundishapur J. Microbiol.* **2015**, *8*, e21257. [[CrossRef](#)]
95. Sriram, M.I.; Kalishwaralal, K.; Deepak, V.; Gracerosept, R.; Srisakthi, K.; Gurunathan, S. Biofilm Inhibition and Antimicrobial Action of Lipopeptide Biosurfactant Produced by Heavy Metal Tolerant Strain *Bacillus cereus* NK1. *Colloids Surf. B Biointerfaces* **2011**, *85*, 174–181. [[CrossRef](#)]
96. Das, P.; Mukherjee, S.; Sen, R. Antimicrobial Potential of a Lipopeptide Biosurfactant Derived from a Marine *Bacillus circulans*. *J. Appl. Microbiol.* **2008**, *104*, 1675–1684. [[CrossRef](#)] [[PubMed](#)]
97. Ruangprachaya, F.; Chuenchomrat, P. Isolation and Characterization of Biosurfactant Produced by Lactic Acid Bacteria from Indigenous Thai Fermented Foods. *Int. J. Food Eng.* **2018**, *4*. [[CrossRef](#)]
98. Saravanakumari, P.; Mani, K. Structural Characterization of a Novel Xylolipid Biosurfactant from *Lactococcus lactis* and Analysis of Antibacterial Activity against Multi-Drug Resistant Pathogens. *Bioresour. Technol.* **2010**, *101*, 8851–8854. [[CrossRef](#)] [[PubMed](#)]
99. Sharma, D.; Singh Saharan, B. Simultaneous Production of Biosurfactants and Bacteriocins by Probiotic *Lactobacillus casei* MRTL3. *Int. J. Microbiol.* **2014**, *2014*, 698713. [[CrossRef](#)]
100. Madhu, A.N.; Prapulla, S.G. Evaluation and Functional Characterization of a Biosurfactant Produced by *Lactobacillus plantarum* CFR 2194. *Appl. Biochem. Biotechnol.* **2014**, *172*, 1777–1789. [[CrossRef](#)] [[PubMed](#)]
101. Gudiña, E.J.; Fernandes, E.C.; Teixeira, J.A.; Rodrigues, L.R. Antimicrobial and Anti-Adhesive Activities of Cell-Bound Biosurfactant from *Lactobacillus agilis* CCUG31450. *RSC Adv.* **2015**, *5*, 90960–90968. [[CrossRef](#)]
102. Mnif, I.; Hammami, I.; Triki, M.A.; Azabou, M.C.; Ellouze-Chaabouni, S.; Ghribi, D. Antifungal Efficiency of a Lipopeptide Biosurfactant Derived from *Bacillus subtilis* SPB1 versus the Phytopathogenic Fungus, *Fusarium solani*. *Environ. Sci. Pollut. Res. Int.* **2015**, *22*, 18137–18147. [[CrossRef](#)]
103. Pretorius, D.; van Rooyen, J.; Clarke, K.G. Enhanced Production of Antifungal Lipopeptides by *Bacillus amyloliquefaciens* for Biocontrol of Postharvest Disease. *N. Biotechnol.* **2015**, *32*, 243–252. [[CrossRef](#)]
104. Kilani-Feki, O.; Ben Khedher, S.; Dammak, M.; Kamoun, A.; Jabnoun-Khiareddine, H.; Daami-Remadi, M.; Tounsi, S. Improvement of Antifungal Metabolites Production by *Bacillus subtilis* V26 for Biocontrol of Tomato Postharvest Disease. *Biol. Control* **2016**, *95*, 73–82. [[CrossRef](#)]
105. Gu, K.-B.; Zhang, D.-J.; Guan, C.; Xu, J.-H.; Li, S.-L.; Shen, G.-M.; Luo, Y.-C.; Li, Y.-G. Safe Antifungal Lipopeptides Derived from *Bacillus marinus* B-9987 Against Grey Mold Caused by *Botrytis cinerea*. *J. Integr. Agric.* **2017**, *16*, 1999–2008. [[CrossRef](#)]
106. Torres, M.J.; Brandan, C.P.; Petroselli, G.; Erra-Balsells, R.; Audisio, M.C. Antagonistic Effects of *Bacillus subtilis* subsp. *subtilis* and *B. amyloliquefaciens* against *Macrophomina phaseolina*: SEM Study of Fungal Changes and UV-MALDI-TOF MS Analysis of Their Bioactive Compounds. *Microbiol. Res.* **2016**, *182*, 31–39. [[CrossRef](#)]
107. Sachdev, D.P.; Cameotra, S.S. Biosurfactants in Agriculture. *Appl. Microbiol. Biotechnol.* **2013**, *97*, 1005–1016. [[CrossRef](#)] [[PubMed](#)]
108. González-Jaramillo, L.M.; Aranda, F.J.; Teruel, J.A.; Villegas-Escobar, V.; Ortiz, A. Antimycotic Activity of Fengycin C Biosurfactant and Its Interaction with Phosphatidylcholine Model Membranes. *Colloids Surf. B Biointerfaces* **2017**, *156*, 114–122. [[CrossRef](#)] [[PubMed](#)]

109. Jadhav, M.; Kagalkar, A.; Jadhav, S.; Govindwar, S. Isolation, Characterization, and Antifungal Application of a Biosurfactant Produced by *Enterobacter* sp. MS16. *Eur. J. Lipid Sci. Technol.* **2011**, *113*, 1347–1356. [[CrossRef](#)]
110. Yoo, D.S.; Lee, B.S.; Kim, E.K. Characteristics of Microbial Biosurfactant as an Antifungal Agent Against Plant Pathogenic Fungus. *J. Microbiol. Biotechnol.* **2005**, *15*, 1164–1169.
111. Gond, S.K.; Bergen, M.S.; Torres, M.S.; White, J.F., Jr. Endophytic *Bacillus* spp. Produce Antifungal Lipopeptides and Induce Host Defence Gene Expression in Maize. *Microbiol. Res.* **2015**, *172*, 79–87. [[CrossRef](#)] [[PubMed](#)]
112. Eskola, M.; Kos, G.; Elliott, C.T.; Hajšlová, J.; Mayar, S.; Krska, R. Worldwide Contamination of Food-Crops with Mycotoxins: Validity of the Widely Cited ‘FAO Estimate’ of 25. *Crit. Rev. Food Sci. Nutr.* **2020**, *60*, 2773–2789. [[CrossRef](#)] [[PubMed](#)]
113. Muhialdin, B.J.; Saari, N.; Meor Hussin, A.S. Review on the Biological Detoxification of Mycotoxins Using Lactic Acid Bacteria to Enhance the Sustainability of Foods Supply. *Molecules* **2020**, *25*, 2655. [[CrossRef](#)]
114. Watanakij, N.; Visessanguan, W.; Petchkongkaew, A. Aflatoxin B₁-Degrading Activity from *Bacillus subtilis* BCC 42005 Isolated from Fermented Cereal Products. *Food Addit. Contam. Part. A Chem. Anal. Control. Expo. Risk Assess.* **2020**, *37*, 1579–1589. [[CrossRef](#)]
115. Fazeli, M.R.; Hajimohammadali, M.; Moshkani, A.; Samadi, N.; Jamalifar, H.; Khoshayand, M.R.; Vaghari, E.; Pouragahi, S. Aflatoxin B₁ Binding Capacity of Autochthonous Strains of Lactic Acid Bacteria. *J. Food Prot.* **2009**, *72*, 189–192. [[CrossRef](#)] [[PubMed](#)]
116. Hernandez-Mendoza, A.; Garcia, H.S.; Steele, J.L. Screening of *Lactobacillus casei* Strains for Their Ability to Bind Aflatoxin B₁. *Food Chem. Toxicol.* **2009**, *47*, 1064–1068. [[CrossRef](#)] [[PubMed](#)]
117. El-Nezami, H.; Kankaanpaa, P.; Salminen, S.; Ahokas, J. Ability of Dairy Strains of Lactic Acid Bacteria to Bind a Common Food Carcinogen, Aflatoxin B₁. *Food Chem. Toxicol.* **1998**, *36*, 321–326. [[CrossRef](#)]
118. Peltonen, K.; El-Nezami, H.; Haskard, C.; Ahokas, J.; Salminen, S. Aflatoxin B₁ Binding by Dairy Strains of Lactic Acid Bacteria and Bifidobacteria. *J. Dairy Sci.* **2001**, *84*, 2152–2156. [[CrossRef](#)]
119. Topcu, A.; Bulat, T.; Wishah, R.; Boyacı, I.H. Detoxification of Aflatoxin B₁ and Patulin by *Enterococcus faecium* Strains. *Int. J. Food Microbiol.* **2010**, *139*, 202–205. [[CrossRef](#)]
120. Khanafari, A.; Soudi, H.; Miraboufathi, M.; Osboo, R.K. An in vitro Investigation of Aflatoxin B₁ Biological Control by *Lactobacillus plantarum*. *Pak. J. Biol. Sci.* **2007**, *10*, 2553–2556. [[CrossRef](#)]
121. Shah, N.; Wu, S. Aflatoxin B₁ Binding Abilities of Probiotic Bacteria. *Biosci. Microflora* **1999**, *18*, 43–48. [[CrossRef](#)]
122. Sarimehmetoğlu, B.; Küplülü, Ö. Binding Ability of Aflatoxin M₁ to Yoghurt Bacteria. *Ankara Üniversitesi Veteriner Fakültesi Dergisi* **2004**, *51*, 195–198.
123. Sezer, Ç.; Güven, A.; Oral, N.B.; Vatansver, L. Detoxification of Aflatoxin B₁ by Bacteriocins and Bacteriocinogenic Lactic Acid Bacteria. *Turk. J. Vet. Anim. Sci.* **2013**, *37*, 594–601. [[CrossRef](#)]
124. Elsanhoty, R.M.; Salam, S.A.; Ramadan, M.F.; Badr, F.H. Detoxification of Aflatoxin M₁ in Yoghurt Using Probiotics and Lactic Acid Bacteria. *Food Control* **2014**, *43*, 129–134. [[CrossRef](#)]
125. Ślizewska, K.; Smulikowska, S. Detoxification of Aflatoxin B₁ and Change in Microflora Pattern by Probiotic in vitro Fermentation of Broiler Feed. *J. Anim. Feed Sci.* **2011**, *20*, 300–309. [[CrossRef](#)]
126. Huang, L.; Duan, C.; Zhao, Y.; Gao, L.; Niu, C.; Xu, J.; Li, S. Reduction of Aflatoxin B₁ Toxicity by *Lactobacillus plantarum* C88: A Potential Probiotic Strain Isolated from Chinese Traditional Fermented Food “Tofu”. *PLoS ONE* **2017**, *12*, e0170109. [[CrossRef](#)]
127. Niderkorn, V.; Morgavi, D.P.; Aboab, B.; Lemaire, M.; Boudra, H. Cell Wall Component and Mycotoxin Moieties Involved in the Binding of Fumonisin B₁ and B₂ by Lactic Acid Bacteria. *J. Appl. Microbiol.* **2009**, *106*, 977–985. [[CrossRef](#)]
128. Niderkorn, V.; Boudra, H.; Morgavi, D.P. Binding of Fusarium Mycotoxins by Fermentative Bacteria in vitro. *J. Appl. Microbiol.* **2006**, *101*, 849–856. [[CrossRef](#)]
129. El-Nezami, H.; Polychronaki, N.; Salminen, S.; Mykkänen, H. Binding Rather than Metabolism May Explain the Interaction of Two Food-Grade *Lactobacillus* strains with Zearalenone and Its Derivative Alpha-Zearalenol. *Appl. Environ. Microbiol.* **2002**, *68*, 3545–3549. [[CrossRef](#)]
130. Long, M.; Li, P.; Zhang, W.; Li, X.B.; Zhang, Y.; Wang, Z.; Liu, G. Removal of Zearalenone by Strains of *Lactobacillus* sp. Isolated from Rumen in vitro. *J. Anim. Vet. Adv.* **2012**, *11*, 2417–2422.
131. Rogowska, A.; Pomastowski, P.; Walczak, J.; Railean-Plugaru, V.; Rudnicka, J.; Buszewski, B. Investigation of Zearalenone Adsorption and Biotransformation by Microorganisms Cultured under Cellular Stress Conditions. *Toxins* **2019**, *11*, 463. [[CrossRef](#)]
132. Franco, T.S.; Garcia, S.; Hirooka, E.Y.; Ono, Y.S.; dos Santos, J.S. Lactic acid Bacteria in the Inhibition of *Fusarium graminearum* and Deoxynivalenol Detoxification. *J. Appl. Microbiol.* **2011**, *111*, 739–748. [[CrossRef](#)]
133. Fuchs, S.; Sontag, G.; Stidl, R.; Ehrlich, V.; Kundi, M.; Knasmüller, S. Detoxification of Patulin and Ochratoxin A, Two Abundant Mycotoxins, by Lactic Acid Bacteria. *Food Chem. Toxicol.* **2008**, *46*, 1398–1407. [[CrossRef](#)] [[PubMed](#)]
134. Wang, L.; Yue, T.; Yuan, Y.; Wang, Z.; Ye, M.; Cai, R. A New Insight into the Adsorption Mechanism of Patulin by the Heat-Inactive Lactic Acid Bacteria Cells. *Food Control* **2015**, *50*, 104–110. [[CrossRef](#)]
135. Del Prete, V.; Rodriguez, H.; Carrascosa, A.V.; de las Rivas, B.; Garcia-Moruno, E.; Muñoz, R. In vitro Removal of Ochratoxin A by Wine Lactic Acid Bacteria. *J. Food Prot.* **2007**, *70*, 2155–2160. [[CrossRef](#)]
136. Mateo, E.M.; Medina, Á.; Mateo, F.; Valle-Algarra, F.M.; Pardo, I.; Jiménez, M. Ochratoxin A Removal in Synthetic Media by Living and Heat-Inactivated Cells of *Oenococcus oeni* Isolated from Wines. *Food Control* **2010**, *21*, 23–28. [[CrossRef](#)]
137. Abrunhosa, L.; Inês, A.; Rodrigues, A.I.; Guimarães, A.; Pereira, V.L.; Parpot, P.; Mendes-Faia, A.; Venâncio, A. Biodegradation of Ochratoxin A by *Pediococcus parvulus* Isolated from Douro Wines. *Int. J. Food Microbiol.* **2014**, *188*, 45–52. [[CrossRef](#)] [[PubMed](#)]

138. Luz, C.; Ferrer, J.; Mañes, J.; Meca, G. Toxicity Reduction of Ochratoxin A by Lactic Acid Bacteria. *Food Chem. Toxicol.* **2018**, *112*, 60–66. [[CrossRef](#)] [[PubMed](#)]
139. Petchkongkaew, A.; Taillandier, P.; Gasaluck, P.; Lebrihi, A. Isolation of *Bacillus* spp. from Thai Fermented Soybean (Thua-Nao): Screening for Aflatoxin B₁ and Ochratoxin A Detoxification. *J. Appl. Microbiol.* **2008**, *104*, 1495–1502. [[CrossRef](#)]
140. Liu, J.R.; Yi, P.J. *Bacillus licheniformis* and Method for Detoxification of Zearalenone. U.S. Patent 8,404,477, 26 March 2013.
141. Cho, K.J.; Kang, J.S.; Cho, W.T.; Lee, C.H.; Ha, J.K.; Song, K.B. In Vitro Degradation of Zearalenone by *Bacillus subtilis*. *Biotechnol. Lett.* **2010**, *32*, 1921–1924. [[CrossRef](#)] [[PubMed](#)]
142. Robert, R. When Multinationals Pivot: Three Strategies from the Food Industry. *Paris Innovation Review*, 2016.
143. Makkar, H.P.S. Antinutritional Factors in Foods for Livestock. *BSAP Occas. Publ.* **1993**, *16*, 69–85. [[CrossRef](#)]
144. Gilani, G.S.; Cockell, K.A.; Sepehr, E. Effects of Antinutritional Factors on Protein Digestibility and Amino Acid Availability in Foods. *J. AOAC Int.* **2005**, *88*, 967–987. [[CrossRef](#)] [[PubMed](#)]
145. Fekadu Gemedo, H. Antinutritional Factors in Plant Foods: Potential Health Benefits and Adverse Effects. *Int. J. Nutr. Food Sci.* **2014**, *3*, 284. [[CrossRef](#)]
146. El-Adawy, T.A. Nutritional Composition and Antinutritional Factors of Chickpeas (*Cicer arietinum* L.) Undergoing Different Cooking Methods and Germination. *Plant. Foods Hum. Nutr.* **2002**, *57*, 83–97. [[CrossRef](#)]
147. Ibrahim, S.; Habiba, R.; Shatta, A.; Embaby, H. Effect of Soaking, Germination, Cooking and Fermentation on Antinutritional Factors in Cowpeas. *Nahrung* **2002**, *46*, 92–95. [[CrossRef](#)]
148. Ghavidel, R.A.; Prakash, J. The Impact of Germination and Dehulling on Nutrients, Antinutrients, In Vitro Iron and Calcium Bioavailability and In Vitro Starch and Protein Digestibility of Some Legume Seeds. *LWT Food Sci. Technol.* **2007**, *40*, 1292–1299. [[CrossRef](#)]
149. Martín-Cabrejas, M.A.; Sanfiz, B.; Vidal, A.; Mollá, E.; Esteban, R.; López-Andréu, F.J. Effect of Fermentation and Autoclaving on Dietary Fiber Fractions and Antinutritional Factors of Beans (*Phaseolus vulgaris* L.). *J. Agric. Food Chem.* **2004**, *52*, 261–266. [[CrossRef](#)] [[PubMed](#)]
150. Refstie, S.; Sahlström, S.; Bråthen, E.; Baeverfjord, G.; Krogedal, P. Lactic Acid Fermentation Eliminates Indigestible Carbohydrates and Antinutritional Factors in Soybean Meal for Atlantic Salmon (*Salmo salar*). *Aquaculture* **2005**, *246*, 331–345. [[CrossRef](#)]
151. Granito, M.; Álvarez, G. Lactic Acid Fermentation of Black Beans (*Phaseolus vulgaris*): Microbiological and Chemical Characterization. *J. Sci. Food Agric.* **2006**, *86*, 1164–1171. [[CrossRef](#)]
152. El Hag, M.E.; El Tinay, A.H.; Yousif, N.E. Effect of Fermentation and Dehulling on Starch, Total Polyphenols, Phytic Acid Content and In Vitro Protein Digestibility of Pearl Millet. *Food Chem.* **2002**, *77*, 193–196. [[CrossRef](#)]
153. Bartkiene, E.; Krungleviciute, V.; Juodeikiene, G.; Vidmantiene, D.; Maknickiene, Z. Solid State Fermentation with Lactic Acid Bacteria to Improve the Nutritional Quality of Lupin and Soya Bean. *J. Sci. Food Agric.* **2015**, *95*, 1336–1342. [[CrossRef](#)]
154. Wu, W.; Zhang, B. Lactic Acid Bacteria and B Vitamins. In *Lactic Acid Bacteria*; Chen, W., Ed.; Springer: Singapore, 2019.
155. Ogbonna, J.C. Microbiological Production of Tocopherols: Current State and Prospects. *Appl. Microbiol. Biotechnol.* **2009**, *84*, 217–225. [[CrossRef](#)]
156. Verwaal, R.; Wang, J.; Meijnen, J.P.; Visser, H.; Sandmann, G.; Van Den Berg, J.A.; Van Ooyen, A.J. High-Level Production of Beta-Carotene in *Saccharomyces cerevisiae* by Successive Transformation with Carotenogenic Genes from *Xanthophyllomyces dendrorhous*. *Appl. Environ. Microbiol.* **2007**, *73*, 4342–4350. [[CrossRef](#)]
157. Yan, G.L.; Wen, K.R.; Duan, C.Q. Enhancement of Beta-Carotene Production by Over-Expression of HMG-CoA Reductase Coupled with Addition of Ergosterol Biosynthesis Inhibitors in Recombinant *Saccharomyces cerevisiae*. *Curr. Microbiol.* **2012**, *64*, 159–163. [[CrossRef](#)]
158. Araya-Garay, J.M.; Feijoo-Siota, L.; Rosa-Dos-Santos, F.; Veiga-Crespo, P.; Villa, T.G. Construction of New *Pichia pastoris* X-33 Strains for Production of Lycopene and Beta-Carotene. *Appl. Microbiol. Biotechnol.* **2012**, *93*, 2483–2492. [[CrossRef](#)] [[PubMed](#)]
159. Bhatayaa, A.; Schmidt-Dannerta, C.; Leeb, P. Metabolic Engineering of *Pichia pastoris* X-33 for Lycopene Production. *Process Biochem.* **2009**, *44*, 1095–1102. [[CrossRef](#)]
160. Sabirova, J.S.; Haddouche, R.; Van Bogaert, I.N.; Mulaa, F.; Verstraete, W.; Timmis, K.N.; Schmidt-Dannert, C.; Nicaud, J.M.; Soetaert, W. The “LipoYeasts” Project: Using the Oleaginous Yeast *Yarrowia lipolytica* in Combination with Specific Bacterial Genes for the Bioconversion of Lipids, Fats and Oils into Highvalue Products. *Microb. Biotechnol.* **2011**, *4*, 47–54. [[CrossRef](#)] [[PubMed](#)]
161. Souza, C.M.; Schwabe, T.M.; Pichler, H.; Ploier, B.; Leitner, E.; Guan, X.L.; Wenk, M.R.; Riezman, I.; Riezman, H. A Stable Yeast Strain Efficiently Producing Cholesterol Instead of Ergosterol is Functional for Tryptophan Uptake, but not Weak Organic Acid Resistance. *Metab. Eng.* **2011**, *13*, 555–569. [[CrossRef](#)]
162. Taghuchi, H.; Shibata, T.; Duangmanee, C.; Tani, Y. Menaquinone-4 Production by a Sulfonamide-Resistant Mutant of *Favobacterium* sp. 238-7. *Agric. Biol. Chem.* **1989**, *53*, 3017–3023.
163. Furuichi, K.; Hojo, K.; Katakura, Y.; Ninomiya, K.; Shioya, S. Aerobic Culture of *Propionibacterium freudenreichii* ET-3 Can Increase Production Ratio of 1,4-dihydroxy-2-Naphthoic Acid to Menaquinone. *J. Biosci. Bioeng.* **2006**, *101*, 464–470. [[CrossRef](#)] [[PubMed](#)]
164. Sato, T.; Yamada, Y.; Ohtani, Y.; Mitsui, N.; Murasawa, H.; Araki, S. Efficient Production of Menaquinone (Vitamin K2) by a Menadione-resistant Mutant of *Bacillus subtilis*. *J. Ind. Microbiol. Biotechnol.* **2001**, *26*, 115–120. [[CrossRef](#)]
165. Ledesma-Amaro, R.; Jimenez, A.; Santos, M.A.; Revuelta, J.L. Microbial Production of Vitamins. In *Microbial Production of Food Ingredients, Enzymes and Nutraceuticals*; Series in Food Science; Woodhead Publishing: Cambridge, UK, 2013.

166. Urbance, J.W.; Bratina, B.J.; Stoddard, S.F.; Schmidt, T.M. Taxonomic Characterization of *Ketogulonigenium vulgare* gen. nov. sp. nov. and *Ketogulonigenium robustum* sp. nov. which Oxidize l-Sorbose to 2-Keto-l-Gulonic Acid. *Int. J. Syst. Evol. Microbiol.* **2001**, *51*, 1059–1107. [[CrossRef](#)]
167. Sugisawa, T.; Hoshino, T.; Masuda, S.; Nomura, S.; Setoguchi, Y.; Tazoe, M.; Shinjoh, M.; Someha, S.; Fujiwara, A. Microbial Production of 2-Keto-l-Gulonic Acid from l-Sorbose and d-Sorbitol by *Gluconobacter melanogenus*. *Agric. Biol. Chem.* **1990**, *54*, 1201–1209. [[CrossRef](#)]
168. World Health Organization. *Guideline: Daily Iron and Folic Acid Supplementation in Pregnant Women*; World Health Organization: Geneva, Switzerland, 2012.
169. Saubade, F.; Hemery, Y.M.; Guyot, J.P.; Humblot, C. Lactic Acid Fermentation as a Tool for Increasing the Folate Content of Foods. *Crit. Rev. Food Sci. Nutr.* **2017**, *57*, 3894–3910. [[CrossRef](#)]
170. Bhutta, Z.A.; Salam, R.A.; Das, J.K. Meeting the Challenges of Micronutrient Malnutrition in the Developing World. *Br. Med. Bull.* **2013**, *106*, 7–17. [[CrossRef](#)]
171. Padalino, M.; Perez-Conesa, D.; Lopez-Nicol, R.; Frontela-Saseta, C.; Ros-Berruezo, G. Effect of Fructooligosaccharides and Galactooligosaccharides on the Folate Production of Some Folate-Producing Bacteria in Media Cultures or Milk. *Int. Dairy J.* **2012**, *27*, 27–33. [[CrossRef](#)]
172. Pompei, A.; Cordisco, L.; Amaretti, A.; Zanoni, S.; Matteuzzi, D.; Rossi, M. Folate Production by *Bifidobacteria* as a Potential Probiotic Property. *Appl. Env. Microbiol.* **2007**, *73*, 179–185. [[CrossRef](#)] [[PubMed](#)]
173. Lin, M.Y.; Young, C.M. Folate Levels in Cultures of Lactic Acid Bacteria. *Int. Dairy J.* **2000**, *10*, 409–413. [[CrossRef](#)]
174. Laino, J.E.; Juarez del Valle, M.; Savoy de Giori, G.; LeBlanc, J.G.J. Applicability of a *Lactobacillus amylovorus* Strain as Co-Culture for Natural Folate Bio-Enrichment of Fermented Milk. *Int. J. Food Microbiol.* **2014**, *191*, 10–16. [[CrossRef](#)]
175. Hugenschmidt, S.; Schwenninger, S.M.; Gnehm, N.; Lacroix, C. Screening of a Natural Biodiversity of Lactic and Propionic Acid Bacteria for Folate and Vitamin B12 Production in Supplemented Whey Permeate. *Int. Dairy J.* **2010**, *20*, 852–857. [[CrossRef](#)]
176. Masuda, M.; Ide, M.; Utsumi, H.; Niuro, T.; Shimamura, Y.; Murata, M. Production Potency of Folate, Vitamin B-12, and Thiamine by Lactic Acid Bacteria Isolated from Japanese Pickles. *Biosci. Biotechnol. Biochem.* **2012**, *76*, 2061–2067. [[CrossRef](#)]
177. Cardenas, N.; Laino, J.E.; Delgado, S.; Jimenez, E.; Del Valle, M.J.; De Giori, G.S.; Sesma, F.; Mayo, B.; Fernandez, L.; LeBlanc, J.G.; et al. Relationships Between the Genome and Some Phenotypical Properties of *Lactobacillus fermentum* CECT 5716, a Probiotic Strain Isolated from Human Milk. *Appl. Microbiol. Biotechnol.* **2015**, *99*, 4343–4353. [[CrossRef](#)]
178. Nor, N.M.; Mohamad, R.; Foo, H.L.; Rahim, R.A. Improvement of Folate Biosynthesis by Lactic Acid Bacteria Using Response Surface Methodology. *Food Technol. Biotechnol.* **2010**, *48*, 243–250.
179. Sybesma, W.; Starrenburg, M.; Tijsseling, L.; Hoefnagel, M.H.N.; Hugenholtz, J. Effects of Cultivation Conditions on Folate Production by Lactic Acid Bacteria. *Appl. Environ. Microbiol.* **2003**, *69*, 4542–4548. [[CrossRef](#)] [[PubMed](#)]
180. Laino, J.E.; Guy LeBlanc, J.; Savoy de Giori, G. Production of Natural Foliates by Lactic Acid Bacteria Starter Cultures Isolated from Artisanal Argentinean Yogurts. *Can. J. Microbiol.* **2012**, *58*, 581–588. [[CrossRef](#)]
181. Guyton, A.C. *Textbook of Medical Physiology*, 8th ed.; W.B. Saunders: Philadelphia, PA, USA, 1991.
182. McLaughlin, S.; Margolskee, R.F. The Sense of Taste. *Am. Sci.* **1994**, *82*, 538–545.
183. Yamaguchi, S. Basic Properties of Umami and Effects on Humans. *Physiol. Behav.* **1991**, *49*, 833–841. [[CrossRef](#)]
184. Yamaguchi, S. Fundamental Properties of Umami on Human Taste Sensation. In *Umami: A Basic Taste*; Kawamura, Y., Kare, M.R., Eds.; Marcel Dekker: New York, NY, USA, 1987; pp. 41–73.
185. Giacometti, T. Free and Bound Glutamate in Natural Products. In *Glutamic Acid: Advances in Biochemistry and Physiology*; Filer, L.J., Garattini, S., Kare, M.R., Reynolds, A.W., Wurtman, R.J., Eds.; Raven Press: New York, NY, USA, 1979; pp. 25–34.
186. Yamaguchi, S.; Ninomiya, K. Umami and Food Palatability. *J. Nutr.* **2000**, *130* (Suppl. S4), 921s–926s. [[CrossRef](#)] [[PubMed](#)]
187. Ngoc, N.T.T.; Thuan, D.V.; Thanh, D.V.; Quan, L.-H. Optimization for Batch Proteolytic Hydrolysis of Spent Brewer's Yeast by Using Proteases. *J. Sci. Technol.* **2016**, *54*, 181–188.
188. Ngoc, N.T.T.; Quan, L.-H.; Thuan, D.V.; Thanh, D.V. Influences of Technological Hydrolysis Condition on Nucleic Acid Content of Spent Brewer's Yeast Hydrolysate. *Vietnam J. Sci. Technol.* **2016**, *55*, 169–177. [[CrossRef](#)]
189. Hung, W.C.; Lee, M.T.; Chung, H.; Sun, Y.T.; Chen, H.; Charron, N.E.; Huang, H.W. Comparative Study of the Condensing Effects of Ergosterol and Cholesterol. *Biophys. J.* **2016**, *110*, 2026–2033. [[CrossRef](#)]
190. Hull, C.M.; Loveridge, E.J.; Donnison, I.S.; Kelly, D.E.; Kelly, S.L. Co-Production of Bioethanol and Probiotic Yeast Biomass from Agricultural Feedstock: Application of the Rural Biorefinery Concept. *AMB Express* **2014**, *4*, 64. [[CrossRef](#)] [[PubMed](#)]
191. Zhang, K.; Tong, M.; Gao, K.; Di, Y.; Wang, P.; Zhang, C.; Wu, X.; Zheng, D. Genomic Reconstruction to Improve Bioethanol and Ergosterol Production of Industrial Yeast *Saccharomyces Cerevisiae*. *J. Ind. Microbiol. Biotechnol.* **2015**, *42*, 207–218. [[CrossRef](#)] [[PubMed](#)]
192. Blaga, A.C.; Ciobanu, C.; Caşcaval, D.; Galaction, A.-I. Enhancement of Ergosterol Production by *Saccharomyces cerevisiae* in Batch and Fed-Batch Fermentation Processes Using n-Dodecane as Oxygen-Vector. *Biochem. Eng. J.* **2018**, *131*, 70–76. [[CrossRef](#)]