

**Table S1: Mean number of *T. gondii* tachyzoites ( $\times 10^4$ ) per well in culture supernatant of Vero cells among all studied groups:**

Tachyzoite count	Group I (n = 3)	Group II (n = 3)	Group III (n = 3)	Group IV (n = 3)	Group V (n = 3)	F <sub>p</sub>
Mean	46.67	33.33	25.17	25.67	16.33	<0.001*
$\pm$ SD.	3.06	1.15	1.26	0.58	1.53	
% reduction		28.6	46.1	45.0	65.0	
$p_1$		<0.001*	<0.001*	<0.001*	<0.001*	
$p_2$			0.001*	0.002*	<0.001*	
Sig. bet. grps.				$p_3 = 0.996, p_4 = 0.001^*, p_5 < 0.001^*$		

SD: Standard deviation, F: F for ANOVA test, Pairwise comparison bet each 2 groups was done using Post Hoc Test (Tukey)

$p$ :  $p$  value for comparing between the different studied groups

$p_1$ ;  $p$  value for comparing between **Group I** and each other group

$p_2$ ;  $p$  value for comparing between **Group II** and each other group

$p_3$ ;  $p$  value for comparing between **Group III** and **Group IV**

$p_4$ ;  $p$  value for comparing between **Group III** and **Group V**

$p_5$ ;  $p$  value for comparing between **Group IV** and **Group V**

\*: Statistically significant at  $p \leq 0.05$

**Group I:** Control

**Group II:** RA–SLNs, **Group III:** TLA

**Group IV:** TLA/ plain SLNs,

**Group V:** TLA/ RA–SLNs,