

Proceedings

Observing Post-Fire Vegetation Regeneration Dynamics Exploiting High-Resolution Sentinel-2 Data[†]

Federico Filipponi^{1,*} and Giacinto Manfron²

¹ Istituto Superiore per la Protezione e la Ricerca Ambientale, 00144 Roma, Italy

² European Commission Joint Research Centre, 21027 Ispra (VA), Italy; giacinto.manfron@ec.europa.eu

* Correspondence: federico.filipponi@isprambiente.it; Tel.: +39-06-5007-2438

† Presented at the 3rd International Electronic Conference on Remote Sensing, 22 May–5 June 2019;

Available online: <https://ecrs-3.sciforum.net/>.

Supplementary Materials




Figure S1. Temporal distribution as number of wildfire events during year 2017.




Figure S2. Distribution of burned area pixels with respect to hazard Italian national landslide hazard PAI classes.




Figure S3. Density scatterplot highlighting average LAI profiles differences (LAI 2016 minus LAI 2018) as function of the slope of the fire-affected area. Samples (pixels) belongs to 2017 fire affected areas.



© 2019 by the authors; licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC-BY) license (<http://creativecommons.org/licenses/by/4.0/>).