

\*language \*

LOGISTIC REGRESSION VARIABLESneutral  
/METHOD=ENTER Spanish  
/PRINT=GOODFIT CI(95)  
/CRITERIA=PIN(0.05) POUT(0.10) ITERATE(20) CUT(0.5).

## Logistic Regression

### Notes

Output Created		28-AUG-2024 17:46:59
Comments		
Input	Data	C: \Users\lapalacios\Document s\Media manuscript\Database.sav
	Active Dataset	DataSet1
	Filter	<none>
	Weight	<none>
	Split File	<none>
	N of Rows in Working Data File	407
Missing Value Handling	Definition of Missing	User-defined missing values are treated as missing
Syntax		LOGISTIC REGRESSION VARIABLES neutral /METHOD=ENTER Spanish /PRINT=GOODFIT CI(95) /CRITERIA=PIN(0.05) POUT(0.10) ITERATE(20) CUT(0.5).
Resources	Processor Time	00:00:00.00
	Elapsed Time	00:00:00.00

### Case Processing Summary

Unweighted Cases <sup>a</sup>		N	Percent
Selected Cases	Included in Analysis	407	100.0
	Missing Cases	0	.0
	Total	407	100.0
Unselected Cases		0	.0
Total		407	100.0

a. If weight is in effect, see classification table for the total number of cases.

### Dependent Variable Encoding

Original Value	Internal Value
not neutral	0
neutral	1

### Block 0: Beginning Block

**Classification Table<sup>a,b</sup>**

Observed		Predicted		Percentage Correct
		not neutral	neutral	
Step 0	neutral	253	0	100.0
	not neutral	154	0	.0
	Overall Percentage			62.2

a. Constant is included in the model.

b. The cut value is .500

### Variables in the Equation

		B	S.E.	Wald	df	Sig.	Exp(B)
Step 0	Constant	-.496	.102	23.593	1	<.001	.609

### Variables not in the Equation

			Score	df	Sig.
Step 0	Variables	Spanish	.019	1	.891
	Overall Statistics		.019	1	.891

### Block 1: Method = Enter

#### Omnibus Tests of Model Coefficients

		Chi-square	df	Sig.
Step 1	Step	.019	1	.890
	Block	.019	1	.890
	Model	.019	1	.890

### Model Summary

Step	-2 Log likelihood	Cox & Snell R Square	Nagelkerke R Square
1	539.878 <sup>a</sup>	.000	.000

a. Estimation terminated at iteration number 3 because parameter estimates changed by less than .001.

### Hosmer and Lemeshow Test

Step	Chi-square	df	Sig.
1	.000	0	.

### Contingency Table for Hosmer and Lemeshow Test

		neutral = not neutral		neutral = neutral		Total
		Observed	Expected	Observed	Expected	
Step 1	1	59	59.000	35	35.000	94
	2	194	194.000	119	119.000	313

### Classification Table<sup>a</sup>

		Predicted		Percentage Correct
		Observed	neutral	
Step 1	neutral	not neutral	neutral	100.0
	not neutral	253	0	.0
	neutral	154	0	.0
Overall Percentage				62.2

a. The cut value is .500

### Variables in the Equation

		B	S.E.	Wald	df	Sig.	Exp(B)	95% C.I. for
								Lower
Step 1 <sup>a</sup>	Spanish	.033	.243	.019	1	.891	1.034	.642
	Constant	-.522	.213	5.990	1	.014	.593	

### Variables in the Equation

		95% C.I. for ..
		Upper
Step 1 <sup>a</sup>	Spanish	1.665
	Constant	

a. Variable(s) entered on step 1: Spanish.

LOGISTIC REGRESSION VARIABLESneutral  
/METHOD=ENTER Portuguese  
/PRINT=GOODFIT CI(95)  
/CRITERIA=PIN(0.05) POUT(0.10) ITERATE(20) CUT(0.5).

## Logistic Regression

### Notes

Output Created		28-AUG-2024 17:46:59
Comments		
Input	Data	C: \Users\apalacios\Documents\Media manuscript\Database.sav
	Active Dataset	DataSet1
	Filter	<none>
	Weight	<none>
	Split File	<none>
	N of Rows in Working Data File	407
Missing Value Handling	Definition of Missing	User-defined missing values are treated as missing
Syntax		LOGISTIC REGRESSION VARIABLES neutral /METHOD=ENTER Portuguese /PRINT=GOODFIT CI(95) /CRITERIA=PIN(0.05) POUT(0.10) ITERATE(20) CUT(0.5).
Resources	Processor Time	00:00:00.03
	Elapsed Time	00:00:00.02

### Case Processing Summary

Unweighted Cases <sup>a</sup>		N	Percent
Selected Cases	Included in Analysis	407	100.0
	Missing Cases	0	.0
	Total	407	100.0
Unselected Cases		0	.0
Total		407	100.0

a. If weight is in effect, see classification table for the total number of cases.

### Dependent Variable Encoding

Original Value	Internal Value
not neutral	0
neutral	1

### Block 0: Beginning Block

**Classification Table<sup>a,b</sup>**

Observed		Predicted		Percentage Correct
		not neutral	neutral	
Step 0	not neutral	253	0	100.0
	neutral	154	0	.0
	Overall Percentage			62.2

a. Constant is included in the model.

b. The cut value is .500

### Variables in the Equation

		B	S.E.	Wald	df	Sig.	Exp(B)
Step 0	Constant	-.496	.102	23.593	1	<.001	.609

### Variables not in the Equation

		Score	df	Sig.
Step 0	Variables Portuguese	.019	1	.891
	Overall Statistics	.019	1	.891

### Block 1: Method = Enter

#### Omnibus Tests of Model Coefficients

		Chi-square	df	Sig.
Step 1	Step	.019	1	.890
	Block	.019	1	.890
	Model	.019	1	.890

### Model Summary

Step	-2 Log likelihood	Cox & Snell R Square	Nagelkerke R Square
1	539.878 <sup>a</sup>	.000	.000

a. Estimation terminated at iteration number 3 because parameter estimates changed by less than .001.

### Hosmer and Lemeshow Test

Step	Chi-square	df	Sig.
1	.000	0	.

### Contingency Table for Hosmer and Lemeshow Test

		neutral = not neutral		neutral = neutral		Total
		Observed	Expected	Observed	Expected	
Step 1	1	59	59.000	35	35.000	94
	2	194	194.000	119	119.000	313

### Classification Table<sup>a</sup>

			Predicted		Percentage Correct
Observed		not neutral	neutral		
Step 1	neutral	not neutral	253	0	100.0
		neutral	154	0	.0
	Overall Percentage				62.2

a. The cut value is .500

### Variables in the Equation

		B	S.E.	Wald	df	Sig.	Exp(B)	95% C.I. for Lower
Step 1 <sup>a</sup>	Portuguese	-.033	.243	.019	1	.891	.967	.601
	Constant	-.489	.116	17.618	1	<.001	.613	

### Variables in the Equation

		95% C.I. for .. Upper
Step 1 <sup>a</sup>	Portuguese	1.557
	Constant	

a. Variable(s) entered on step 1: Portuguese.

LOGISTIC REGRESSION VARIABLESneutral  
/METHOD=ENTER newsmedia  
/PRINT=GOODFIT CI(95)  
/CRITERIA=PIN(0.05) POUT(0.10) ITERATE(20) CUT(0.5).

## Logistic Regression

### Notes

Output Created		28-AUG-2024 17:47:09
Comments		
Input	Data	C: \Users\apalacios\Document s\Media manuscript\Database.sav
	Active Dataset	DataSet1
	Filter	<none>
	Weight	<none>
	Split File	<none>
	N of Rows in Working Data File	407
Missing Value Handling	Definition of Missing	User-defined missing values are treated as missing
Syntax		LOGISTIC REGRESSION VARIABLES neutral /METHOD=ENTER newsmedia /PRINT=GOODFIT CI(95) /CRITERIA=PIN(0.05) POUT(0.10) ITERATE(20) CUT(0.5).
Resources	Processor Time	00:00:00.03
	Elapsed Time	00:00:00.02

### Case Processing Summary

Unweighted Cases <sup>a</sup>		N	Percent
Selected Cases	Included in Analysis	402	98.8
	Missing Cases	5	1.2
	Total	407	100.0
Unselected Cases		0	.0
Total		407	100.0

a. If weight is in effect, see classification table for the total number of cases.

**Dependent Variable  
Encoding**

Original Value	Internal Value
not neutral	0
neutral	1

**Block 0: Beginning Block**

**Classification Table<sup>a,b</sup>**

		Predicted		Percentage Correct
		not neutral	neutral	
Step 0	neutral	250	0	100.0
	neutral	152	0	.0
	Overall Percentage			62.2

a. Constant is included in the model.

b. The cut value is .500

**Variables in the Equation**

		B	S.E.	Wald	df	Sig.	Exp(B)
Step 0	Constant	-.498	.103	23.404	1	<.001	.608

**Variables not in the Equation**

			Score	df	Sig.
Step 0	Variables	newsmedia	1.469	1	.225
	Overall Statistics		1.469	1	.225

**Block 1: Method = Enter**

**Omnibus Tests of Model Coefficients**

		Chi-square	df	Sig.
Step 1	Step	1.419	1	.234
	Block	1.419	1	.234
	Model	1.419	1	.234



### Model Summary

Step	-2 Log likelihood	Cox & Snell R Square	Nagelkerke R Square
1	531.739 <sup>a</sup>	.004	.005

a. Estimation terminated at iteration number 3 because parameter estimates changed by less than .001.

### Hosmer and Lemeshow Test

Step	Chi-square	df	Sig.
1	.000	0	.

### Contingency Table for Hosmer and Lemeshow Test

		neutral = not neutral		neutral = neutral		Total
		Observed	Expected	Observed	Expected	
Step 1	1	250	250.000	152	152.000	402

### Classification Table<sup>a</sup>

		Predicted		Percentage Correct
Observed		not neutral	neutral	
Step 1	neutral	244	6	97.6
	not neutral	145	7	4.6
Overall Percentage				62.4

a. The cut value is .500

### Variables in the Equation

		B	S.E.	Wald	df	Sig.	Exp(B)	95% C.I. for
								Lower
Step 1 <sup>a</sup>	newsmedia	-.675	.566	1.420	1	.233	.509	.168
	Constant	.154	.556	.077	1	.782	1.167	

### Variables in the Equation

		95% C.I. for ..
		Upper
Step 1 <sup>a</sup>	newsmedia	1.545
	Constant	

a. Variable(s) entered on step 1: newsmedia.

LOGISTIC REGRESSION VARIABLESneutral  
/METHOD=ENTER Op\_Ed  
/PRINT=GOODFIT CI(95)  
/CRITERIA=PIN(0.05) POUT(0.10) ITERATE(20) CUT(0.5).

## Logistic Regression

### Notes

Output Created		28-AUG-2024 17:47:09
Comments		
Input	Data	C: \Users\apalacios\Documents\Media manuscript\Database.sav
	Active Dataset	DataSet1
	Filter	<none>
	Weight	<none>
	Split File	<none>
	N of Rows in Working Data File	407
Missing Value Handling	Definition of Missing	User-defined missing values are treated as missing
Syntax		LOGISTIC REGRESSION VARIABLES neutral /METHOD=ENTER Op_Ed /PRINT=GOODFIT CI(95) /CRITERIA=PIN(0.05) POUT(0.10) ITERATE(20) CUT(0.5).
Resources	Processor Time	00:00:00.03
	Elapsed Time	00:00:00.03

### Case Processing Summary

Unweighted Cases <sup>a</sup>		N	Percent
Selected Cases	Included in Analysis	402	98.8
	Missing Cases	5	1.2
	Total	407	100.0
Unselected Cases		0	.0
Total		407	100.0

a. If weight is in effect, see classification table for the total number of cases.

### Dependent Variable Encoding

Original Value	Internal Value
not neutral	0
neutral	1

### Block 0: Beginning Block

#### Classification Table<sup>a,b</sup>

		Predicted		Percentage Correct
		not neutral	neutral	
Step 0	neutral	250	0	100.0
	neutral	152	0	.0
	Overall Percentage			62.2

a. Constant is included in the model.

b. The cut value is .500

#### Variables in the Equation

		B	S.E.	Wald	df	Sig.	Exp(B)
Step 0	Constant	-.498	.103	23.404	1	<.001	.608

#### Variables not in the Equation

		Score	df	Sig.
Step 0	Variables	.570	1	.450
	Overall Statistics	.570	1	.450

### Block 1: Method = Enter

#### Omnibus Tests of Model Coefficients

		Chi-square	df	Sig.
Step 1	Step	.604	1	.437
	Block	.604	1	.437
	Model	.604	1	.437

### Model Summary

Step	-2 Log likelihood	Cox & Snell R Square	Nagelkerke R Square
1	532.553 <sup>a</sup>	.002	.002

a. Estimation terminated at iteration number 4 because parameter estimates changed by less than .001.

### Hosmer and Lemeshow Test

Step	Chi-square	df	Sig.
1	.000	0	.

### Contingency Table for Hosmer and Lemeshow Test

		neutral = not neutral		neutral = neutral		Total
		Observed	Expected	Observed	Expected	
Step 1	1	6	6.000	2	2.000	8
	2	244	244.000	150	150.000	394

### Classification Table<sup>a</sup>

		Predicted		Percentage Correct
		Observed	neutral	
Step 1	neutral	not neutral	neutral	100.0
	not neutral	250	0	.0
	Overall Percentage	152	0	62.2

a. The cut value is .500

### Variables in the Equation

		B	S.E.	Wald	df	Sig.	Exp(B)	95% C.I. for
								Lower
Step 1 <sup>a</sup>	Op_Ed	-.612	.823	.553	1	.457	.542	.108
	Constant	-.487	.104	21.989	1	<.001	.615	

### Variables in the Equation

		95% C.I. for ..
		Upper
Step 1 <sup>a</sup>	Op_Ed	2.721
	Constant	

a. Variable(s) entered on step 1: Op\_Ed.

LOGISTIC REGRESSION VARIABLESneutral  
/METHOD=ENTER New\_treatment  
/PRINT=GOODFIT CI(95)  
/CRITERIA=PIN(0.05) POUT(0.10) ITERATE(20) CUT(0.5).

## Logistic Regression

### Notes

Output Created		28-AUG-2024 17:49:38
Comments		
Input	Data	C: \Users\apalacios\Documents\Media manuscript\Database.sav
	Active Dataset	DataSet1
	Filter	<none>
	Weight	<none>
	Split File	<none>
	N of Rows in Working Data File	407
Missing Value Handling	Definition of Missing	User-defined missing values are treated as missing
Syntax		LOGISTIC REGRESSION VARIABLES neutral /METHOD=ENTER New_treatment /PRINT=GOODFIT CI(95) /CRITERIA=PIN(0.05) POUT(0.10) ITERATE(20) CUT(0.5).
Resources	Processor Time	00:00:00.00
	Elapsed Time	00:00:00.00

### Case Processing Summary

Unweighted Cases <sup>a</sup>		N	Percent
Selected Cases	Included in Analysis	407	100.0
	Missing Cases	0	.0
	Total	407	100.0
Unselected Cases		0	.0
Total		407	100.0

a. If weight is in effect, see classification table for the total number of cases.

### Dependent Variable Encoding

Original Value	Internal Value
not neutral	0
neutral	1

### Block 0: Beginning Block

#### Classification Table<sup>a,b</sup>

		Predicted		Percentage Correct
		not neutral	neutral	
Step 0	neutral	253	0	100.0
	neutral	154	0	.0
	Overall Percentage			62.2

a. Constant is included in the model.

b. The cut value is .500

#### Variables in the Equation

		B	S.E.	Wald	df	Sig.	Exp(B)
Step 0	Constant	-.496	.102	23.593	1	<.001	.609

#### Variables not in the Equation

		Score	df	Sig.
Step 0	Variables MI- New Treatment	32.960	1	<.001
	Overall Statistics	32.960	1	<.001

### Block 1: Method = Enter

#### Omnibus Tests of Model Coefficients

		Chi-square	df	Sig.
Step 1	Step	34.691	1	<.001
	Block	34.691	1	<.001
	Model	34.691	1	<.001

### Model Summary

Step	-2 Log likelihood	Cox & Snell R Square	Nagelkerke R Square
1	505.207 <sup>a</sup>	.082	.111

a. Estimation terminated at iteration number 4 because parameter estimates changed by less than .001.

### Hosmer and Lemeshow Test

Step	Chi-square	df	Sig.
1	.000	0	.

### Contingency Table for Hosmer and Lemeshow Test

		neutral = not neutral		neutral = neutral		Total
		Observed	Expected	Observed	Expected	
Step 1	1	121	121.000	30	30.000	151
	2	132	132.000	124	124.000	256

### Classification Table<sup>a</sup>

		Predicted		Percentage Correct
Observed		not neutral	neutral	
Step 1	neutral	253	0	100.0
	not neutral	154	0	.0
	Overall Percentage			62.2

a. The cut value is .500

### Variables in the Equation

		B	S.E.	Wald	df	Sig.	Exp(B)
Step 1 <sup>a</sup>	MI- New Treatment	-1.332	.239	31.001	1	<.001	.264
	Constant	-.063	.125	.250	1	.617	.939

### Variables in the Equation

		95% C.I. for EXP(B)	
		Lower	Upper
Step 1 <sup>a</sup>	MI- New Treatment	.165	.422
	Constant		

a. Variable(s) entered on step 1: MI- New Treatment.

LOGISTIC REGRESSION VARIABLESneutral  
 /METHOD=ENTER Drug\_abuse  
 /PRINT=GOODFIT CI(95)  
 /CRITERIA=PIN(0.05) POUT(0.10) ITERATE(20) CUT(0.5).

## Logistic Regression

### Notes

Output Created		28-AUG-2024 17:49:38
Comments		
Input	Data	C: \Users\apalacios\Documents\Media manuscript\Database.sav
	Active Dataset	DataSet1
	Filter	<none>
	Weight	<none>
	Split File	<none>
	N of Rows in Working Data File	407
Missing Value Handling	Definition of Missing	User-defined missing values are treated as missing
Syntax		LOGISTIC REGRESSION VARIABLES neutral /METHOD=ENTER Drug_abuse /PRINT=GOODFIT CI(95) /CRITERIA=PIN(0.05) POUT(0.10) ITERATE(20) CUT(0.5).
Resources	Processor Time	00:00:00.03
	Elapsed Time	00:00:00.01

### Case Processing Summary

Unweighted Cases <sup>a</sup>		N	Percent
Selected Cases	Included in Analysis	407	100.0
	Missing Cases	0	.0
	Total	407	100.0
Unselected Cases		0	.0
Total		407	100.0

a. If weight is in effect, see classification table for the total number of cases.



### Dependent Variable Encoding

Original Value	Internal Value
not neutral	0
neutral	1

### Block 0: Beginning Block

**Classification Table<sup>a,b</sup>**

Observed		Predicted		Percentage Correct
		not neutral	neutral	
Step 0	neutral	253	0	100.0
	not neutral	154	0	.0
	Overall Percentage			62.2

a. Constant is included in the model.

b. The cut value is .500

### Variables in the Equation

		B	S.E.	Wald	df	Sig.	Exp(B)
Step 0	Constant	-.496	.102	23.593	1	<.001	.609

### Variables not in the Equation

			Score	df	Sig.
Step 0	Variables	MI-Drug Abuse	20.453	1	<.001
	Overall Statistics		20.453	1	<.001

### Block 1: Method = Enter

#### Omnibus Tests of Model Coefficients

		Chi-square	df	Sig.
Step 1	Step	19.980	1	<.001
	Block	19.980	1	<.001
	Model	19.980	1	<.001

### Model Summary

Step	-2 Log likelihood	Cox & Snell R Square	Nagelkerke R Square
1	519.917 <sup>a</sup>	.048	.065

a. Estimation terminated at iteration number 3 because parameter estimates changed by less than .001.

### Hosmer and Lemeshow Test

Step	Chi-square	df	Sig.
1	.000	0	.

### Contingency Table for Hosmer and Lemeshow Test

		neutral = not neutral		neutral = neutral		Total
		Observed	Expected	Observed	Expected	
Step 1	1	211	211.000	98	98.000	309
	2	42	42.000	56	56.000	98

### Classification Table<sup>a</sup>

		Predicted		Percentage Correct
		Observed	neutral not neutral	
Step 1	neutral	not neutral	211	83.4
		neutral	98	36.4
	Overall Percentage			65.6

a. The cut value is .500

### Variables in the Equation

		B	S.E.	Wald	df	Sig.	Exp(B)
Step 1 <sup>a</sup>	MI-Drug Abuse	1.055	.238	19.645	1	<.001	2.871
	Constant	-.767	.122	39.357	1	<.001	.464

### Variables in the Equation

		95% C.I. for EXP(B)	
		Lower	Upper
Step 1 <sup>a</sup>	MI-Drug Abuse	1.801	4.576
	Constant		

a. Variable(s) entered on step 1: MI-Drug Abuse.

LOGISTIC REGRESSION VARIABLESneutral  
/METHOD=ENTER Health  
/PRINT=GOODFIT CI(95)  
/CRITERIA=PIN(0.05) POUT(0.10) ITERATE(20) CUT(0.5).

## Logistic Regression

### Notes

Output Created		28-AUG-2024 17:49:38
Comments		
Input	Data	C: \Users\apalacios\Documents\Media manuscript\Database.sav
	Active Dataset	DataSet1
	Filter	<none>
	Weight	<none>
	Split File	<none>
	N of Rows in Working Data File	407
Missing Value Handling	Definition of Missing	User-defined missing values are treated as missing
Syntax		LOGISTIC REGRESSION VARIABLES neutral /METHOD=ENTER Health /PRINT=GOODFIT CI(95) /CRITERIA=PIN(0.05) POUT(0.10) ITERATE(20) CUT(0.5).
Resources	Processor Time	00:00:00.00
	Elapsed Time	00:00:00.00

### Case Processing Summary

Unweighted Cases <sup>a</sup>		N	Percent
Selected Cases	Included in Analysis	407	100.0
	Missing Cases	0	.0
	Total	407	100.0
Unselected Cases		0	.0
Total		407	100.0

a. If weight is in effect, see classification table for the total number of cases.

### Dependent Variable Encoding

Original Value	Internal Value
not neutral	0
neutral	1

### Block 0: Beginning Block

**Classification Table<sup>a,b</sup>**

		Predicted		Percentage Correct
		not neutral	neutral	
Step 0	neutral	253	0	100.0
	neutral	154	0	.0
	Overall Percentage			62.2

a. Constant is included in the model.

b. The cut value is .500

### Variables in the Equation

		B	S.E.	Wald	df	Sig.	Exp(B)
Step 0	Constant	-.496	.102	23.593	1	<.001	.609

### Variables not in the Equation

		Score	df	Sig.
Step 0	Variables MI- Health	5.545	1	.019
	Overall Statistics	5.545	1	.019

### Block 1: Method = Enter

#### Omnibus Tests of Model Coefficients

		Chi-square	df	Sig.
Step 1	Step	5.396	1	.020
	Block	5.396	1	.020
	Model	5.396	1	.020

### Model Summary

Step	-2 Log likelihood	Cox & Snell R Square	Nagelkerke R Square
1	534.501 <sup>a</sup>	.013	.018

a. Estimation terminated at iteration number 3 because parameter estimates changed by less than .001.

### Hosmer and Lemeshow Test

Step	Chi-square	df	Sig.
1	.000	0	.

### Contingency Table for Hosmer and Lemeshow Test

		neutral = not neutral		neutral = neutral		Total
		Observed	Expected	Observed	Expected	
Step 1	1	225	225.000	124	124.000	349
	2	28	28.000	30	30.000	58

### Classification Table<sup>a</sup>

		Predicted		Percentage Correct
		Observed	neutral	
Step 1	neutral	not neutral	neutral	88.9
	neutral	225	28	19.5
	Overall Percentage	124	30	62.7

a. The cut value is .500

### Variables in the Equation

		B	S.E.	Wald	df	Sig.	Exp(B)	95% C.I. for
								Lower
Step 1 <sup>a</sup>	MI- Health	.665	.286	5.419	1	.020	1.944	1.111
	Constant	-.596	.112	28.380	1	<.001	.551	

### Variables in the Equation

		95% C.I. for ..
		Upper
Step 1 <sup>a</sup>	MI- Health	3.403
	Constant	

a. Variable(s) entered on step 1: MI- Health.

LOGISTIC REGRESSION VARIABLESneutral  
/METHOD=ENTER Regulation  
/PRINT=GOODFIT CI(95)  
/CRITERIA=PIN(0.05) POUT(0.10) ITERATE(20) CUT(0.5).

## Logistic Regression

### Notes

Output Created		28-AUG-2024 17:49:38
Comments		
Input	Data	C: \Users\apalacios\Document s\Media manuscript\Database.sav
	Active Dataset	DataSet1
	Filter	<none>
	Weight	<none>
	Split File	<none>
	N of Rows in Working Data File	407
Missing Value Handling	Definition of Missing	User-defined missing values are treated as missing
Syntax		LOGISTIC REGRESSION VARIABLES neutral /METHOD=ENTER Regulation /PRINT=GOODFIT CI(95) /CRITERIA=PIN(0.05) POUT(0.10) ITERATE(20) CUT(0.5).
Resources	Processor Time	00:00:00.02
	Elapsed Time	00:00:00.02

### Case Processing Summary

Unweighted Cases <sup>a</sup>		N	Percent
Selected Cases	Included in Analysis	407	100.0
	Missing Cases	0	.0
	Total	407	100.0
Unselected Cases		0	.0
Total		407	100.0

a. If weight is in effect, see classification table for the total number of cases.

**Dependent Variable  
Encoding**

Original Value	Internal Value
not neutral	0
neutral	1

**Block 0: Beginning Block**

**Classification Table<sup>a,b</sup>**

		Predicted		Percentage Correct
Observed		not neutral	neutral	
Step 0	not neutral	253	0	100.0
	neutral	154	0	.0
	Overall Percentage			62.2

a. Constant is included in the model.

b. The cut value is .500

**Variables in the Equation**

		B	S.E.	Wald	df	Sig.	Exp(B)
Step 0	Constant	-.496	.102	23.593	1	<.001	.609

**Variables not in the Equation**

		Score	df	Sig.
Step 0	Variables in the Equation	.076	1	.782
	Overall Statistics	.076	1	.782

**Block 1: Method = Enter**

**Omnibus Tests of Model Coefficients**

		Chi-square	df	Sig.
Step 1	Step	.077	1	.782
	Block	.077	1	.782
	Model	.077	1	.782

### Model Summary

Step	-2 Log likelihood	Cox & Snell R Square	Nagelkerke R Square
1	539.821 <sup>a</sup>	.000	.000

a. Estimation terminated at iteration number 3 because parameter estimates changed by less than .001.

### Hosmer and Lemeshow Test

Step	Chi-square	df	Sig.
1	.000	0	.

### Contingency Table for Hosmer and Lemeshow Test

		neutral = not neutral		neutral = neutral		Total
		Observed	Expected	Observed	Expected	
Step 1	1	37	37.000	21	21.000	58
	2	216	216.000	133	133.000	349

### Classification Table<sup>a</sup>

		Predicted		Percentage Correct
		Observed	neutral	
Step 1	neutral	not neutral	neutral	100.0
	not neutral	253	0	.0
	neutral	154	0	62.2
Overall Percentage				

a. The cut value is .500

### Variables in the Equation

		B	S.E.	Wald	df	Sig.	Exp(B)
Step 1 <sup>a</sup>	MI - Regulation	-.081	.295	.076	1	.782	.922
	Constant	-.485	.110	19.357	1	<.001	.616

### Variables in the Equation

		95% C.I. for EXP(B)	
		Lower	Upper
Step 1 <sup>a</sup>	MI - Regulation	.517	1.642
	Constant		

a. Variable(s) entered on step 1: MI - Regulation.



LOGISTIC REGRESSION VARIABLESneutral  
 /METHOD=ENTER Price  
 /PRINT=GOODFIT CI(95)  
 /CRITERIA=PIN(0.05) POUT(0.10) ITERATE(20) CUT(0.5).

## Logistic Regression

### Notes

Output Created		28-AUG-2024 17:49:38
Comments		
Input	Data	C: \Users\apalacios\Documents\Media manuscript\Database.sav
	Active Dataset	DataSet1
	Filter	<none>
	Weight	<none>
	Split File	<none>
	N of Rows in Working Data File	407
Missing Value Handling	Definition of Missing	User-defined missing values are treated as missing
Syntax		LOGISTIC REGRESSION VARIABLES neutral /METHOD=ENTER Price /PRINT=GOODFIT CI(95) /CRITERIA=PIN(0.05) POUT(0.10) ITERATE(20) CUT(0.5).
Resources	Processor Time	00:00:00.02
	Elapsed Time	00:00:00.02

### Case Processing Summary

Unweighted Cases <sup>a</sup>		N	Percent
Selected Cases	Included in Analysis	407	100.0
	Missing Cases	0	.0
	Total	407	100.0
Unselected Cases		0	.0
Total		407	100.0

a. If weight is in effect, see classification table for the total number of cases.

### Dependent Variable Encoding

Original Value	Internal Value
not neutral	0
neutral	1

### Block 0: Beginning Block

**Classification Table<sup>a,b</sup>**

Observed		Predicted		Percentage Correct
		not neutral	neutral	
Step 0	not neutral	253	0	100.0
	neutral	154	0	.0
	Overall Percentage			62.2

a. Constant is included in the model.

b. The cut value is .500

### Variables in the Equation

		B	S.E.	Wald	df	Sig.	Exp(B)
Step 0	Constant	-.496	.102	23.593	1	<.001	.609

### Variables not in the Equation

		Score	df	Sig.
Step 0	Variables in the Model	.279	1	.597
	Overall Statistics	.279	1	.597

### Block 1: Method = Enter

#### Omnibus Tests of Model Coefficients

		Chi-square	df	Sig.
Step 1	Step	.284	1	.594
	Block	.284	1	.594
	Model	.284	1	.594

### Model Summary

Step	-2 Log likelihood	Cox & Snell R Square	Nagelkerke R Square
1	539.614 <sup>a</sup>	.001	.001

a. Estimation terminated at iteration number 3 because parameter estimates changed by less than .001.

### Hosmer and Lemeshow Test

Step	Chi-square	df	Sig.
1	.000	0	.

### Contingency Table for Hosmer and Lemeshow Test

		neutral = not neutral		neutral = neutral		Total
		Observed	Expected	Observed	Expected	
Step 1	1	20	20.000	10	10.000	30
	2	233	233.000	144	144.000	377

### Classification Table<sup>a</sup>

		Predicted		Percentage Correct
		Observed	neutral	
Step 1	neutral	not neutral	neutral	100.0
	neutral	253	0	.0
	Overall Percentage	154	0	62.2

a. The cut value is .500

### Variables in the Equation

		B	S.E.	Wald	df	Sig.	Exp(B)	95% C.I. for
								Lower
Step 1 <sup>a</sup>	MI - Price	-.212	.402	.279	1	.598	.809	.368
	Constant	-.481	.106	20.610	1	<.001	.618	

### Variables in the Equation

		95% C.I. for ..
		Upper
Step 1 <sup>a</sup>	MI - Price	1.777
	Constant	

a. Variable(s) entered on step 1: MI - Price.

LOGISTIC REGRESSION VARIABLESneutral  
/METHOD=ENTER Business\_news  
/PRINT=GOODFIT CI(95)  
/CRITERIA=PIN(0.05) POUT(0.10) ITERATE(20) CUT(0.5).

## Logistic Regression

### Notes

Output Created		28-AUG-2024 17:49:38
Comments		
Input	Data	C: \Users\apalacios\Documents\Media manuscript\Database.sav
	Active Dataset	DataSet1
	Filter	<none>
	Weight	<none>
	Split File	<none>
	N of Rows in Working Data File	407
Missing Value Handling	Definition of Missing	User-defined missing values are treated as missing
Syntax		LOGISTIC REGRESSION VARIABLES neutral /METHOD=ENTER Business_news /PRINT=GOODFIT CI(95) /CRITERIA=PIN(0.05) POUT(0.10) ITERATE(20) CUT(0.5).
Resources	Processor Time	00:00:00.02
	Elapsed Time	00:00:00.02

### Case Processing Summary

Unweighted Cases <sup>a</sup>		N	Percent
Selected Cases	Included in Analysis	407	100.0
	Missing Cases	0	.0
	Total	407	100.0
Unselected Cases		0	.0
Total		407	100.0

a. If weight is in effect, see classification table for the total number of cases.

### Dependent Variable Encoding

Original Value	Internal Value
not neutral	0
neutral	1

### Block 0: Beginning Block

#### Classification Table<sup>a,b</sup>

		Predicted		Percentage Correct
		not neutral	neutral	
Step 0	Observed not neutral	253	0	100.0
	neutral	154	0	.0
	Overall Percentage			62.2

a. Constant is included in the model.

b. The cut value is .500

#### Variables in the Equation

		B	S.E.	Wald	df	Sig.	Exp(B)
Step 0	Constant	-.496	.102	23.593	1	<.001	.609

#### Variables not in the Equation

		Score	df	Sig.
Step 0	Variables Mi - Business News	2.208	1	.137
	Overall Statistics	2.208	1	.137

### Block 1: Method = Enter

#### Omnibus Tests of Model Coefficients

		Chi-square	df	Sig.
Step 1	Step	2.125	1	.145
	Block	2.125	1	.145
	Model	2.125	1	.145

### Model Summary

Step	-2 Log likelihood	Cox & Snell R Square	Nagelkerke R Square
1	537.773 <sup>a</sup>	.005	.007

a. Estimation terminated at iteration number 3 because parameter estimates changed by less than .001.

### Hosmer and Lemeshow Test

Step	Chi-square	df	Sig.
1	.000	0	.

### Contingency Table for Hosmer and Lemeshow Test

		neutral = not neutral		neutral = neutral		Total
		Observed	Expected	Observed	Expected	
Step 1	1	253	253.000	154	154.000	407

### Variables in the Equation

		B	S.E.	Wald	df	Sig.	Exp(B)
Step 1 <sup>a</sup>	Mi - Business News	.859	.595	2.088	1	.148	2.362
	Constant	-.523	.104	25.245	1	<.001	.593

### Variables in the Equation

		95% C.I. for EXP(B)	
		Lower	Upper
Step 1 <sup>a</sup>	Mi - Business News	.736	7.577
	Constant		

a. Variable(s) entered on step 1: Mi - Business News.

### Classification Table<sup>a</sup>

			Predicted		Percentage Correct
			not neutral	neutral	
Step 1	Observed		248	5	98.0
	neutral	not neutral			
		neutral	147	7	4.5
	Overall Percentage				62.7

a. The cut value is .500

SORT CASES BY PropagandYES (D).  
 SORT CASES BY Propaganda (D).  
 LOGISTIC REGRESSION VARIABLESneutral  
 /METHOD=ENTER PropagandNO  
 /PRINT=GOODFIT CI(95)  
 /CRITERIA=PIN(0.05) POUT(0.10) ITERATE(20) CUT(0.5).

## Logistic Regression

### Notes

Output Created		28-AUG-2024 17:57:23
Comments		
Input	Data	C: \Users\apalacios\Documents\Media manuscript\Database.sav
	Active Dataset	DataSet1
	Filter	<none>
	Weight	<none>
	Split File	<none>
	N of Rows in Working Data File	407
Missing Value Handling	Definition of Missing	User-defined missing values are treated as missing
Syntax		LOGISTIC REGRESSION VARIABLES neutral /METHOD=ENTER PropagandNO /PRINT=GOODFIT CI(95) /CRITERIA=PIN(0.05) POUT(0.10) ITERATE(20) CUT(0.5).
Resources	Processor Time	00:00:00.00
	Elapsed Time	00:00:00.00

### Case Processing Summary

Unweighted Cases <sup>a</sup>		N	Percent
Selected Cases	Included in Analysis	407	100.0
	Missing Cases	0	.0
	Total	407	100.0
Unselected Cases		0	.0
Total		407	100.0

a. If weight is in effect, see classification table for the total number of cases.

### Dependent Variable Encoding

Original Value	Internal Value
not neutral	0
neutral	1

### Block 0: Beginning Block

#### Classification Table<sup>a,b</sup>

Observed		Predicted		Percentage Correct
		not neutral	neutral	
Step 0	not neutral	253	0	100.0
	neutral	154	0	.0
	Overall Percentage			62.2

a. Constant is included in the model.

b. The cut value is .500

#### Variables in the Equation

		B	S.E.	Wald	df	Sig.	Exp(B)
Step 0	Constant	-.496	.102	23.593	1	<.001	.609

#### Variables not in the Equation

		Score	df	Sig.
Step 0	Variables PropagandNO	53.165	1	<.001
	Overall Statistics	53.165	1	<.001

### Block 1: Method = Enter

#### Omnibus Tests of Model Coefficients

		Chi-square	df	Sig.
Step 1	Step	59.511	1	<.001
	Block	59.511	1	<.001
	Model	59.511	1	<.001



### Model Summary

Step	-2 Log likelihood	Cox & Snell R Square	Nagelkerke R Square
1	480.386 <sup>a</sup>	.136	.185

a. Estimation terminated at iteration number 5 because parameter estimates changed by less than .001.

### Hosmer and Lemeshow Test

Step	Chi-square	df	Sig.
1	.000	0	.

### Contingency Table for Hosmer and Lemeshow Test

		neutral = not neutral		neutral = neutral		Total
		Observed	Expected	Observed	Expected	
Step 1	1	112	112.000	15	15.000	127
	2	141	141.000	139	139.000	280

### Classification Table<sup>a</sup>

		Predicted		Percentage Correct
Observed		not neutral	neutral	
Step 1	neutral	253	0	100.0
	not neutral	154	0	.0
	Overall Percentage			62.2

a. The cut value is .500

### Variables in the Equation

		B	S.E.	Wald	df	Sig.	Exp(B)
Step 1 <sup>a</sup>	PropagandNO	1.996	.300	44.332	1	<.001	7.361
	Constant	-2.010	.275	53.468	1	<.001	.134

### Variables in the Equation

		95% C.I. for EXP(B)	
		Lower	Upper
Step 1 <sup>a</sup>	PropagandNO	4.090	13.247
	Constant		

a. Variable(s) entered on step 1: PropagandNO.

DATASETACTIVATEDataSet1.

SAVEOUTFILE='C:\Users\apalacios\Documents\Media manuscript\Database.sav '  
/COMPRESSED.  
LOGISTIC REGRESSION VARIABLESnneutral  
/METHOD=ENTER Propaganda  
/PRINT=GOODFIT CI(95)  
/CRITERIA=PIN(0.05) POUT(0.10) ITERATE(20) CUT(0.5).

## Logistic Regression

### Notes

Output Created		28-AUG-2024 17:58:58
Comments		
Input	Data	C: \Users\apalacios\Document s\Media manuscript\Database.sav
	Active Dataset	DataSet1
	Filter	<none>
	Weight	<none>
	Split File	<none>
	N of Rows in Working Data File	407
Missing Value Handling	Definition of Missing	User-defined missing values are treated as missing
Syntax		LOGISTIC REGRESSION VARIABLES neutral /METHOD=ENTER Propaganda /PRINT=GOODFIT CI(95) /CRITERIA=PIN(0.05) POUT(0.10) ITERATE(20) CUT(0.5).
Resources	Processor Time	00:00:00.00
	Elapsed Time	00:00:00.00

### Case Processing Summary

Unweighted Cases <sup>a</sup>		N	Percent
Selected Cases	Included in Analysis	407	100.0
	Missing Cases	0	.0
	Total	407	100.0
Unselected Cases		0	.0
Total		407	100.0

a. If weight is in effect, see classification table for the total number of cases.

### Dependent Variable Encoding

Original Value	Internal Value
not neutral	0
neutral	1

### Block 0: Beginning Block

#### Classification Table<sup>a,b</sup>

Observed		Predicted		Percentage Correct
		not neutral	neutral	
Step 0	neutral	253	0	100.0
	not neutral	154	0	.0
Overall Percentage				62.2

a. Constant is included in the model.

b. The cut value is .500

### Variables in the Equation

		B	S.E.	Wald	df	Sig.	Exp(B)
Step 0	Constant	-.496	.102	23.593	1	<.001	.609

### Variables not in the Equation

			Score	df	Sig.
Step 0	Variables	propaganda yes or maybe	53.165	1	<.001
	Overall Statistics		53.165	1	<.001

### Block 1: Method = Enter

### Omnibus Tests of Model Coefficients

		Chi-square	df	Sig.
Step 1	Step	59.511	1	<.001
	Block	59.511	1	<.001
	Model	59.511	1	<.001

### Model Summary

Step	-2 Log likelihood	Cox & Snell R Square	Nagelkerke R Square
1	480.386 <sup>a</sup>	.136	.185

a. Estimation terminated at iteration number 5 because parameter estimates changed by less than .001.

### Hosmer and Lemeshow Test

Step	Chi-square	df	Sig.
1	.000	0	.

### Contingency Table for Hosmer and Lemeshow Test

		neutral = not neutral		neutral = neutral		Total
		Observed	Expected	Observed	Expected	
Step 1	1	112	112.000	15	15.000	127
	2	141	141.000	139	139.000	280

### Classification Table<sup>a</sup>

			Predicted		Percentage Correct
Observed		not neutral	neutral		
Step 1	neutral	not neutral	253	0	100.0
		neutral	154	0	.0
	Overall Percentage				62.2

a. The cut value is .500

### Variables in the Equation

		B	S.E.	Wald	df	Sig.
Step 1 <sup>a</sup>	propaganda yes or maybe	-1.996	.300	44.332	1	<.001
	Constant	-.014	.120	.014	1	.905

### Variables in the Equation

		Exp(B)	95% C.I. for EXP(B)	
			Lower	Upper
Step 1 <sup>a</sup>	propaganda yes or maybe	.136	.075	.244
	Constant	.986		

a. Variable(s) entered on step 1: propaganda yes or maybe.

```
LOGISTIC REGRESSION VARIABLES New_treatment
/METHOD=ENTER adversereact
/PRINT=GOODFIT CI(95)
/CRITERIA=PIN(0.05) POUT(0.10) ITERATE(20) CUT(0.5).
```

### Logistic Regression

#### Notes

Output Created		28-AUG-2024 18:51:25
Comments		
Input	Data	C: \Users\apalacios\Documents\Media manuscript\Database.sav
	Active Dataset	DataSet1
	Filter	<none>
	Weight	<none>
	Split File	<none>
	N of Rows in Working Data File	407
Missing Value Handling	Definition of Missing	User-defined missing values are treated as missing
Syntax		LOGISTIC REGRESSION VARIABLES New_treatment /METHOD=ENTER adversereact /PRINT=GOODFIT CI(95) /CRITERIA=PIN(0.05) POUT(0.10) ITERATE(20) CUT(0.5).
Resources	Processor Time	00:00:00.00
	Elapsed Time	00:00:00.00

### Case Processing Summary

Unweighted Cases <sup>a</sup>		N	Percent
Selected Cases	Included in Analysis	407	100.0
	Missing Cases	0	.0
	Total	407	100.0
Unselected Cases		0	.0
Total		407	100.0

a. If weight is in effect, see classification table for the total number of cases.

### Dependent Variable Encoding

Original Value	Internal Value
Not new	0
New treatment	1

### Block 0: Beginning Block

#### Classification Table<sup>a,b</sup>

Observed			Predicted		Percentage Correct
			MI- New Treatment Not new	New treatment	
Step 0	MI- New Treatment	Not new	256	0	100.0
		New treatment	151	0	.0
	Overall Percentage				62.9

a. Constant is included in the model.

b. The cut value is .500

### Variables in the Equation

		B	S.E.	Wald	df	Sig.	Exp(B)
Step 0	Constant	-.528	.103	26.468	1	<.001	.590

### Variables not in the Equation

			Score	df	Sig.
Step 0	Variables	adversereact	15.223	1	<.001
Overall Statistics			15.223	1	<.001

### Block 1: Method = Enter

### Omnibus Tests of Model Coefficients

		Chi-square	df	Sig.
Step 1	Step	16.160	1	<.001
	Block	16.160	1	<.001
	Model	16.160	1	<.001

### Model Summary

Step	-2 Log likelihood	Cox & Snell R Square	Nagelkerke R Square
1	520.665 <sup>a</sup>	.039	.053

a. Estimation terminated at iteration number 4 because parameter estimates changed by less than .001.

### Hosmer and Lemeshow Test

Step	Chi-square	df	Sig.
1	.000	0	.

### Contingency Table for Hosmer and Lemeshow Test

		MI- New Treatment = Not new		MI- New Treatment = New treatment		Total
		Observed	Expected	Observed	Expected	
Step 1	1	82	82.000	22	22.000	104
	2	174	174.000	129	129.000	303

### Classification Table<sup>a</sup>

		Predicted		Percentage Correct
		MI- New Treatment Not new	MI- New Treatment New treatment	
Step 1	Observed MI- New Treatment	Not new	New treatment	
	Not new	256	0	100.0
	New treatment	151	0	.0
	Overall Percentage			62.9

a. The cut value is .500

### Variables in the Equation

		B	S.E.	Wald	df	Sig.	Exp(B)	95% C.I. for Lower
Step 1 <sup>a</sup>	adversereact	-1.016	.267	14.521	1	<.001	.362	.215
	Constant	-.299	.116	6.634	1	.010	.741	

## Variables in the Equation

		95% C.I. for ..
		Upper
Step 1 <sup>a</sup>	adversereact	.610
	Constant	

a. Variable(s) entered on step 1: adversereact.

LOGISTIC REGRESSION VARIABLES New\_treatment  
 /METHOD=ENTER price\_included  
 /PRINT=GOODFIT CI(95)  
 /CRITERIA=PIN(0.05) POUT(0.10) ITERATE(20) CUT(0.5).

## Logistic Regression

### Notes

Output Created		28-AUG-2024 18:51:25
Comments		
Input	Data	C: \Users\apalacios\Documents\Media manuscript\Database.sav
	Active Dataset	DataSet1
	Filter	<none>
	Weight	<none>
	Split File	<none>
	N of Rows in Working Data File	407
Missing Value Handling	Definition of Missing	User-defined missing values are treated as missing
Syntax		LOGISTIC REGRESSION VARIABLES New_treatment /METHOD=ENTER price_included /PRINT=GOODFIT CI(95) /CRITERIA=PIN(0.05) POUT(0.10) ITERATE(20) CUT(0.5).
Resources	Processor Time	00:00:00.00
	Elapsed Time	00:00:00.00



### Case Processing Summary

Unweighted Cases <sup>a</sup>		N	Percent
Selected Cases	Included in Analysis	384	94.3
	Missing Cases	23	5.7
	Total	407	100.0
Unselected Cases		0	.0
Total		407	100.0

a. If weight is in effect, see classification table for the total number of cases.

### Dependent Variable Encoding

Original Value	Internal Value
Not new	0
New treatment	1

### Block 0: Beginning Block

#### Classification Table<sup>a,b</sup>

Observed			Predicted		Percentage Correct
			MI- New Treatment Not new	New treatment	
Step 0	MI- New Treatment	Not new	240	0	100.0
		New treatment	144	0	.0
	Overall Percentage				62.5

a. Constant is included in the model.

b. The cut value is .500

### Variables in the Equation

		B	S.E.	Wald	df	Sig.	Exp(B)
Step 0	Constant	-.511	.105	23.485	1	<.001	.600

### Variables not in the Equation

			Score	df	Sig.
Step 0	Variables	price_included	.040	1	.841
Overall Statistics			.040	1	.841

### Block 1: Method = Enter

### Omnibus Tests of Model Coefficients

		Chi-square	df	Sig.
Step 1	Step	.040	1	.841
	Block	.040	1	.841
	Model	.040	1	.841

### Model Summary

Step	-2 Log likelihood	Cox & Snell R Square	Nagelkerke R Square
1	508.040 <sup>a</sup>	.000	.000

a. Estimation terminated at iteration number 3 because parameter estimates changed by less than .001.

### Hosmer and Lemeshow Test

Step	Chi-square	df	Sig.
1	.000	0	.

### Contingency Table for Hosmer and Lemeshow Test

		MI- New Treatment = Not new		MI- New Treatment = New treatment		Total
		Observed	Expected	Observed	Expected	
Step 1	1	47	47.000	27	27.000	74
	2	193	193.000	117	117.000	310

### Classification Table<sup>a</sup>

			Predicted		Percentage Correct
Observed		MI- New Treatment Not new	New treatment		
Step 1	MI- New Treatment	Not new	240	0	100.0
		New treatment	144	0	.0
	Overall Percentage				62.5

a. The cut value is .500

### Variables in the Equation

		B	S.E.	Wald	df	Sig.	Exp(B)
Step 1 <sup>a</sup>	price_included	-.054	.268	.040	1	.841	.948
	Constant	-.501	.117	18.248	1	<.001	.606

### Variables in the Equation

		95% C.I. for EXP(B)	
		Lower	Upper
Step 1 <sup>a</sup>	price_included	.560	1.604
	Constant		

a. Variable(s) entered on step 1: price\_included.

LOGISTIC REGRESSION VARIABLES legacy\_digital  
 /METHOD=ENTER neutral  
 /PRINT=GOODFIT CI(95)  
 /CRITERIA=PIN(0.05) POUT(0.10) ITERATE(20) CUT(0.5).

### Logistic Regression

#### Notes

Output Created		30-AUG-2024 15:50:32
Comments		
Input	Data	C: \Users\apalacios\Documents\Media manuscript\Database.sav
	Active Dataset	DataSet1
	Filter	<none>
	Weight	<none>
	Split File	<none>
	N of Rows in Working Data File	407
Missing Value Handling	Definition of Missing	User-defined missing values are treated as missing
Syntax		LOGISTIC REGRESSION VARIABLES legacy_digital /METHOD=ENTER neutral /PRINT=GOODFIT CI(95) /CRITERIA=PIN(0.05) POUT(0.10) ITERATE(20) CUT(0.5).
Resources	Processor Time	00:00:00.00
	Elapsed Time	00:00:00.01

### Case Processing Summary

Unweighted Cases <sup>a</sup>		N	Percent
Selected Cases	Included in Analysis	407	100.0
	Missing Cases	0	.0
	Total	407	100.0
Unselected Cases		0	.0
Total		407	100.0

a. If weight is in effect, see classification table for the total number of cases.

### Dependent Variable Encoding

Original Value	Internal Value
legacy	0
digital	1

### Block 0: Beginning Block

#### Classification Table<sup>a,b</sup>

Observed			Predicted		Percentage Correct
			legacy_digital legacy	digital	
Step 0	legacy_digital	legacy	0	199	.0
		digital	0	208	100.0
	Overall Percentage				51.1

a. Constant is included in the model.

b. The cut value is .500

### Variables in the Equation

	B	S.E.	Wald	df	Sig.	Exp(B)
Step 0 Constant	.044	.099	.199	1	.656	1.045

### Variables not in the Equation

	Score	df	Sig.
Step 0 Variables neutral	.455	1	.500
Overall Statistics	.455	1	.500

### Block 1: Method = Enter

### Omnibus Tests of Model Coefficients

		Chi-square	df	Sig.
Step 1	Step	.455	1	.500
	Block	.455	1	.500
	Model	.455	1	.500

### Model Summary

Step	-2 Log likelihood	Cox & Snell R Square	Nagelkerke R Square
1	563.568 <sup>a</sup>	.001	.001

a. Estimation terminated at iteration number 2 because parameter estimates changed by less than .001.

### Hosmer and Lemeshow Test

Step	Chi-square	df	Sig.
1	.000	0	.

### Contingency Table for Hosmer and Lemeshow Test

		legacy_digital = legacy		legacy_digital = digital		Total
		Observed	Expected	Observed	Expected	
Step 1	1	127	127.000	126	126.000	253
	2	72	72.000	82	82.000	154

### Classification Table<sup>a</sup>

			Predicted		Percentage Correct
			legacy_digital legacy	digital	
Step 1	Observed				
	legacy_digital	legacy	127	72	63.8
		digital	126	82	39.4
Overall Percentage					51.4

a. The cut value is .500

### Variables in the Equation

		B	S.E.	Wald	df	Sig.	Exp(B)	95% C.I. for Lower
Step 1 <sup>a</sup>	neutral	.138	.205	.454	1	.500	1.148	.769
	Constant	-.008	.126	.004	1	.950	.992	

## Variables in the Equation

		95% C.I. for ..
		Upper
Step 1 <sup>a</sup>	neutral	1.715
	Constant	

a. Variable(s) entered on step 1: neutral.

DATASETACTIVATEDataSet1.

SAVEOUTFILE='C:\Users\apalacios\Documents\Media manuscript\Database.sav '  
/COMPRESSED.

DATASETACTIVATEDataSet1.

SAVEOUTFILE='C:\Users\apalacios\Documents\Media manuscript\Database.sav '  
/COMPRESSED.

LOGISTIC REGRESSION VARIABLESlegacy

/METHOD=ENTER neutral

/PRINT=GOODFIT CI(95)

/CRITERIA=PIN(0.05) POUT(0.10) ITERATE(20) CUT(0.5).

## Logistic Regression

### Notes

Output Created		30-AUG-2024 15:58:22
Comments		
Input	Data	C: \Users\apalacios\Document s\Media manuscript\Database.sav
	Active Dataset	DataSet1
	Filter	<none>
	Weight	<none>
	Split File	<none>
	N of Rows in Working Data File	407
Missing Value Handling	Definition of Missing	User-defined missing values are treated as missing
Syntax		LOGISTIC REGRESSION VARIABLES legacy /METHOD=ENTER neutral /PRINT=GOODFIT CI(95) /CRITERIA=PIN(0.05) POUT(0.10) ITERATE(20) CUT(0.5).

### Notes

Resources	Processor Time	00:00:00.00
	Elapsed Time	00:00:00.00

### Case Processing Summary

Unweighted Cases <sup>a</sup>		N	Percent
Selected Cases	Included in Analysis	407	100.0
	Missing Cases	0	.0
	Total	407	100.0
Unselected Cases		0	.0
Total		407	100.0

a. If weight is in effect, see classification table for the total number of cases.

### Dependent Variable Encoding

Original Value	Internal Value
Digital	0
Legacy	1

### Block 0: Beginning Block

#### Classification Table<sup>a,b</sup>

			Predicted		Percentage Correct
			legacy		
Step 0	Observed	Digital	Digital	Legacy	
	legacy	Digital	208	0	100.0
		Legacy	199	0	.0
	Overall Percentage				51.1

a. Constant is included in the model.

b. The cut value is .500

### Variables in the Equation

		B	S.E.	Wald	df	Sig.	Exp(B)
Step 0	Constant	-.044	.099	.199	1	.656	.957

### Variables not in the Equation

		Score	df	Sig.
Step 0	Variables: neutral	.455	1	.500
	Overall Statistics	.455	1	.500

### Block 1: Method = Enter

#### Omnibus Tests of Model Coefficients

		Chi-square	df	Sig.
Step 1	Step	.455	1	.500
	Block	.455	1	.500
	Model	.455	1	.500

#### Model Summary

Step	-2 Log likelihood	Cox & Snell R Square	Nagelkerke R Square
1	563.568 <sup>a</sup>	.001	.001

a. Estimation terminated at iteration number 2 because parameter estimates changed by less than .001.

#### Hosmer and Lemeshow Test

Step	Chi-square	df	Sig.
1	.000	0	.

#### Contingency Table for Hosmer and Lemeshow Test

		legacy = Digital		legacy = Legacy		Total
		Observed	Expected	Observed	Expected	
Step 1	1	82	82.000	72	72.000	154
	2	126	126.000	127	127.000	253

#### Classification Table<sup>a</sup>

			Predicted		Percentage Correct
			legacy		
Observed	Digital	Legacy			
	Digital	Legacy			
Step 1	legacy	Digital	82	126	39.4
		Legacy	72	127	63.8
	Overall Percentage				51.4

a. The cut value is .500



### Variables in the Equation

		B	S.E.	Wald	df	Sig.	Exp(B)	95% C.I. for Lower
Step 1 <sup>a</sup>	neutral	-.138	.205	.454	1	.500	.871	.583
	Constant	.008	.126	.004	1	.950	1.008	

### Variables in the Equation

		95% C.I. for .. Upper
Step 1 <sup>a</sup>	neutral	1.301
	Constant	

a. Variable(s) entered on step 1: neutral.

LOGISTIC REGRESSION VARIABLES digital  
 /METHOD=ENTER PropagandYES  
 /PRINT=GOODFIT CI(95)  
 /CRITERIA=PIN(0.05) POUT(0.10) ITERATE(20) CUT(0.5).

### Logistic Regression

#### Notes

Output Created	06-SEP-2024 08:50:47	
Comments		
Input	Data	C: \Users\lapalacios\Documents\Media manuscript\Database.sav
	Active Dataset	DataSet1
	Filter	<none>
	Weight	<none>
	Split File	<none>
	N of Rows in Working Data File	407
Missing Value Handling	Definition of Missing	User-defined missing values are treated as missing
Syntax	LOGISTIC REGRESSION VARIABLES digital /METHOD=ENTER PropagandYES /PRINT=GOODFIT CI(95) /CRITERIA=PIN(0.05) POUT(0.10) ITERATE(20) CUT(0.5).	

### Notes

Resources	Processor Time	00:00:00.00
	Elapsed Time	00:00:00.00

### Case Processing Summary

Unweighted Cases <sup>a</sup>		N	Percent
Selected Cases	Included in Analysis	407	100.0
	Missing Cases	0	.0
	Total	407	100.0
Unselected Cases		0	.0
Total		407	100.0

a. If weight is in effect, see classification table for the total number of cases.

### Dependent Variable Encoding

Original Value	Internal Value
legacy	0
digital	1

### Block 0: Beginning Block

### Classification Table<sup>a,b</sup>

Observed		Predicted		Percentage Correct
		legacy	digital	
Step 0	digital	0	199	.0
	legacy	0	208	100.0
	Overall Percentage			51.1

a. Constant is included in the model.

b. The cut value is .500

### Variables in the Equation

	B	S.E.	Wald	df	Sig.	Exp(B)
Step 0 Constant	.044	.099	.199	1	.656	1.045

### Variables not in the Equation

			Score	df	Sig.
Step 0	Variables	PropagandYES	2.915	1	.088
	Overall Statistics		2.915	1	.088

### Block 1: Method = Enter

#### Omnibus Tests of Model Coefficients

		Chi-square	df	Sig.
Step 1	Step	2.921	1	.087
	Block	2.921	1	.087
	Model	2.921	1	.087

#### Model Summary

Step	-2 Log likelihood	Cox & Snell R Square	Nagelkerke R Square
1	561.102 <sup>a</sup>	.007	.010

a. Estimation terminated at iteration number 3 because parameter estimates changed by less than .001.

#### Hosmer and Lemeshow Test

Step	Chi-square	df	Sig.
1	.000	0	.

#### Contingency Table for Hosmer and Lemeshow Test

		digital = legacy		digital = digital		Total
		Observed	Expected	Observed	Expected	
Step 1	1	47	47.000	35	35.000	82
	2	152	152.000	173	173.000	325

#### Classification Table<sup>a</sup>

		Predicted		Percentage Correct
		digital	digital	
Observed		legacy	digital	
Step 1	digital	47	152	23.6
	legacy	35	173	83.2
	Overall Percentage			54.1

a. The cut value is .500

### Variables in the Equation

		B	S.E.	Wald	df	Sig.	Exp(B)
Step 1 <sup>a</sup>	PropagandYES	-.424	.249	2.893	1	.089	.654
	Constant	.129	.111	1.355	1	.244	1.138

### Variables in the Equation

		95% C.I. for EXP(B)	
		Lower	Upper
Step 1 <sup>a</sup>	PropagandYES	.401	1.067
	Constant		

a. Variable(s) entered on step 1: PropagandYES.

LOGISTIC REGRESSION VARIABLESdigital  
 /METHOD=ENTER adversereact  
 /PRINT=GOODFIT CI(95)  
 /CRITERIA=PIN(0.05) POUT(0.10) ITERATE(20) CUT(0.5).

### Logistic Regression

## Notes

Output Created		06-SEP-2024 08:50:47
Comments		
Input	Data	C:\Users\apalacios\Documents\Media manuscript\Database.sav
	Active Dataset	DataSet1
	Filter	<none>
	Weight	<none>
	Split File	<none>
	N of Rows in Working Data File	407
Missing Value Handling	Definition of Missing	User-defined missing values are treated as missing
Syntax		LOGISTIC REGRESSION VARIABLES digital /METHOD=ENTER adversereact /PRINT=GOODFIT CI(95) /CRITERIA=PIN(0.05) POUT(0.10) ITERATE(20) ...
Resources	Processor Time	00:00:00.02
	Elapsed Time	00:00:00.02

## Case Processing Summary

Unweighted Cases <sup>a</sup>		N	Percent
Selected Cases	Included in Analysis	407	100.0
	Missing Cases	0	.0
	Total	407	100.0
Unselected Cases		0	.0
Total		407	100.0

a. If weight is in effect, see classification table for the total number of cases.

## Dependent Variable Encoding

Original Value	Internal Value
legacy	0
digital	1

## Block 0: Beginning Block

**Classification Table<sup>a,b</sup>**

Observed		Predicted		Percentage Correct
		legacy	digital	
Step 0	digital	0	199	.0
	legacy	0	208	100.0
	Overall Percentage			51.1

a. Constant is included in the model.

b. The cut value is .500

**Variables in the Equation**

		B	S.E.	Wald	df	Sig.	Exp(B)
Step 0	Constant	.044	.099	.199	1	.656	1.045

**Variables not in the Equation**

			Score	df	Sig.
Step 0	Variables	adversereact	1.955	1	.162
	Overall Statistics		1.955	1	.162

## Block 1: Method = Enter

**Omnibus Tests of Model Coefficients**

		Chi-square	df	Sig.
Step 1	Step	1.956	1	.162
	Block	1.956	1	.162
	Model	1.956	1	.162

**Model Summary**

Step	-2 Log likelihood	Cox & Snell R Square	Nagelkerke R Square
1	562.067 <sup>a</sup>	.005	.006

a. Estimation terminated at iteration number 2 because parameter estimates changed by less than .001.

**Hosmer and Lemeshow Test**

Step	Chi-square	df	Sig.
1	.000	0	.

### Contingency Table for Hosmer and Lemeshow Test

		digital = legacy		digital = digital		Total
		Observed	Expected	Observed	Expected	
Step 1	1	57	57.000	47	47.000	104
	2	142	142.000	161	161.000	303

### Classification Table<sup>a</sup>

		Predicted		Percentage Correct
		digital	legacy	
Step 1	Observed	digital	legacy	
	digital	57	142	28.6
	legacy	47	161	77.4
Overall Percentage				53.6

a. The cut value is .500

### Variables in the Equation

		B	S.E.	Wald	df	Sig.	Exp(B)	95% C.I. for
								Lower
Step 1 <sup>a</sup>	adversereact	-.318	.228	1.948	1	.163	.727	.465
	Constant	.126	.115	1.190	1	.275	1.134	

### Variables in the Equation

		95% C.I. for ..
		Upper
Step 1 <sup>a</sup>	adversereact	1.137
	Constant	

a. Variable(s) entered on step 1: adversereact.

```
LOGISTIC REGRESSION VARIABLESNew_treatment
/METHOD=ENTER propaganda
/PRINT=GOODFIT CI(95)
/CRITERIA=PIN(0.05) POUT(0.10) ITERATE(20) CUT(0.5).
```

### Logistic Regression

## Notes

Output Created		06-SEP-2024 09:32:50
Comments		
Input	Data	C: \Users\apalacios\Documents\Media manuscript\Database.sav
	Active Dataset	DataSet1
	Filter	<none>
	Weight	<none>
	Split File	<none>
	N of Rows in Working Data File	407
Missing Value Handling	Definition of Missing	User-defined missing values are treated as missing
Syntax		LOGISTIC REGRESSION VARIABLES New_treatment /METHOD=ENTER propaganda /PRINT=GOODFIT CI(95) /CRITERIA=PIN(0.05) POUT(0.10) ITERATE(20) CUT(0.5).
Resources	Processor Time	00:00:00.00
	Elapsed Time	00:00:00.00

## Case Processing Summary

Unweighted Cases <sup>a</sup>		N	Percent
Selected Cases	Included in Analysis	407	100.0
	Missing Cases	0	.0
	Total	407	100.0
Unselected Cases		0	.0
Total		407	100.0

a. If weight is in effect, see classification table for the total number of cases.

## Dependent Variable Encoding

Original Value	Internal Value
Not new	0
New treatment	1



## Block 0: Beginning Block

**Classification Table<sup>a,b</sup>**

Observed			Predicted		Percentage Correct
			MI- New Treatment Not new	New treatment	
Step 0	MI- New Treatment	Not new	256	0	100.0
		New treatment	151	0	.0
	Overall Percentage				62.9

a. Constant is included in the model.

b. The cut value is .500

**Variables in the Equation**

		B	S.E.	Wald	df	Sig.	Exp(B)
Step 0	Constant	-.528	.103	26.468	1	<.001	.590

**Variables not in the Equation**

			Score	df	Sig.
Step 0	Variables	propaganda yes or maybe	86.032	1	<.001
	Overall Statistics		86.032	1	<.001

## Block 1: Method = Enter

**Omnibus Tests of Model Coefficients**

		Chi-square	df	Sig.
Step 1	Step	85.757	1	<.001
	Block	85.757	1	<.001
	Model	85.757	1	<.001

**Model Summary**

Step	-2 Log likelihood	Cox & Snell R Square	Nagelkerke R Square
1	451.067 <sup>a</sup>	.190	.259

a. Estimation terminated at iteration number 4 because parameter estimates changed by less than .001.

**Hosmer and Lemeshow Test**

Step	Chi-square	df	Sig.
1	.000	0	.

### Contingency Table for Hosmer and Lemeshow Test

		MI- New Treatment = Not new		MI- New Treatment = New treatment		Total
		Observed	Expected	Observed	Expected	
Step 1	1	218	218.000	62	62.000	280
	2	38	38.000	89	89.000	127

### Classification Table<sup>a</sup>

		Predicted		Percentage Correct
		MI- New Treatment Not new	MI- New Treatment New treatment	
Observed				
Step 1	MI- New Treatment Not new	218	38	85.2
	MI- New Treatment New treatment	62	89	58.9
Overall Percentage				75.4

a. The cut value is .500

### Variables in the Equation

		B	S.E.	Wald	df	Sig.
Step 1 <sup>a</sup>	propaganda yes or maybe	2.108	.241	76.292	1	<.001
	Constant	-1.257	.144	76.315	1	<.001

### Variables in the Equation

		Exp(B)	95% C.I. for EXP(B)	
			Lower	Upper
Step 1 <sup>a</sup>	propaganda yes or maybe	8.235	5.131	13.217
	Constant	.284		

a. Variable(s) entered on step 1: propaganda yes or maybe.

LOGISTIC REGRESSION VARIABLES shared\_in\_media  
 /METHOD=ENTER Spanish  
 /PRINT=GOODFIT CI(95)  
 /CRITERIA=PIN(0.05) POUT(0.10) ITERATE(20) CUT(0.5).

### Logistic Regression

## Notes

Output Created		10-SEP-2024 21:15:15
Comments		
Input	Data	C:\Users\apalacios\Documents\Media manuscript\Database.sav
	Active Dataset	DataSet1
	Filter	<none>
	Weight	<none>
	Split File	<none>
	N of Rows in Working Data File	407
Missing Value Handling	Definition of Missing	User-defined missing values are treated as missing
Syntax		LOGISTIC REGRESSION VARIABLES shared_in_media /METHOD=ENTER Spanish /PRINT=GOODFIT CI(95) /CRITERIA=PIN(0.05) POUT(0.10) ITERATE(20) CUT(0.5).
Resources	Processor Time	00:00:00.00
	Elapsed Time	00:00:00.00

## Case Processing Summary

Unweighted Cases <sup>a</sup>		N	Percent
Selected Cases	Included in Analysis	406	99.8
	Missing Cases	1	.2
	Total	407	100.0
Unselected Cases		0	.0
Total		407	100.0

a. If weight is in effect, see classification table for the total number of cases.

## Dependent Variable Encoding

Original Value	Internal Value
not shared	0
shared	1

## Block 0: Beginning Block

**Classification Table<sup>a,b</sup>**

			Predicted		Percentage Correct
Observed			Sharted at least once in media not shared	shared	
Step 0	Sharted at least once in media	not shared	0	198	.0
		shared	0	208	100.0
	Overall Percentage				51.2

a. Constant is included in the model.

b. The cut value is .500

**Variables in the Equation**

		B	S.E.	Wald	df	Sig.	Exp(B)
Step 0	Constant	.049	.099	.246	1	.620	1.051

**Variables not in the Equation**

			Score	df	Sig.
Step 0	Variables	Spanish	30.449	1	<.001
	Overall Statistics		30.449	1	<.001

## Block 1: Method = Enter

**Omnibus Tests of Model Coefficients**

		Chi-square	df	Sig.
Step 1	Step	31.794	1	<.001
	Block	31.794	1	<.001
	Model	31.794	1	<.001

**Model Summary**

Step	-2 Log likelihood	Cox & Snell R Square	Nagelkerke R Square
1	530.795 <sup>a</sup>	.075	.100

a. Estimation terminated at iteration number 4 because parameter estimates changed by less than .001.

**Hosmer and Lemeshow Test**

Step	Chi-square	df	Sig.
1	.000	0	.

### Contingency Table for Hosmer and Lemeshow Test

		Sharted at least once in media = not shared		Sharted at least once in media = shared		Total
		Observed	Expected	Observed	Expected	
Step 1	1	176	176.000	137	137.000	313
	2	22	22.000	71	71.000	93

### Classification Table<sup>a</sup>

		Predicted		Percentage Correct
		Sharted at least once in media not shared	shared	
Step 1	Observed			
	Sharted at least once in media	not shared	176	22
		shared	137	71
Overall Percentage				60.8

a. The cut value is .500

### Variables in the Equation

		B	S.E.	Wald	df	Sig.	Exp(B)	95% C.I. for Lower
Step 1 <sup>a</sup>	Spanish	-1.422	.269	27.889	1	<.001	.241	.142
	Constant	1.172	.244	23.056	1	<.001	3.227	

### Variables in the Equation

		95% C.I. for .. Upper
Step 1 <sup>a</sup>	Spanish	.409
	Constant	

a. Variable(s) entered on step 1: Spanish.

```
LOGISTIC REGRESSION VARIABLESshared_in_media
/METHOD=ENTER Portuguese
/PRINT=GOODFIT CI(95)
/CRITERIA=PIN(0.05) POUT(0.10) ITERATE(20) CUT(0.5).
```

### Logistic Regression

## Notes

Output Created		10-SEP-2024 21:15:15
Comments		
Input	Data	C:\Users\apalacios\Documents\Media manuscript\Database.sav
	Active Dataset	DataSet1
	Filter	<none>
	Weight	<none>
	Split File	<none>
	N of Rows in Working Data File	407
Missing Value Handling	Definition of Missing	User-defined missing values are treated as missing
Syntax		LOGISTIC REGRESSION VARIABLES shared_in_media /METHOD=ENTER Portuguese /PRINT=GOODFIT CI(95) /CRITERIA=PIN(0.05) POUT(0.10) ITERATE(20) CUT(0.5).
Resources	Processor Time	00:00:00.02
	Elapsed Time	00:00:00.02

## Case Processing Summary

Unweighted Cases <sup>a</sup>		N	Percent
Selected Cases	Included in Analysis	406	99.8
	Missing Cases	1	.2
	Total	407	100.0
Unselected Cases		0	.0
Total		407	100.0

a. If weight is in effect, see classification table for the total number of cases.

## Dependent Variable Encoding

Original Value	Internal Value
not shared	0
shared	1

## Block 0: Beginning Block

**Classification Table<sup>a,b</sup>**

Observed			Predicted		Percentage Correct
			Sharted at least once in media not shared	shared	
Step 0	Sharted at least once in media	not shared	0	198	.0
		shared	0	208	100.0
	Overall Percentage				51.2

a. Constant is included in the model.

b. The cut value is .500

**Variables in the Equation**

		B	S.E.	Wald	df	Sig.	Exp(B)
Step 0	Constant	.049	.099	.246	1	.620	1.051

**Variables not in the Equation**

			Score	df	Sig.
Step 0	Variables	Portuguese	30.449	1	<.001
	Overall Statistics		30.449	1	<.001

## Block 1: Method = Enter

**Omnibus Tests of Model Coefficients**

		Chi-square	df	Sig.
Step 1	Step	31.794	1	<.001
	Block	31.794	1	<.001
	Model	31.794	1	<.001

**Model Summary**

Step	-2 Log likelihood	Cox & Snell R Square	Nagelkerke R Square
1	530.795 <sup>a</sup>	.075	.100

a. Estimation terminated at iteration number 4 because parameter estimates changed by less than .001.

**Hosmer and Lemeshow Test**

Step	Chi-square	df	Sig.
1	.000	0	.

### Contingency Table for Hosmer and Lemeshow Test

		Sharted at least once in media = not shared		Sharted at least once in media = shared		Total
		Observed	Expected	Observed	Expected	
Step 1	1	176	176.000	137	137.000	313
	2	22	22.000	71	71.000	93

### Classification Table<sup>a</sup>

			Predicted		Percentage Correct
			Sharted at least once in media not shared	shared	
Step 1	Observed				
	Sharted at least once in media	not shared	176	22	88.9
		shared	137	71	34.1
Overall Percentage					60.8

a. The cut value is .500

### Variables in the Equation

		B	S.E.	Wald	df	Sig.	Exp(B)	95% C.I. for Lower
Step 1 <sup>a</sup>	Portuguese	1.422	.269	27.889	1	<.001	4.146	2.446
	Constant	-.251	.114	4.834	1	.028	.778	

### Variables in the Equation

		95% C.I. for .. Upper
Step 1 <sup>a</sup>	Portuguese	7.028
	Constant	

a. Variable(s) entered on step 1: Portuguese.

```
LOGISTIC REGRESSION VARIABLESshared_in_media
/METHOD=ENTER legacy
/PRINT=GOODFIT CI(95)
/CRITERIA=PIN(0.05) POUT(0.10) ITERATE(20) CUT(0.5).
```

### Logistic Regression



## Notes

Output Created		10-SEP-2024 21:19:20
Comments		
Input	Data	C:\Users\apalacios\Documents\Media manuscript\Database.sav
	Active Dataset	DataSet1
	Filter	<none>
	Weight	<none>
	Split File	<none>
	N of Rows in Working Data File	407
Missing Value Handling	Definition of Missing	User-defined missing values are treated as missing
Syntax		LOGISTIC REGRESSION VARIABLES shared_in_media /METHOD=ENTER legacy /PRINT=GOODFIT CI(95) /CRITERIA=PIN(0.05) POUT(0.10) ITERATE(20) CUT(0.5).
Resources	Processor Time	00:00:00.00
	Elapsed Time	00:00:00.00

## Case Processing Summary

Unweighted Cases <sup>a</sup>		N	Percent
Selected Cases	Included in Analysis	406	99.8
	Missing Cases	1	.2
	Total	407	100.0
Unselected Cases		0	.0
Total		407	100.0

a. If weight is in effect, see classification table for the total number of cases.

## Dependent Variable Encoding

Original Value	Internal Value
not shared	0
shared	1

## Block 0: Beginning Block

**Classification Table<sup>a,b</sup>**

			Predicted		Percentage Correct
Observed			Sharted at least once in media not shared	shared	
Step 0	Sharted at least once in media	not shared	0	198	.0
		shared	0	208	100.0
	Overall Percentage				51.2

a. Constant is included in the model.

b. The cut value is .500

**Variables in the Equation**

		B	S.E.	Wald	df	Sig.	Exp(B)
Step 0	Constant	.049	.099	.246	1	.620	1.051

**Variables not in the Equation**

			Score	df	Sig.
Step 0	Variables	legacy	.466	1	.495
	Overall Statistics		.466	1	.495

## Block 1: Method = Enter

**Omnibus Tests of Model Coefficients**

		Chi-square	df	Sig.
Step 1	Step	.467	1	.495
	Block	.467	1	.495
	Model	.467	1	.495

**Model Summary**

Step	-2 Log likelihood	Cox & Snell R Square	Nagelkerke R Square
1	562.123 <sup>a</sup>	.001	.002

a. Estimation terminated at iteration number 2 because parameter estimates changed by less than .001.

**Hosmer and Lemeshow Test**

Step	Chi-square	df	Sig.
1	.000	0	.

### Contingency Table for Hosmer and Lemeshow Test

		Sharted at least once in media = not shared		Sharted at least once in media = shared		Total
		Observed	Expected	Observed	Expected	
Step 1	1	100	100.000	98	98.000	198
	2	98	98.000	110	110.000	208

### Classification Table<sup>a</sup>

			Predicted		Percentage Correct
			Sharted at least once in media not shared	shared	
Step 1	Observed				
	Sharted at least once in media	not shared	100	98	50.5
		shared	98	110	52.9
Overall Percentage					51.7

a. The cut value is .500

### Variables in the Equation

		B	S.E.	Wald	df	Sig.	Exp(B)	95% C.I. for Lower
Step 1 <sup>a</sup>	legacy	-.136	.199	.466	1	.495	.873	.591
	Constant	.116	.139	.692	1	.406	1.122	

### Variables in the Equation

		95% C.I. for .. Upper
Step 1 <sup>a</sup>	legacy	1.289
	Constant	

a. Variable(s) entered on step 1: legacy.

```
LOGISTIC REGRESSION VARIABLESshared_in_media
/METHOD=ENTER digital
/PRINT=GOODFIT CI(95)
/CRITERIA=PIN(0.05) POUT(0.10) ITERATE(20) CUT(0.5).
```

### Logistic Regression

## Notes

Output Created		10-SEP-2024 21:19:20
Comments		
Input	Data	C: \Users\apalacios\Documents\Media manuscript\Database.sav
	Active Dataset	DataSet1
	Filter	<none>
	Weight	<none>
	Split File	<none>
	N of Rows in Working Data File	407
Missing Value Handling	Definition of Missing	User-defined missing values are treated as missing
Syntax		LOGISTIC REGRESSION VARIABLES shared_in_media /METHOD=ENTER digital /PRINT=GOODFIT CI(95) /CRITERIA=PIN(0.05) POUT(0.10) ITERATE(20) CUT(0.5).
Resources	Processor Time	00:00:00.00
	Elapsed Time	00:00:00.00

## Case Processing Summary

Unweighted Cases <sup>a</sup>		N	Percent
Selected Cases	Included in Analysis	406	99.8
	Missing Cases	1	.2
	Total	407	100.0
Unselected Cases		0	.0
Total		407	100.0

a. If weight is in effect, see classification table for the total number of cases.

## Dependent Variable Encoding

Original Value	Internal Value
not shared	0
shared	1

## Block 0: Beginning Block

**Classification Table<sup>a,b</sup>**

		Predicted		Percentage Correct
Observed		Sharted at least once in media not shared	shared	
Step 0	Sharted at least once in media	not shared	0	.0
		shared	0	100.0
	Overall Percentage			51.2

a. Constant is included in the model.

b. The cut value is .500

**Variables in the Equation**

		B	S.E.	Wald	df	Sig.	Exp(B)
Step 0	Constant	.049	.099	.246	1	.620	1.051

**Variables not in the Equation**

		Score	df	Sig.
Step 0	Variables digital	.466	1	.495
	Overall Statistics	.466	1	.495

## Block 1: Method = Enter

**Omnibus Tests of Model Coefficients**

		Chi-square	df	Sig.
Step 1	Step	.467	1	.495
	Block	.467	1	.495
	Model	.467	1	.495

**Model Summary**

Step	-2 Log likelihood	Cox & Snell R Square	Nagelkerke R Square
1	562.123 <sup>a</sup>	.001	.002

a. Estimation terminated at iteration number 2 because parameter estimates changed by less than .001.

**Hosmer and Lemeshow Test**

Step	Chi-square	df	Sig.
1	.000	0	.

### Contingency Table for Hosmer and Lemeshow Test

		Sharted at least once in media = not shared		Sharted at least once in media = shared		Total
		Observed	Expected	Observed	Expected	
Step 1	1	100	100.000	98	98.000	198
	2	98	98.000	110	110.000	208

### Classification Table<sup>a</sup>

			Predicted		Percentage Correct
			Sharted at least once in media not shared	shared	
Step 1	Observed				
	Sharted at least once in media	not shared	100	98	50.5
		shared	98	110	52.9
	Overall Percentage				51.7

a. The cut value is .500

### Variables in the Equation

		B	S.E.	Wald	df	Sig.	Exp(B)	95% C.I. for Lower
Step 1 <sup>a</sup>	digital	.136	.199	.466	1	.495	1.145	.776
	Constant	-.020	.142	.020	1	.887	.980	

### Variables in the Equation

		95% C.I. for .. Upper
Step 1 <sup>a</sup>	digital	1.691
	Constant	

a. Variable(s) entered on step 1: digital.

```
LOGISTIC REGRESSION VARIABLESshared_in_media
/METHOD=ENTER newsmmedia
/PRINT=GOODFIT CI(95)
/CRITERIA=PIN(0.05) POUT(0.10) ITERATE(20) CUT(0.5).
```

### Logistic Regression

## Notes

Output Created		10-SEP-2024 21:21:20
Comments		
Input	Data	C:\Users\apalacios\Documents\Media manuscript\Database.sav
	Active Dataset	DataSet1
	Filter	<none>
	Weight	<none>
	Split File	<none>
	N of Rows in Working Data File	407
Missing Value Handling	Definition of Missing	User-defined missing values are treated as missing
Syntax		LOGISTIC REGRESSION VARIABLES shared_in_media /METHOD=ENTER newsmedia /PRINT=GOODFIT CI(95) /CRITERIA=PIN(0.05) POUT(0.10) ITERATE(20) CUT(0.5).
Resources	Processor Time	00:00:00.00
	Elapsed Time	00:00:00.00

## Case Processing Summary

Unweighted Cases <sup>a</sup>		N	Percent
Selected Cases	Included in Analysis	406	99.8
	Missing Cases	1	.2
	Total	407	100.0
Unselected Cases		0	.0
Total		407	100.0

a. If weight is in effect, see classification table for the total number of cases.

## Dependent Variable Encoding

Original Value	Internal Value
not shared	0
shared	1

## Block 0: Beginning Block

**Classification Table<sup>a,b</sup>**

			Predicted		Percentage Correct
Observed			Sharted at least once in media not shared	shared	
Step 0	Sharted at least once in media	not shared	0	198	.0
		shared	0	208	100.0
	Overall Percentage				51.2

a. Constant is included in the model.

b. The cut value is .500

**Variables in the Equation**

		B	S.E.	Wald	df	Sig.	Exp(B)
Step 0	Constant	.049	.099	.246	1	.620	1.051

**Variables not in the Equation**

			Score	df	Sig.
Step 0	Variables	newsmedia	.124	1	.725
	Overall Statistics		.124	1	.725

## Block 1: Method = Enter

**Omnibus Tests of Model Coefficients**

		Chi-square	df	Sig.
Step 1	Step	.124	1	.725
	Block	.124	1	.725
	Model	.124	1	.725

**Model Summary**

Step	-2 Log likelihood	Cox & Snell R Square	Nagelkerke R Square
1	562.466 <sup>a</sup>	.000	.000

a. Estimation terminated at iteration number 2 because parameter estimates changed by less than .001.

**Hosmer and Lemeshow Test**

Step	Chi-square	df	Sig.
1	.000	0	.



### Contingency Table for Hosmer and Lemeshow Test

		Sharted at least once in media = not shared		Sharted at least once in media = shared		Total
		Observed	Expected	Observed	Expected	
Step 1	1	9	9.000	8	8.000	17
	2	189	189.000	200	200.000	389

### Classification Table<sup>a</sup>

			Predicted		Percentage Correct
			Sharted at least once in media not shared	shared	
Observed					
Step 1	Sharted at least once in media	not shared	9	189	4.5
		shared	8	200	96.2
	Overall Percentage				51.5

a. The cut value is .500

### Variables in the Equation

		B	S.E.	Wald	df	Sig.	Exp(B)	95% C.I. for Lower
Step 1 <sup>a</sup>	newsmedia	.174	.496	.123	1	.725	1.190	.450
	Constant	-.118	.486	.059	1	.808	.889	

### Variables in the Equation

		95% C.I. for .. Upper
Step 1 <sup>a</sup>	newsmedia	3.150
	Constant	

a. Variable(s) entered on step 1: newsmedia.

```
LOGISTIC REGRESSION VARIABLESshared_in_media
/METHOD=ENTER Government
/PRINT=GOODFIT CI(95)
/CRITERIA=PIN(0.05) POUT(0.10) ITERATE(20) CUT(0.5).
```

### Logistic Regression

## Notes

Output Created		10-SEP-2024 21:21:20
Comments		
Input	Data	C:\Users\apalacios\Documents\Media manuscript\Database.sav
	Active Dataset	DataSet1
	Filter	<none>
	Weight	<none>
	Split File	<none>
	N of Rows in Working Data File	407
Missing Value Handling	Definition of Missing	User-defined missing values are treated as missing
Syntax		LOGISTIC REGRESSION VARIABLES shared_in_media /METHOD=ENTER Government /PRINT=GOODFIT CI(95) /CRITERIA=PIN(0.05) POUT(0.10) ITERATE(20) CUT(0.5).
Resources	Processor Time	00:00:00.00
	Elapsed Time	00:00:00.00

## Case Processing Summary

Unweighted Cases <sup>a</sup>		N	Percent
Selected Cases	Included in Analysis	406	99.8
	Missing Cases	1	.2
	Total	407	100.0
Unselected Cases		0	.0
Total		407	100.0

a. If weight is in effect, see classification table for the total number of cases.

## Dependent Variable Encoding

Original Value	Internal Value
not shared	0
shared	1

## Block 0: Beginning Block

**Classification Table<sup>a,b</sup>**

			Predicted		Percentage Correct
Observed			Sharted at least once in media not shared	shared	
Step 0	Sharted at least once in media	not shared	0	198	.0
		shared	0	208	100.0
	Overall Percentage				51.2

a. Constant is included in the model.

b. The cut value is .500

**Variables in the Equation**

		B	S.E.	Wald	df	Sig.	Exp(B)
Step 0	Constant	.049	.099	.246	1	.620	1.051

**Variables not in the Equation**

			Score	df	Sig.
Step 0	Variables	Government	.316	1	.574
	Overall Statistics		.316	1	.574

## Block 1: Method = Enter

**Omnibus Tests of Model Coefficients**

		Chi-square	df	Sig.
Step 1	Step	.318	1	.573
	Block	.318	1	.573
	Model	.318	1	.573

**Model Summary**

Step	-2 Log likelihood	Cox & Snell R Square	Nagelkerke R Square
1	562.271 <sup>a</sup>	.001	.001

a. Estimation terminated at iteration number 3 because parameter estimates changed by less than .001.

**Hosmer and Lemeshow Test**

Step	Chi-square	df	Sig.
1	.000	0	.

### Contingency Table for Hosmer and Lemeshow Test

		Sharted at least once in media = not shared		Sharted at least once in media = shared		Total
		Observed	Expected	Observed	Expected	
Step 1	1	198	198.000	208	208.000	406

### Classification Table<sup>a</sup>

			Predicted		Percentage Correct
			Sharted at least once in media not shared	shared	
Step 1	Sharted at least once in media	not shared	0	198	.0
		shared	0	208	100.0
	Overall Percentage				51.2

a. The cut value is .500

### Variables in the Equation

		B	S.E.	Wald	df	Sig.	Exp(B)	95% C.I. for Lower
Step 1 <sup>a</sup>	Government	.365	.653	.312	1	.576	1.441	.400
	Constant	.040	.101	.162	1	.688	1.041	

### Variables in the Equation

		95% C.I. for .. Upper
Step 1 <sup>a</sup>	Government	5.183
	Constant	

a. Variable(s) entered on step 1: Government.

```
LOGISTIC REGRESSION VARIABLESshared_in_media
/METHOD=ENTER Op_Ed
/PRINT=GOODFITCI(95)
/CRITERIA=PIN(0.05) POUT(0.10) ITERATE(20) CUT(0.5).
```

### Logistic Regression

## Notes

Output Created		10-SEP-2024 21:24:01
Comments		
Input	Data	C:\Users\apalacios\Documents\Media manuscript\Database.sav
	Active Dataset	DataSet1
	Filter	<none>
	Weight	<none>
	Split File	<none>
	N of Rows in Working Data File	407
Missing Value Handling	Definition of Missing	User-defined missing values are treated as missing
Syntax		LOGISTIC REGRESSION VARIABLES shared_in_media /METHOD=ENTER Op_Ed /PRINT=GOODFIT CI(95) /CRITERIA=PIN(0.05) POUT(0.10) ITERATE(20) CUT(0.5).
Resources	Processor Time	00:00:00.00
	Elapsed Time	00:00:00.00

## Case Processing Summary

Unweighted Cases <sup>a</sup>		N	Percent
Selected Cases	Included in Analysis	406	99.8
	Missing Cases	1	.2
	Total	407	100.0
Unselected Cases		0	.0
Total		407	100.0

a. If weight is in effect, see classification table for the total number of cases.

## Dependent Variable Encoding

Original Value	Internal Value
not shared	0
shared	1

## Block 0: Beginning Block

**Classification Table<sup>a,b</sup>**

			Predicted		Percentage Correct
Observed			Sharted at least once in media not shared	shared	
Step 0	Sharted at least once in media	not shared	0	198	.0
		shared	0	208	100.0
	Overall Percentage				51.2

a. Constant is included in the model.

b. The cut value is .500

**Variables in the Equation**

		B	S.E.	Wald	df	Sig.	Exp(B)
Step 0	Constant	.049	.099	.246	1	.620	1.051

**Variables not in the Equation**

			Score	df	Sig.
Step 0	Variables	Op_Ed	1.464	1	.226
	Overall Statistics		1.464	1	.226

## Block 1: Method = Enter

**Omnibus Tests of Model Coefficients**

		Chi-square	df	Sig.
Step 1	Step	1.506	1	.220
	Block	1.506	1	.220
	Model	1.506	1	.220

**Model Summary**

Step	-2 Log likelihood	Cox & Snell R Square	Nagelkerke R Square
1	561.084 <sup>a</sup>	.004	.005

a. Estimation terminated at iteration number 3 because parameter estimates changed by less than .001.

**Hosmer and Lemeshow Test**

Step	Chi-square	df	Sig.
1	.000	0	.

### Contingency Table for Hosmer and Lemeshow Test

		Sharted at least once in media = not shared		Sharted at least once in media = shared		Total
		Observed	Expected	Observed	Expected	
Step 1	1	5	5.000	2	2.000	7
	2	193	193.000	206	206.000	399

### Classification Table<sup>a</sup>

			Predicted		Percentage Correct
			Sharted at least once in media not shared	shared	
Step 1	Observed				
	Sharted at least once in media	not shared	5	193	2.5
		shared	2	206	99.0
	Overall Percentage				52.0

a. The cut value is .500

### Variables in the Equation

		B	S.E.	Wald	df	Sig.	Exp(B)	95% C.I. for Lower
Step 1 <sup>a</sup>	Op_Ed	-.981	.843	1.357	1	.244	.375	.072
	Constant	.065	.100	.423	1	.515	1.067	

### Variables in the Equation

		95% C.I. for .. Upper
Step 1 <sup>a</sup>	Op_Ed	1.954
	Constant	

a. Variable(s) entered on step 1: Op\_Ed.

```
LOGISTIC REGRESSION VARIABLESshared_in_media
/METHOD=ENTER New_treatment
/PRINT=GOODFIT CI(95)
/CRITERIA=PIN(0.05) POUT(0.10) ITERATE(20) CUT(0.5).
```

### Logistic Regression

## Notes

Output Created		10-SEP-2024 21:25:42
Comments		
Input	Data	C:\Users\apalacios\Documents\Media manuscript\Database.sav
	Active Dataset	DataSet1
	Filter	<none>
	Weight	<none>
	Split File	<none>
	N of Rows in Working Data File	407
Missing Value Handling	Definition of Missing	User-defined missing values are treated as missing
Syntax		LOGISTIC REGRESSION VARIABLES shared_in_media /METHOD=ENTER New_treatment /PRINT=GOODFIT CI(95) /CRITERIA=PIN(0.05) POUT(0.10) ITERATE(20) CUT(0.5).
Resources	Processor Time	00:00:00.00
	Elapsed Time	00:00:00.00

## Case Processing Summary

Unweighted Cases <sup>a</sup>		N	Percent
Selected Cases	Included in Analysis	406	99.8
	Missing Cases	1	.2
	Total	407	100.0
Unselected Cases		0	.0
Total		407	100.0

a. If weight is in effect, see classification table for the total number of cases.

## Dependent Variable Encoding

Original Value	Internal Value
not shared	0
shared	1



## Block 0: Beginning Block

**Classification Table<sup>a,b</sup>**

			Predicted		Percentage Correct
Observed			Sharted at least once in media not shared	shared	
Step 0	Sharted at least once in media	not shared	0	198	.0
		shared	0	208	100.0
	Overall Percentage				51.2

a. Constant is included in the model.

b. The cut value is .500

**Variables in the Equation**

		B	S.E.	Wald	df	Sig.	Exp(B)
Step 0	Constant	.049	.099	.246	1	.620	1.051

**Variables not in the Equation**

			Score	df	Sig.
Step 0	Variables	MI- New Treatment	.343	1	.558
	Overall Statistics		.343	1	.558

## Block 1: Method = Enter

**Omnibus Tests of Model Coefficients**

		Chi-square	df	Sig.
Step 1	Step	.343	1	.558
	Block	.343	1	.558
	Model	.343	1	.558

**Model Summary**

Step	-2 Log likelihood	Cox & Snell R Square	Nagelkerke R Square
1	562.246 <sup>a</sup>	.001	.001

a. Estimation terminated at iteration number 2 because parameter estimates changed by less than .001.

**Hosmer and Lemeshow Test**

Step	Chi-square	df	Sig.
1	.000	0	.

### Contingency Table for Hosmer and Lemeshow Test

		Sharted at least once in media = not shared		Sharted at least once in media = shared		Total
		Observed	Expected	Observed	Expected	
Step 1	1	76	76.000	74	74.000	150
	2	122	122.000	134	134.000	256

### Classification Table<sup>a</sup>

			Predicted		Percentage Correct
			Sharted at least once in media not shared	shared	
Step 1	Observed				
	Sharted at least once in media	not shared	76	122	38.4
		shared	74	134	64.4
Overall Percentage					51.7

a. The cut value is .500

### Variables in the Equation

		B	S.E.	Wald	df	Sig.	Exp(B)
Step 1 <sup>a</sup>	MI- New Treatment	-.120	.206	.343	1	.558	.886
	Constant	.094	.125	.562	1	.453	1.098

### Variables in the Equation

		95% C.I. for EXP(B)	
		Lower	Upper
Step 1 <sup>a</sup>	MI- New Treatment	.592	1.327
	Constant		

a. Variable(s) entered on step 1: MI- New Treatment.

```
LOGISTIC REGRESSION VARIABLESshared_in_media
/METHOD=ENTER Drug_abuse
/PRINT=GOODFIT CI(95)
/CRITERIA=PIN(0.05) POUT(0.10) ITERATE(20) CUT(0.5).
```

### Logistic Regression

## Notes

Output Created		10-SEP-2024 21:25:42
Comments		
Input	Data	C:\Users\apalacios\Documents\Media manuscript\Database.sav
	Active Dataset	DataSet1
	Filter	<none>
	Weight	<none>
	Split File	<none>
	N of Rows in Working Data File	407
Missing Value Handling	Definition of Missing	User-defined missing values are treated as missing
Syntax		LOGISTIC REGRESSION VARIABLES shared_in_media /METHOD=ENTER Drug_abuse /PRINT=GOODFIT CI(95) /CRITERIA=PIN(0.05) POUT(0.10) ITERATE(20) CUT(0.5).
Resources	Processor Time	00:00:00.00
	Elapsed Time	00:00:00.00

## Case Processing Summary

Unweighted Cases <sup>a</sup>		N	Percent
Selected Cases	Included in Analysis	406	99.8
	Missing Cases	1	.2
	Total	407	100.0
Unselected Cases		0	.0
Total		407	100.0

a. If weight is in effect, see classification table for the total number of cases.

## Dependent Variable Encoding

Original Value	Internal Value
not shared	0
shared	1

## Block 0: Beginning Block

**Classification Table<sup>a,b</sup>**

			Predicted		Percentage Correct
Observed			Sharted at least once in media not shared	shared	
Step 0	Sharted at least once in media	not shared	0	198	.0
		shared	0	208	100.0
	Overall Percentage				51.2

a. Constant is included in the model.

b. The cut value is .500

**Variables in the Equation**

		B	S.E.	Wald	df	Sig.	Exp(B)
Step 0	Constant	.049	.099	.246	1	.620	1.051

**Variables not in the Equation**

			Score	df	Sig.
Step 0	Variables	MI-Drug Abuse	.078	1	.779
	Overall Statistics		.078	1	.779

## Block 1: Method = Enter

**Omnibus Tests of Model Coefficients**

		Chi-square	df	Sig.
Step 1	Step	.078	1	.779
	Block	.078	1	.779
	Model	.078	1	.779

**Model Summary**

Step	-2 Log likelihood	Cox & Snell R Square	Nagelkerke R Square
1	562.511 <sup>a</sup>	.000	.000

a. Estimation terminated at iteration number 2 because parameter estimates changed by less than .001.

**Hosmer and Lemeshow Test**

Step	Chi-square	df	Sig.
1	.000	0	.

### Contingency Table for Hosmer and Lemeshow Test

		Sharted at least once in media = not shared		Sharted at least once in media = shared		Total
		Observed	Expected	Observed	Expected	
Step 1	1	49	49.000	49	49.000	98
	2	149	149.000	159	159.000	308

### Classification Table<sup>a</sup>

			Predicted		Percentage Correct
			Sharted at least once in media not shared	shared	
Step 1	Observed				
	Sharted at least once in media	not shared	0	198	.0
		shared	0	208	100.0
	Overall Percentage				51.2

a. The cut value is .500

### Variables in the Equation

		B	S.E.	Wald	df	Sig.	Exp(B)
Step 1 <sup>a</sup>	MI-Drug Abuse	-.065	.232	.078	1	.779	.937
	Constant	.065	.114	.325	1	.569	1.067

### Variables in the Equation

		95% C.I. for EXP(B)	
		Lower	Upper
Step 1 <sup>a</sup>	MI-Drug Abuse	.595	1.477
	Constant		

a. Variable(s) entered on step 1: MI-Drug Abuse.

```
LOGISTIC REGRESSION VARIABLESshared_in_media
/METHOD=ENTER Regulation
/PRINT=GOODFIT CI(95)
/CRITERIA=PIN(0.05) POUT(0.10) ITERATE(20) CUT(0.5).
```

### Logistic Regression

## Notes

Output Created		10-SEP-2024 21:25:42
Comments		
Input	Data	C:\Users\apalacios\Documents\Media manuscript\Database.sav
	Active Dataset	DataSet1
	Filter	<none>
	Weight	<none>
	Split File	<none>
	N of Rows in Working Data File	407
Missing Value Handling	Definition of Missing	User-defined missing values are treated as missing
Syntax		LOGISTIC REGRESSION VARIABLES shared_in_media /METHOD=ENTER Regulation /PRINT=GOODFIT CI(95) /CRITERIA=PIN(0.05) POUT(0.10) ITERATE(20) CUT(0.5).
Resources	Processor Time	00:00:00.00
	Elapsed Time	00:00:00.00

## Case Processing Summary

Unweighted Cases <sup>a</sup>		N	Percent
Selected Cases	Included in Analysis	406	99.8
	Missing Cases	1	.2
	Total	407	100.0
Unselected Cases		0	.0
Total		407	100.0

a. If weight is in effect, see classification table for the total number of cases.

## Dependent Variable Encoding

Original Value	Internal Value
not shared	0
shared	1

## Block 0: Beginning Block

**Classification Table<sup>a,b</sup>**

			Predicted		Percentage Correct
Observed			Sharted at least once in media not shared	shared	
Step 0	Sharted at least once in media	not shared	0	198	.0
		shared	0	208	100.0
	Overall Percentage				51.2

a. Constant is included in the model.

b. The cut value is .500

**Variables in the Equation**

		B	S.E.	Wald	df	Sig.	Exp(B)
Step 0	Constant	.049	.099	.246	1	.620	1.051

**Variables not in the Equation**

			Score	df	Sig.
Step 0	Variables	MI - Regulation	4.274	1	.039
	Overall Statistics		4.274	1	.039

## Block 1: Method = Enter

**Omnibus Tests of Model Coefficients**

		Chi-square	df	Sig.
Step 1	Step	4.329	1	.037
	Block	4.329	1	.037
	Model	4.329	1	.037

**Model Summary**

Step	-2 Log likelihood	Cox & Snell R Square	Nagelkerke R Square
1	558.261 <sup>a</sup>	.011	.014

a. Estimation terminated at iteration number 3 because parameter estimates changed by less than .001.

**Hosmer and Lemeshow Test**

Step	Chi-square	df	Sig.
1	.000	0	.

### Contingency Table for Hosmer and Lemeshow Test

		Sharted at least once in media = not shared		Sharted at least once in media = shared		Total
		Observed	Expected	Observed	Expected	
Step 1	1	177	177.000	171	171.000	348
	2	21	21.000	37	37.000	58

### Classification Table<sup>a</sup>

			Predicted		Percentage Correct
			Sharted at least once in media not shared	shared	
Step 1	Observed				
	Sharted at least once in media	not shared	177	21	89.4
		shared	171	37	17.8
Overall Percentage					52.7

a. The cut value is .500

### Variables in the Equation

		B	S.E.	Wald	df	Sig.	Exp(B)
Step 1 <sup>a</sup>	MI - Regulation	.601	.294	4.191	1	.041	1.824
	Constant	-.034	.107	.103	1	.748	.966

### Variables in the Equation

		95% C.I. for EXP(B)	
		Lower	Upper
Step 1 <sup>a</sup>	MI - Regulation	1.026	3.242
	Constant		

a. Variable(s) entered on step 1: MI - Regulation.

```
LOGISTIC REGRESSION VARIABLESshared_in_media
/METHOD=ENTER Health
/PRINT=GOODFIT CI(95)
/CRITERIA=PIN(0.05) POUT(0.10) ITERATE(20) CUT(0.5).
```

### Logistic Regression



## Notes

Output Created		10-SEP-2024 21:25:42
Comments		
Input	Data	C: \Users\apalacios\Documents\Media manuscript\Database.sav
	Active Dataset	DataSet1
	Filter	<none>
	Weight	<none>
	Split File	<none>
	N of Rows in Working Data File	407
Missing Value Handling	Definition of Missing	User-defined missing values are treated as missing
Syntax		LOGISTIC REGRESSION VARIABLES shared_in_media /METHOD=ENTER Health /PRINT=GOODFIT CI(95) /CRITERIA=PIN(0.05) POUT(0.10) ITERATE(20) CUT(0.5).
Resources	Processor Time	00:00:00.00
	Elapsed Time	00:00:00.00

## Case Processing Summary

Unweighted Cases <sup>a</sup>		N	Percent
Selected Cases	Included in Analysis	406	99.8
	Missing Cases	1	.2
	Total	407	100.0
Unselected Cases		0	.0
Total		407	100.0

a. If weight is in effect, see classification table for the total number of cases.

## Dependent Variable Encoding

Original Value	Internal Value
not shared	0
shared	1

## Block 0: Beginning Block

**Classification Table<sup>a,b</sup>**

			Predicted		Percentage Correct
Observed			Sharted at least once in media not shared	shared	
Step 0	Sharted at least once in media	not shared	0	198	.0
		shared	0	208	100.0
	Overall Percentage				51.2

a. Constant is included in the model.

b. The cut value is .500

**Variables in the Equation**

		B	S.E.	Wald	df	Sig.	Exp(B)
Step 0	Constant	.049	.099	.246	1	.620	1.051

**Variables not in the Equation**

			Score	df	Sig.
Step 0	Variables	MI- Health	.237	1	.627
	Overall Statistics		.237	1	.627

## Block 1: Method = Enter

**Omnibus Tests of Model Coefficients**

		Chi-square	df	Sig.
Step 1	Step	.237	1	.627
	Block	.237	1	.627
	Model	.237	1	.627

**Model Summary**

Step	-2 Log likelihood	Cox & Snell R Square	Nagelkerke R Square
1	562.353 <sup>a</sup>	.001	.001

a. Estimation terminated at iteration number 2 because parameter estimates changed by less than .001.

**Hosmer and Lemeshow Test**

Step	Chi-square	df	Sig.
1	.000	0	.

### Contingency Table for Hosmer and Lemeshow Test

		Sharted at least once in media = not shared		Sharted at least once in media = shared		Total
		Observed	Expected	Observed	Expected	
Step 1	1	30	30.000	28	28.000	58
	2	168	168.000	180	180.000	348

### Classification Table<sup>a</sup>

			Predicted		Percentage Correct
Observed			Sharted at least once in media not shared	shared	
Step 1	Sharted at least once in media	not shared	30	168	15.2
		shared	28	180	86.5
	Overall Percentage				

a. The cut value is .500

### Variables in the Equation

		B	S.E.	Wald	df	Sig.	Exp(B)	95% C.I. for Lower
Step 1 <sup>a</sup>	MI- Health	-.138	.284	.236	1	.627	.871	.499
	Constant	.069	.107	.414	1	.520	1.071	

### Variables in the Equation

		95% C.I. for .. Upper
Step 1 <sup>a</sup>	MI- Health	1.519
	Constant	

a. Variable(s) entered on step 1: MI- Health.

```
LOGISTIC REGRESSION VARIABLESshared_in_media
/METHOD=ENTER Price
/PRINT=GOODFIT CI(95)
/CRITERIA=PIN(0.05) POUT(0.10) ITERATE(20) CUT(0.5).
```

### Logistic Regression

## Notes

Output Created		10-SEP-2024 21:25:42
Comments		
Input	Data	C: \Users\apalacios\Documents\Media manuscript\Database.sav
	Active Dataset	DataSet1
	Filter	<none>
	Weight	<none>
	Split File	<none>
	N of Rows in Working Data File	407
Missing Value Handling	Definition of Missing	User-defined missing values are treated as missing
Syntax		LOGISTIC REGRESSION VARIABLES shared_in_media /METHOD=ENTER Price /PRINT=GOODFIT CI(95) /CRITERIA=PIN(0.05) POUT(0.10) ITERATE(20) CUT(0.5).
Resources	Processor Time	00:00:00.00
	Elapsed Time	00:00:00.00

## Case Processing Summary

Unweighted Cases <sup>a</sup>		N	Percent
Selected Cases	Included in Analysis	406	99.8
	Missing Cases	1	.2
	Total	407	100.0
Unselected Cases		0	.0
Total		407	100.0

a. If weight is in effect, see classification table for the total number of cases.

## Dependent Variable Encoding

Original Value	Internal Value
not shared	0
shared	1

## Block 0: Beginning Block

**Classification Table<sup>a,b</sup>**

			Predicted		Percentage Correct
Observed			Sharted at least once in media not shared	shared	
Step 0	Sharted at least once in media	not shared	0	198	.0
		shared	0	208	100.0
	Overall Percentage				51.2

a. Constant is included in the model.

b. The cut value is .500

**Variables in the Equation**

		B	S.E.	Wald	df	Sig.	Exp(B)
Step 0	Constant	.049	.099	.246	1	.620	1.051

**Variables not in the Equation**

			Score	df	Sig.
Step 0	Variables	MI - Price	1.636	1	.201
	Overall Statistics		1.636	1	.201

## Block 1: Method = Enter

**Omnibus Tests of Model Coefficients**

		Chi-square	df	Sig.
Step 1	Step	1.643	1	.200
	Block	1.643	1	.200
	Model	1.643	1	.200

**Model Summary**

Step	-2 Log likelihood	Cox & Snell R Square	Nagelkerke R Square
1	560.946 <sup>a</sup>	.004	.005

a. Estimation terminated at iteration number 3 because parameter estimates changed by less than .001.

**Hosmer and Lemeshow Test**

Step	Chi-square	df	Sig.
1	.000	0	.

### Contingency Table for Hosmer and Lemeshow Test

		Sharted at least once in media = not shared		Sharted at least once in media = shared		Total
		Observed	Expected	Observed	Expected	
Step 1	1	18	18.000	12	12.000	30
	2	180	180.000	196	196.000	376

### Classification Table<sup>a</sup>

			Predicted		Percentage Correct
			Sharted at least once in media not shared	shared	
Step 1	Observed				
	Sharted at least once in media	not shared	18	180	9.1
		shared	12	196	94.2
Overall Percentage					52.7

a. The cut value is .500

### Variables in the Equation

		B	S.E.	Wald	df	Sig.	Exp(B)	95% C.I. for Lower
Step 1 <sup>a</sup>	MI - Price	-.491	.387	1.610	1	.205	.612	.287
	Constant	.085	.103	.680	1	.409	1.089	

### Variables in the Equation

		95% C.I. for .. Upper
Step 1 <sup>a</sup>	MI - Price	1.306
	Constant	

a. Variable(s) entered on step 1: MI - Price.

```
LOGISTIC REGRESSION VARIABLESshared_in_media
/METHOD=ENTER Business_news
/PRINT=GOODFIT CI(95)
/CRITERIA=PIN(0.05) POUT(0.10) ITERATE(20) CUT(0.5).
```

### Logistic Regression

## Notes

Output Created		10-SEP-2024 21:31:17
Comments		
Input	Data	C:\Users\apalacios\Documents\Media manuscript\Database.sav
	Active Dataset	DataSet1
	Filter	<none>
	Weight	<none>
	Split File	<none>
	N of Rows in Working Data File	407
Missing Value Handling	Definition of Missing	User-defined missing values are treated as missing
Syntax		LOGISTIC REGRESSION VARIABLES shared_in_media /METHOD=ENTER Business_news /PRINT=GOODFIT CI(95) /CRITERIA=PIN(0.05) POUT(0.10) ITERATE(20) CUT(0.5).
Resources	Processor Time	00:00:00.00
	Elapsed Time	00:00:00.00

## Case Processing Summary

Unweighted Cases <sup>a</sup>		N	Percent
Selected Cases	Included in Analysis	406	99.8
	Missing Cases	1	.2
	Total	407	100.0
Unselected Cases		0	.0
Total		407	100.0

a. If weight is in effect, see classification table for the total number of cases.

## Dependent Variable Encoding

Original Value	Internal Value
not shared	0
shared	1

## Block 0: Beginning Block

**Classification Table<sup>a,b</sup>**

			Predicted		Percentage Correct
Observed			Sharted at least once in media not shared	shared	
Step 0	Sharted at least once in media	not shared	0	198	.0
		shared	0	208	100.0
	Overall Percentage				51.2

a. Constant is included in the model.

b. The cut value is .500

**Variables in the Equation**

		B	S.E.	Wald	df	Sig.	Exp(B)
Step 0	Constant	.049	.099	.246	1	.620	1.051

**Variables not in the Equation**

			Score	df	Sig.
Step 0	Variables	Mi - Business News	1.179	1	.278
	Overall Statistics		1.179	1	.278

## Block 1: Method = Enter

**Omnibus Tests of Model Coefficients**

		Chi-square	df	Sig.
Step 1	Step	1.204	1	.272
	Block	1.204	1	.272
	Model	1.204	1	.272

**Model Summary**

Step	-2 Log likelihood	Cox & Snell R Square	Nagelkerke R Square
1	561.385 <sup>a</sup>	.003	.004

a. Estimation terminated at iteration number 3 because parameter estimates changed by less than .001.

**Hosmer and Lemeshow Test**

Step	Chi-square	df	Sig.
1	.000	0	.



### Contingency Table for Hosmer and Lemeshow Test

		Sharted at least once in media = not shared		Sharted at least once in media = shared		Total
		Observed	Expected	Observed	Expected	
Step 1	1	198	198.000	208	208.000	406

### Classification Table<sup>a</sup>

		Predicted		Percentage Correct
		Sharted at least once in media not shared	Sharted at least once in media shared	
Step 1	Observed			
	Sharted at least once in media	not shared	0	198
		shared	0	208
Overall Percentage				51.2

a. The cut value is .500

### Variables in the Equation

		B	S.E.	Wald	df	Sig.	Exp(B)
Step 1 <sup>a</sup>	Mi - Business News	.663	.621	1.140	1	.286	1.940
	Constant	.030	.101	.091	1	.762	1.031

### Variables in the Equation

		95% C.I. for EXP(B)	
		Lower	Upper
Step 1 <sup>a</sup>	Mi - Business News	.575	6.547
	Constant		

a. Variable(s) entered on step 1: Mi - Business News.

```
LOGISTIC REGRESSION VARIABLESshared_in_media
/METHOD=ENTER Propaganda
/PRINT=GOODFITCI(95)
/CRITERIA=PIN(0.05) POUT(0.10) ITERATE(20) CUT(0.5).
```

### Logistic Regression

## Notes

Output Created		10-SEP-2024 21:33:55
Comments		
Input	Data	C:\Users\apalacios\Documents\Media manuscript\Database.sav
	Active Dataset	DataSet1
	Filter	<none>
	Weight	<none>
	Split File	<none>
	N of Rows in Working Data File	407
Missing Value Handling	Definition of Missing	User-defined missing values are treated as missing
Syntax		LOGISTIC REGRESSION VARIABLES shared_in_media /METHOD=ENTER Propaganda /PRINT=GOODFIT CI(95) /CRITERIA=PIN(0.05) POUT(0.10) ITERATE(20) CUT(0.5).
Resources	Processor Time	00:00:00.00
	Elapsed Time	00:00:00.00

## Case Processing Summary

Unweighted Cases <sup>a</sup>		N	Percent
Selected Cases	Included in Analysis	406	99.8
	Missing Cases	1	.2
	Total	407	100.0
Unselected Cases		0	.0
Total		407	100.0

a. If weight is in effect, see classification table for the total number of cases.

## Dependent Variable Encoding

Original Value	Internal Value
not shared	0
shared	1

## Block 0: Beginning Block

**Classification Table<sup>a,b</sup>**

Observed			Predicted		Percentage Correct
			Sharted at least once in media not shared	shared	
Step 0	Sharted at least once in media	not shared	0	198	.0
		shared	0	208	100.0
	Overall Percentage				51.2

a. Constant is included in the model.

b. The cut value is .500

**Variables in the Equation**

		B	S.E.	Wald	df	Sig.	Exp(B)
Step 0	Constant	.049	.099	.246	1	.620	1.051

**Variables not in the Equation**

				Score	df	Sig.
Step 0	Variables	propaganda yes or maybe		.276	1	.599
	Overall Statistics			.276	1	.599

## Block 1: Method = Enter

**Omnibus Tests of Model Coefficients**

		Chi-square	df	Sig.
Step 1	Step	.276	1	.599
	Block	.276	1	.599
	Model	.276	1	.599

**Model Summary**

Step	-2 Log likelihood	Cox & Snell R Square	Nagelkerke R Square
1	562.313 <sup>a</sup>	.001	.001

a. Estimation terminated at iteration number 2 because parameter estimates changed by less than .001.

**Hosmer and Lemeshow Test**

Step	Chi-square	df	Sig.
1	.000	0	.

### Contingency Table for Hosmer and Lemeshow Test

		Sharted at least once in media = not shared		Sharted at least once in media = shared		Total
		Observed	Expected	Observed	Expected	
Step 1	1	139	139.000	141	141.000	280
	2	59	59.000	67	67.000	126

### Classification Table<sup>a</sup>

		Predicted		Percentage Correct
		Sharted at least once in media not shared	Sharted at least once in media shared	
Step 1	Observed			
	Sharted at least once in media	not shared	0	198
		shared	0	208
Overall Percentage				51.2

a. The cut value is .500

### Variables in the Equation

		B	S.E.	Wald	df	Sig.
Step 1 <sup>a</sup>	propaganda yes or maybe	.113	.215	.276	1	.599
	Constant	.014	.120	.014	1	.905

### Variables in the Equation

		Exp(B)	95% C.I. for EXP(B)	
			Lower	Upper
Step 1 <sup>a</sup>	propaganda yes or maybe	1.119	.735	1.706
	Constant	1.014		

a. Variable(s) entered on step 1: propaganda yes or maybe.

```
LOGISTIC REGRESSION VARIABLESshared_in_media
/METHOD=ENTER PropagandNO
/PRINT=GOODFIT CI(95)
/CRITERIA=PIN(0.05) POUT(0.10) ITERATE(20) CUT(0.5).
```

### Logistic Regression

## Notes

Output Created		10-SEP-2024 21:33:55
Comments		
Input	Data	C: \Users\apalacios\Documents\Media manuscript\Database.sav
	Active Dataset	DataSet1
	Filter	<none>
	Weight	<none>
	Split File	<none>
	N of Rows in Working Data File	407
Missing Value Handling	Definition of Missing	User-defined missing values are treated as missing
Syntax		LOGISTIC REGRESSION VARIABLES shared_in_media /METHOD=ENTER PropagandNO /PRINT=GOODFIT CI(95) /CRITERIA=PIN(0.05) POUT(0.10) ITERATE(20) CUT(0.5).
Resources	Processor Time	00:00:00.00
	Elapsed Time	00:00:00.00

## Case Processing Summary

Unweighted Cases <sup>a</sup>		N	Percent
Selected Cases	Included in Analysis	406	99.8
	Missing Cases	1	.2
	Total	407	100.0
Unselected Cases		0	.0
Total		407	100.0

a. If weight is in effect, see classification table for the total number of cases.

## Dependent Variable Encoding

Original Value	Internal Value
not shared	0
shared	1

## Block 0: Beginning Block

**Classification Table<sup>a,b</sup>**

			Predicted		Percentage Correct
Observed			Sharted at least once in media not shared	shared	
Step 0	Sharted at least once in media	not shared	0	198	.0
		shared	0	208	100.0
	Overall Percentage				51.2

a. Constant is included in the model.

b. The cut value is .500

**Variables in the Equation**

		B	S.E.	Wald	df	Sig.	Exp(B)
Step 0	Constant	.049	.099	.246	1	.620	1.051

**Variables not in the Equation**

			Score	df	Sig.
Step 0	Variables	PropagandNO	.276	1	.599
	Overall Statistics		.276	1	.599

## Block 1: Method = Enter

**Omnibus Tests of Model Coefficients**

		Chi-square	df	Sig.
Step 1	Step	.276	1	.599
	Block	.276	1	.599
	Model	.276	1	.599

**Model Summary**

Step	-2 Log likelihood	Cox & Snell R Square	Nagelkerke R Square
1	562.313 <sup>a</sup>	.001	.001

a. Estimation terminated at iteration number 2 because parameter estimates changed by less than .001.

**Hosmer and Lemeshow Test**

Step	Chi-square	df	Sig.
1	.000	0	.

### Contingency Table for Hosmer and Lemeshow Test

		Sharted at least once in media = not shared		Sharted at least once in media = shared		Total
		Observed	Expected	Observed	Expected	
Step 1	1	139	139.000	141	141.000	280
	2	59	59.000	67	67.000	126

### Classification Table<sup>a</sup>

			Predicted		Percentage Correct
			Sharted at least once in media not shared	shared	
Step 1	Observed				
	Sharted at least once in media	not shared	0	198	.0
		shared	0	208	100.0
	Overall Percentage				51.2

a. The cut value is .500

### Variables in the Equation

		B	S.E.	Wald	df	Sig.	Exp(B)
Step 1 <sup>a</sup>	PropagandNO	-.113	.215	.276	1	.599	.893
	Constant	.127	.179	.507	1	.476	1.136

### Variables in the Equation

		95% C.I. for EXP(B)	
		Lower	Upper
Step 1 <sup>a</sup>	PropagandNO	.586	1.361
	Constant		

a. Variable(s) entered on step 1: PropagandNO.

```
LOGISTIC REGRESSION VARIABLESshared_in_media
/METHOD=ENTER negative
/PRINT=GOODFIT CI(95)
/CRITERIA=PIN(0.05) POUT(0.10) ITERATE(20) CUT(0.5).
```

### Logistic Regression

## Notes

Output Created		10-SEP-2024 21:36:08
Comments		
Input	Data	C:\Users\apalacios\Documents\Media manuscript\Database.sav
	Active Dataset	DataSet1
	Filter	<none>
	Weight	<none>
	Split File	<none>
	N of Rows in Working Data File	407
Missing Value Handling	Definition of Missing	User-defined missing values are treated as missing
Syntax		LOGISTIC REGRESSION VARIABLES shared_in_media /METHOD=ENTER negative /PRINT=GOODFIT CI(95) /CRITERIA=PIN(0.05) POUT(0.10) ITERATE(20) CUT(0.5).
Resources	Processor Time	00:00:00.00
	Elapsed Time	00:00:00.00

## Case Processing Summary

Unweighted Cases <sup>a</sup>		N	Percent
Selected Cases	Included in Analysis	406	99.8
	Missing Cases	1	.2
	Total	407	100.0
Unselected Cases		0	.0
Total		407	100.0

a. If weight is in effect, see classification table for the total number of cases.

## Dependent Variable Encoding

Original Value	Internal Value
not shared	0
shared	1



## Block 0: Beginning Block

**Classification Table<sup>a,b</sup>**

		Predicted		Percentage Correct
Observed		Sharted at least once in media not shared	shared	
Step 0	Sharted at least once in media	not shared	0	.0
		shared	0	100.0
	Overall Percentage			51.2

a. Constant is included in the model.

b. The cut value is .500

**Variables in the Equation**

		B	S.E.	Wald	df	Sig.	Exp(B)
Step 0	Constant	.049	.099	.246	1	.620	1.051

**Variables not in the Equation**

		Score	df	Sig.
Step 0	Variables negative	1.184	1	.277
	Overall Statistics	1.184	1	.277

## Block 1: Method = Enter

**Omnibus Tests of Model Coefficients**

		Chi-square	df	Sig.
Step 1	Step	1.184	1	.277
	Block	1.184	1	.277
	Model	1.184	1	.277

**Model Summary**

Step	-2 Log likelihood	Cox & Snell R Square	Nagelkerke R Square
1	561.405 <sup>a</sup>	.003	.004

a. Estimation terminated at iteration number 2 because parameter estimates changed by less than .001.

**Hosmer and Lemeshow Test**

Step	Chi-square	df	Sig.
1	.000	0	.

### Contingency Table for Hosmer and Lemeshow Test

		Sharted at least once in media = not shared		Sharted at least once in media = shared		Total
		Observed	Expected	Observed	Expected	
Step 1	1	55	55.000	48	48.000	103
	2	143	143.000	160	160.000	303

### Classification Table<sup>a</sup>

			Predicted		Percentage Correct
			Sharted at least once in media not shared	shared	
Step 1	Observed				
	Sharted at least once in media	not shared	55	143	27.8
		shared	48	160	76.9
	Overall Percentage				53.0

a. The cut value is .500

### Variables in the Equation

		B	S.E.	Wald	df	Sig.	Exp(B)	95% C.I. for Lower
Step 1 <sup>a</sup>	negative	-.248	.229	1.181	1	.277	.780	.498
	Constant	.112	.115	.953	1	.329	1.119	

### Variables in the Equation

		95% C.I. for .. Upper
Step 1 <sup>a</sup>	negative	1.221
	Constant	

a. Variable(s) entered on step 1: negative.

```
LOGISTIC REGRESSION VARIABLESshared_in_media
/METHOD=ENTER positive
/PRINT=GOODFIT CI(95)
/CRITERIA=PIN(0.05) POUT(0.10) ITERATE(20) CUT(0.5).
```

### Logistic Regression

## Notes

Output Created		10-SEP-2024 21:36:08
Comments		
Input	Data	C:\Users\apalacios\Documents\Media manuscript\Database.sav
	Active Dataset	DataSet1
	Filter	<none>
	Weight	<none>
	Split File	<none>
	N of Rows in Working Data File	407
Missing Value Handling	Definition of Missing	User-defined missing values are treated as missing
Syntax		LOGISTIC REGRESSION VARIABLES shared_in_media /METHOD=ENTER positive /PRINT=GOODFIT CI(95) /CRITERIA=PIN(0.05) POUT(0.10) ITERATE(20) CUT(0.5).
Resources	Processor Time	00:00:00.02
	Elapsed Time	00:00:00.02

## Case Processing Summary

Unweighted Cases <sup>a</sup>		N	Percent
Selected Cases	Included in Analysis	406	99.8
	Missing Cases	1	.2
	Total	407	100.0
Unselected Cases		0	.0
Total		407	100.0

a. If weight is in effect, see classification table for the total number of cases.

## Dependent Variable Encoding

Original Value	Internal Value
not shared	0
shared	1

## Block 0: Beginning Block

**Classification Table<sup>a,b</sup>**

			Predicted		Percentage Correct
Observed			Sharted at least once in media not shared	shared	
Step 0	Sharted at least once in media	not shared	0	198	.0
		shared	0	208	100.0
	Overall Percentage				51.2

a. Constant is included in the model.

b. The cut value is .500

**Variables in the Equation**

		B	S.E.	Wald	df	Sig.	Exp(B)
Step 0	Constant	.049	.099	.246	1	.620	1.051

**Variables not in the Equation**

			Score	df	Sig.
Step 0	Variables	positive	.797	1	.372
	Overall Statistics		.797	1	.372

## Block 1: Method = Enter

**Omnibus Tests of Model Coefficients**

		Chi-square	df	Sig.
Step 1	Step	.797	1	.372
	Block	.797	1	.372
	Model	.797	1	.372

**Model Summary**

Step	-2 Log likelihood	Cox & Snell R Square	Nagelkerke R Square
1	561.792 <sup>a</sup>	.002	.003

a. Estimation terminated at iteration number 2 because parameter estimates changed by less than .001.

**Hosmer and Lemeshow Test**

Step	Chi-square	df	Sig.
1	.000	0	.

### Contingency Table for Hosmer and Lemeshow Test

		Sharted at least once in media = not shared		Sharted at least once in media = shared		Total
		Observed	Expected	Observed	Expected	
Step 1	1	77	77.000	72	72.000	149
	2	121	121.000	136	136.000	257

### Classification Table<sup>a</sup>

			Predicted		Percentage Correct
			Sharted at least once in media not shared	shared	
Step 1	Observed				
	Sharted at least once in media	not shared	77	121	38.9
		shared	72	136	65.4
	Overall Percentage				52.5

a. The cut value is .500

### Variables in the Equation

		B	S.E.	Wald	df	Sig.	Exp(B)	95% C.I. for Lower
Step 1 <sup>a</sup>	positive	-.184	.206	.797	1	.372	.832	.555
	Constant	.117	.125	.874	1	.350	1.124	

### Variables in the Equation

		95% C.I. for .. Upper
Step 1 <sup>a</sup>	positive	1.246
	Constant	

a. Variable(s) entered on step 1: positive.

```
LOGISTIC REGRESSION VARIABLESshared_in_media
/METHOD=ENTER neutral
/PRINT=GOODFIT CI(95)
/CRITERIA=PIN(0.05) POUT(0.10) ITERATE(20) CUT(0.5).
```

### Logistic Regression

## Notes

Output Created		10-SEP-2024 21:36:08
Comments		
Input	Data	C:\Users\apalacios\Documents\Media manuscript\Database.sav
	Active Dataset	DataSet1
	Filter	<none>
	Weight	<none>
	Split File	<none>
	N of Rows in Working Data File	407
Missing Value Handling	Definition of Missing	User-defined missing values are treated as missing
Syntax		LOGISTIC REGRESSION VARIABLES shared_in_media /METHOD=ENTER neutral /PRINT=GOODFIT CI(95) /CRITERIA=PIN(0.05) POUT(0.10) ITERATE(20) CUT(0.5).
Resources	Processor Time	00:00:00.00
	Elapsed Time	00:00:00.00

## Case Processing Summary

Unweighted Cases <sup>a</sup>		N	Percent
Selected Cases	Included in Analysis	406	99.8
	Missing Cases	1	.2
	Total	407	100.0
Unselected Cases		0	.0
Total		407	100.0

a. If weight is in effect, see classification table for the total number of cases.

## Dependent Variable Encoding

Original Value	Internal Value
not shared	0
shared	1

## Block 0: Beginning Block

**Classification Table<sup>a,b</sup>**

			Predicted		Percentage Correct
Observed			Sharted at least once in media not shared	shared	
Step 0	Sharted at least once in media	not shared	0	198	.0
		shared	0	208	100.0
	Overall Percentage				51.2

a. Constant is included in the model.

b. The cut value is .500

**Variables in the Equation**

		B	S.E.	Wald	df	Sig.	Exp(B)
Step 0	Constant	.049	.099	.246	1	.620	1.051

**Variables not in the Equation**

			Score	df	Sig.
Step 0	Variables	neutral	3.470	1	.062
	Overall Statistics		3.470	1	.062

## Block 1: Method = Enter

**Omnibus Tests of Model Coefficients**

		Chi-square	df	Sig.
Step 1	Step	3.479	1	.062
	Block	3.479	1	.062
	Model	3.479	1	.062

**Model Summary**

Step	-2 Log likelihood	Cox & Snell R Square	Nagelkerke R Square
1	559.110 <sup>a</sup>	.009	.011

a. Estimation terminated at iteration number 3 because parameter estimates changed by less than .001.

**Hosmer and Lemeshow Test**

Step	Chi-square	df	Sig.
1	.000	0	.

### Contingency Table for Hosmer and Lemeshow Test

		Sharted at least once in media = not shared		Sharted at least once in media = shared		Total
		Observed	Expected	Observed	Expected	
Step 1	1	132	132.000	120	120.000	252
	2	66	66.000	88	88.000	154

### Classification Table<sup>a</sup>

			Predicted		Percentage Correct
			Sharted at least once in media not shared	shared	
Step 1	Observed				
	Sharted at least once in media	not shared	132	66	66.7
		shared	120	88	42.3
	Overall Percentage				54.2

a. The cut value is .500

### Variables in the Equation

		B	S.E.	Wald	df	Sig.	Exp(B)	95% C.I. for Lower
Step 1 <sup>a</sup>	neutral	.383	.206	3.458	1	.063	1.467	.980
	Constant	-.095	.126	.571	1	.450	.909	

### Variables in the Equation

		95% C.I. for .. Upper
Step 1 <sup>a</sup>	neutral	2.196
	Constant	

a. Variable(s) entered on step 1: neutral.

FREQUENCIES VARIABLES=Propaganda  
/STATISTICS=RANGE MINIMUMMAXIMUMMODE  
/ORDER=ANALYSIS

### Frequencies



## Notes

Output Created		10-SEP-2024 21:44:05
Comments		
Input	Data	C:\Users\apalacios\Documents\Media manuscript\Database.sav
	Active Dataset	DataSet1
	Filter	<none>
	Weight	<none>
	Split File	<none>
	N of Rows in Working Data File	407
Missing Value Handling	Definition of Missing	User-defined missing values are treated as missing.
	Cases Used	Statistics are based on all cases with valid data.
Syntax		FREQUENCIES VARIABLES=Propaganda /STATISTICS=RANGE MINIMUM MAXIMUM MODE...
Resources	Processor Time	00:00:00.00
	Elapsed Time	00:00:00.00

## Statistics

propaganda yes or maybe

N	Valid	407
	Missing	0
Mode		.00
Range		1.00
Minimum		.00
Maximum		1.00

## propaganda yes or maybe

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No propaganda	280	68.8	68.8	68.8
	Propaganda or maybe propaganda	127	31.2	31.2	100.0
	Total	407	100.0	100.0	