

Lipid-Based Formulations Containing Labrafil M2125-CS: A Deep Investigation on Nanosystem Stability

Martine Tarsitano ¹, Maria Chiara Cristiano ^{2,*}, Antonia Mancuso ², Antonella Barone ², Daniele Torella ² and Donatella Paolino ²

¹ Department of Health Sciences, University of Catanzaro "Magna Graecia", 88100 Catanzaro, Italy; martine.tarsitano@studenti.unicz.it (M.T.)

² Department of Experimental and Clinical Medicine, University "Magna Graecia" of Catanzaro, 88100 Catanzaro, Italy; antonia.mancuso@unicz.it (A.M.); barone@unicz.it (A.B.); dtorella@unicz.it (D.T.); paolino@unicz.it (D.P.)

* Correspondence: mchiara.cristiano@unicz.it (M.C.C.)




Figure S1. Size distribution by intensity (**left**) and raw correlation data (**right**) of samples A, C, and F at t_0 . The name of each panel is related to sample's letter. Data showed are representative of three independent measurements. The plots not visible in the graph but present in the legend were overlaid.




Figure S2. Size distribution by intensity (**left**) and raw correlation data (**right**) of sample F after freeze-drying process with Mannose 5% (panel 1) and Sucrose 5% (panel 2). Data showed are representative of three independent measurements. The plots not visible in the graph but present in legend were overlayed.